



wwPDB EM Validation Summary Report ⓘ

Jun 23, 2026 – 02:56 PM EDT

PDB ID : 10ZZ / pdb_000010zz
EMDB ID : EMD-75578
Title : Yeast co-transcriptional pre-60S assembly intermediate Pwp1 RNP*
Authors : Piwowarczyk, R.; Klinge, S.
Deposited on : 2026-02-13
Resolution : 2.80 Å (reported)
Based on initial model : 8E5T

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY




The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	11806 (2.30 - 3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	3396	 96%
2	G	256	 5% 59% 37%
3	N	204	 7% 63% 32%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	R	576	
5	b	291	
6	i	100	
7	m	427	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12134 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 25S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
1	1	132	2824	1264	514	914	132	0	0

- Molecule 2 is a protein called 60S ribosomal protein L8-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	G	162	1247	800	213	232	2	0	0

- Molecule 3 is a protein called 60S ribosomal protein L15-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	N	138	1187	748	247	191	1	0	0

- Molecule 4 is a protein called Periodic tryptophan protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	R	449	3507	2202	586	702	17	0	0

- Molecule 5 is a protein called Ribosome biogenesis protein BRX1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	b	243	1981	1267	352	356	6	0	0

- Molecule 6 is a protein called 60S ribosomal protein L36-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	i	62	498	310	102	85	1	0	0

- Molecule 7 is a protein called rRNA-processing protein EBP2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	m	107	888	559	154	172	3	0	0


- Molecule 8 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
8	1	2	2	2	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 25S ribosomal RNA



LYS
 LYS
 GLU
 VAL
 ALA
 ALA
 ASP
 THR
 GLU
 TYR
 TYR
 GLN
 SER
 GLN
 ALA
 LEU
 SER
 LYS
 LYS
 GLU
 LYS
 LYS
 ARG
 LYS
 LYS
 LYS
 LEU
 LEU
 MET
 GLN
 GLN
 ASP
 GLU
 THR
 ALA
 ALA
 LYS
 LYS
 HIS
 MET
 SER
 GLY
 ASP
 GLU
 ASP
 GLU
 ASP
 GLU
 SER
 HIS
 MET
 SER
 LYS
 VAL
 VAL
 LYS
 GLU
 ALA
 GLN
 GLU
 ASP
 ASP
 ASP
 GLY
 GLY
 ASP
 ASP
 ASP
 VAL
 VAL
 LEU
 LEU
 VAL
 PHE
 ASP
 GLU
 PHE
 ASP

GLU
 GLU
 GLU
 GLU
 GLY
 ARG
 L128
 D129
 L130
 E131
 K132
 S136
 D137
 S138
 GLU
 SER
 GLU
 ASP
 ASP
 SER
 GLU
 SER
 GLU
 ASN
 ASP
 SER
 GLU
 MET
 GLU
 ASP
 GLU
 ASP
 VAL
 VAL
 LYS
 GLU
 GLN
 GLU
 SER
 GLU
 LYS
 GLU
 GLN
 GLU
 GLN
 VAL
 VAL
 PRO
 LEU
 SER
 VAL
 PHE
 ASP

SER
 ASP
 ALA
 ASP
 VAL
 VAL
 PRO
 HIS
 HIS
 LEU
 LYS
 LEU
 T194
 E219
 D235
 E243
 D286
 K287
 I288
 K289
 GLY
 LYS
 LEU
 LEU
 ILE
 ILE
 ASN
 GLU
 VAL
 VAL
 GLY
 ALA
 SER
 ASP
 LYS
 LYS
 ALA
 ARG
 ARG
 ALA
 ALA
 ARG
 ARG
 GLN
 ARG
 GLN
 ARG
 LYS
 LEU
 LEU
 ASN
 LYS
 LYS
 PHE
 GLY
 GLY
 LYS
 LYS
 VAL
 VAL
 GLN
 GLN
 ASN
 ASN
 ALA
 ALA
 LYS
 THR
 THR
 LEU
 LEU
 GLN
 LYS
 ARG

GLN
 LEU
 GLU
 LYS
 ARG
 ARG
 THR
 LEU
 LEU
 ASN
 ASP
 ALA
 ILE
 ILE
 LYS
 SER
 SER
 LEU
 LYS
 ASN
 ASN
 LYS
 ARG
 LYS
 HIS
 HIS
 ASN
 GLU
 ILE
 ILE
 ASP
 ASP
 HIS
 SER
 LYS
 GLY
 PHE
 LYS
 THR
 ASN
 VAL
 VAL
 GLY
 VAL
 VAL
 GLU
 LYS
 GLU
 VAL
 VAL
 GLY
 LYS
 ARG
 ARG
 PHE
 ASP
 ARG
 GLN
 ARG
 GLN
 ARG
 PRO
 LEU
 ASN
 GLY
 LYS
 ARG
 ARG
 ALA
 ALA
 LYS
 LYS
 VAL
 VAL
 ASN
 ASN
 ALA
 ALA
 LYS
 TYR
 THR
 GLY
 LEU
 GLN
 LYS
 GLY

GLY
 MET
 LYS
 ARG
 PHE
 LYS
 ARG
 LYS
 ASN
 ASP
 ALA
 THR
 SER
 SER
 ALA
 ASP
 VAL
 VAL
 GLY
 PHE
 SER
 SER
 ARG
 ARG
 LYS
 MET
 LYS
 GLY
 LYS
 THR
 ASN
 ARG
 ARG
 PRO
 PRO
 GLY
 LYS
 SER
 SER
 ARG
 ARG
 ALA
 ARG
 ARG
 PHE

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	242741	Depositor
Resolution determination method	OTHER	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	51	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	1700	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	69.787	Depositor
Minimum map value	-38.649	Depositor
Average map value	0.002	Depositor
Map value standard deviation	1.045	Depositor
Recommended contour level	8	Depositor
Map size (\AA)	318.78, 318.78, 318.78	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.0626, 1.0626, 1.0626	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1	0.17	0/3159	0.28	0/4912
2	G	0.17	0/1266	0.34	0/1713
3	N	0.16	0/1212	0.31	0/1626
4	R	0.15	0/3581	0.30	0/4852
5	b	0.18	0/2025	0.37	0/2730
6	i	0.14	0/501	0.33	0/664
7	m	0.15	0/903	0.30	0/1211
All	All	0.16	0/12647	0.31	0/17708

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	2824	0	1423	12	0
2	G	1247	0	1313	7	0
3	N	1187	0	1228	7	0
4	R	3507	0	3354	22	0
5	b	1981	0	2002	10	0
6	i	498	0	544	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	m	888	0	884	2	0
8	1	2	0	0	0	0
All	All	12134	0	10748	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:R:108:SER:O	4:R:111:LYS:NZ	2.16	0.79
5:b:189:ARG:NH1	7:m:243:GLU:OE2	2.20	0.74
5:b:39:ARG:NH2	5:b:66:ASP:OD1	2.20	0.74
4:R:434:ILE:HG21	4:R:437:MET:HE3	1.76	0.67
1:1:269:G:OP1	3:N:47:LYS:NZ	2.21	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	G	160/256 (62%)	159 (99%)	1 (1%)	0	100	100
3	N	134/204 (66%)	129 (96%)	5 (4%)	0	100	100
4	R	437/576 (76%)	432 (99%)	5 (1%)	0	100	100
5	b	237/291 (81%)	235 (99%)	2 (1%)	0	100	100
6	i	60/100 (60%)	60 (100%)	0	0	100	100
7	m	103/427 (24%)	102 (99%)	1 (1%)	0	100	100
All	All	1131/1854 (61%)	1117 (99%)	14 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	G	131/208 (63%)	131 (100%)	0	100	100
3	N	122/176 (69%)	122 (100%)	0	100	100
4	R	388/497 (78%)	388 (100%)	0	100	100
5	b	221/263 (84%)	221 (100%)	0	100	100
6	i	52/82 (63%)	52 (100%)	0	100	100
7	m	100/383 (26%)	100 (100%)	0	100	100
All	All	1014/1609 (63%)	1014 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
3	N	57	GLN
4	R	136	GLN
4	R	390	ASN
5	b	85	ASN
5	b	195	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	1	127/3396 (3%)	8 (6%)	0

5 of 8 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	1	66	A
1	1	109	A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	1	110	G
1	1	121	A
1	1	122	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-75578. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 144



Y Index: 136




Z Index: 159


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 8.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 62 nm³; this corresponds to an approximate mass of 56 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i



*Reported resolution corresponds to spatial frequency of 0.357 \AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-75578 and PDB model 10ZZ. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 8.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (8).

















9.4 Atom inclusion [i](#)



At the recommended contour level, 87% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (8) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8130	 0.5900
1	 0.8870	 0.6020
G	 0.8050	 0.5840
N	 0.8000	 0.5860
R	 0.7510	 0.5790
b	 0.8740	 0.6160
i	 0.7650	 0.5710
m	 0.7400	 0.5650

