# wwPDB EM Validation Summary Report (i)

Nov 19, 2026 – 10:24 AM EST

| | | |
|---|---|---|
| PDB ID | : | 12BB / pdb_000012bb |
| EMDB ID | : | EMD-76274 |
| Title | : | Cryo-EM structure of the iNOS:CaM engaged state |
| Authors | : | Lee, K.; Southworth, D. |
| Deposited on | : | 2026-03-24 |
| Resolution | : | 2.96 Å(reported) |

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

---

The following versions of software and data (see references (i)) were used in the production of this report:

| | | |
|---|---|---|
| EMDB validation analysis | : | 0.0.1.dev133 |
| Mogul | : | 2022.3.0, CSD as543be (2022) |
| MolProbity | : | 4-5-2 with Phenix2.2 |
| Buster-report | : | wwPDB partial adaption of 1.1.7 (2018) |
| Percentile statistics | : | 20250101.v01 (using entries in the PDB archive January 1st 2025) |
| EM percentile statistics | : | 202505.v01 (Using data in the EMDB archive up until May 2025) |
| MapQ | : | 1.9.13 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.51 |

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







| Metric | Whole archive (#Entries) | EM structures (#Entries) | Similar EM resolution (#Entries, resolution range(Å)) |
|---|---|---|---|
| Clashscore | 229148 | 23984 | - |
| Ramachandran outliers | 224038 | 23583 | - |
| Sidechain outliers | 223484 | 23102 | - |
| Q-score | - | 25397 | 13155 ( 2.46 - 3.46 ) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|---|---|---|---|
| 1 | A | 1176 | 88% 11% |
| 1 | B | 1176 | 36% 64% |
| 2 | C | 149 | 13% 95% |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|---|---|---|---|---|---|---|---|
| 6 | FMN | A | 1204 | - | X | - | - |

# 2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 13203 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, inducible.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|---|---|---|---|---|---|---|---|---|---|
| 1 | A | 1043 | Total | C | N | O | S | 0 | 0 |
| | | | 8370 | 5331 | 1474 | 1514 | 51 | | |
| 1 | B | 427 | Total | C | N | O | S | 0 | 0 |
| | | | 3481 | 2223 | 613 | 623 | 22 | | |

There are 48 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|---|---|---|---|---|---|
| A | -22 | MET | - | initiating methionine | UNP P35228 |
| A | -21 | HIS | - | expression tag | UNP P35228 |
| A | -20 | HIS | - | expression tag | UNP P35228 |
| A | -19 | HIS | - | expression tag | UNP P35228 |
| A | -18 | HIS | - | expression tag | UNP P35228 |
| A | -17 | HIS | - | expression tag | UNP P35228 |
| A | -16 | HIS | - | expression tag | UNP P35228 |
| A | -15 | SER | - | expression tag | UNP P35228 |
| A | -14 | SER | - | expression tag | UNP P35228 |
| A | -13 | GLY | - | expression tag | UNP P35228 |
| A | -12 | VAL | - | expression tag | UNP P35228 |
| A | -11 | ASP | - | expression tag | UNP P35228 |
| A | -10 | LEU | - | expression tag | UNP P35228 |
| A | -9 | GLY | - | expression tag | UNP P35228 |
| A | -8 | THR | - | expression tag | UNP P35228 |
| A | -7 | GLU | - | expression tag | UNP P35228 |
| A | -6 | ASN | - | expression tag | UNP P35228 |
| A | -5 | LEU | - | expression tag | UNP P35228 |
| A | -4 | TYR | - | expression tag | UNP P35228 |
| A | -3 | PHE | - | expression tag | UNP P35228 |
| A | -2 | GLN | - | expression tag | UNP P35228 |
| A | -1 | SER | - | expression tag | UNP P35228 |
| A | 0 | ASN | - | expression tag | UNP P35228 |
| A | 1 | ALA | - | expression tag | UNP P35228 |
| B | -22 | MET | - | initiating methionine | UNP P35228 |
| B | -21 | HIS | - | expression tag | UNP P35228 |

*Continued on next page...*

*Continued from previous page...*

| Chain | Residue | Modelled | Actual | Comment | Reference |
|---|---|---|---|---|---|
| B | -20 | HIS | - | expression tag | UNP P35228 |
| B | -19 | HIS | - | expression tag | UNP P35228 |
| B | -18 | HIS | - | expression tag | UNP P35228 |
| B | -17 | HIS | - | expression tag | UNP P35228 |
| B | -16 | HIS | - | expression tag | UNP P35228 |
| B | -15 | SER | - | expression tag | UNP P35228 |
| B | -14 | SER | - | expression tag | UNP P35228 |
| B | -13 | GLY | - | expression tag | UNP P35228 |
| B | -12 | VAL | - | expression tag | UNP P35228 |
| B | -11 | ASP | - | expression tag | UNP P35228 |
| B | -10 | LEU | - | expression tag | UNP P35228 |
| B | -9 | GLY | - | expression tag | UNP P35228 |
| B | -8 | THR | - | expression tag | UNP P35228 |
| B | -7 | GLU | - | expression tag | UNP P35228 |
| B | -6 | ASN | - | expression tag | UNP P35228 |
| B | -5 | LEU | - | expression tag | UNP P35228 |
| B | -4 | TYR | - | expression tag | UNP P35228 |
| B | -3 | PHE | - | expression tag | UNP P35228 |
| B | -2 | GLN | - | expression tag | UNP P35228 |
| B | -1 | SER | - | expression tag | UNP P35228 |
| B | 0 | ASN | - | expression tag | UNP P35228 |
| B | 1 | ALA | - | expression tag | UNP P35228 |

- Molecule 2 is a protein called Calmodulin-1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|---|---|---|---|---|---|---|---|---|---|
| 2 | C | 145 | Total | C | N | O | S | 0 | 0 |
| | | | 1143 | 701 | 184 | 249 | 9 | | |

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

| Mol | Chain | Residues | Atoms | | AltConf |
|---|---|---|---|---|---|
| 3 | A | 1 | Total | Zn | 0 |
| | | | 1 | 1 | |

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | | AltConf |
|---|---|---|---|---|---|---|---|---|
| 4 | A | 1 | Total | C | Fe | N | O | 0 |
| | | | 43 | 34 | 1 | 4 | 4 | |
| 4 | B | 1 | Total | C | Fe | N | O | 0 |
| | | | 43 | 34 | 1 | 4 | 4 | |

- Molecule 5 is 5,6,7,8-TETRAHYDROBIOPTERIN (CCD ID: H4B) (formula: $C_9H_{15}N_5O_3$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | AltConf |
|---|---|---|---|---|---|---|---|
| 5 | A | 1 | Total | C | N | O | 0 |
| | | | 17 | 9 | 5 | 3 | |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Residues | Atoms | | | | AltConf |
|---|---|---|---|---|---|---|---|
| 5 | B | 1 | Total | C | N | O | 0 |
| | | | 17 | 9 | 5 | 3 | |

- Molecule 6 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: $C_{17}H_{21}N_4O_9P$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | | AltConf |
|---|---|---|---|---|---|---|---|---|
| 6 | A | 1 | Total | C | N | O | P | 0 |
| | | | 31 | 17 | 4 | 9 | 1 | |

- Molecule 7 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | | AltConf |
|---|---|---|---|---|---|---|---|---|
| 7 | A | 1 | Total | C | N | O | P | 0 |
| | | | 53 | 27 | 9 | 15 | 2 | |

- Molecule 8 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

| Mol | Chain | Residues | Atoms | | AltConf |
|---|---|---|---|---|---|
| 8 | C | 4 | Total | Ca | 0 |
| | | | 4 | 4 | |

# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nitric oxide synthase, inducible

Chain A:







• Molecule 1: Nitric oxide synthase, inducible

Chain B:



SER CYS GLU ASP GLY GLN GLY LEU ASN TYR LEU PRO GLY GLU HIS LEU GLY VAL CYS PRO GLY ASN GLN PRO ALA LEU VAL GLN GLY ILE LEU GLU ARG VAL VAL ASP GLY PRO THR PRO HIS GLN THR VAL ARG LEU GLU ALA LEU ASP GLU SER GLY SER TYR TRP VAL SER ASP LYS

ARG LEU PRO PRO CYS SER LEU SER GLN ALA LEU THR TYR PHE LEU ASP ILE THR THR PRO PRO THR GLN LEU LEU LEU LEU GLN LYS LEU ALA GLN VAL ALA THR GLU GLU GLU PRO GLU ARG GLN LEU GLU ALA LEU CYS GLN PRO SER GLU TYR SER LYS TRP LYS PHE THR ASN SER PRO

THR PHE LEU GLU VAL LEU GLU GLU PHE PRO SER LEU ARG VAL SER ALA GLY PHE LEU LEU SER GLN LEU PRO ILE LEU LEU LYS PRO ARG PHE TYR SER ILE SER SER SER ARG ASP HIS THR PRO THR GLU ILE HIS LEU THR VAL ALA ALA VAL VAL THR TYR HIS THR ARG ASP GLY GLN GLY

PRO LEU HIS HIS GLY VAL CYS SER THR TRP LEU ASN SER LEU LYS PRO GLN ASP PRO VAL PRO CYS PHE VAL ARG ASN ALA SER GLY PHE HIS LEU PRO GLU ASP PRO SER HIS PRO CYS ILE LEU ILE GLY PRO GLY THR GLY ILE ALA PRO PHE ARG SER PHE TRP GLN GLN ARG LEU

HIS ASP SER GLN HIS LYS GLY VAL ARG GLY GLY ARG MET THR LEU VAL PHE GLY CYS ARG ARG PRO ASP GLU ASP HIS ILE TYR GLN GLU GLU MET LEU GLU MET ALA GLN LYS GLY VAL LEU HIS ALA VAL HIS THR ALA TYR SER ARG LEU PRO GLY LYS PRO LYS VAL TYR VAL GLN

ASP ILE LEU ARG GLN GLN LEU ALA SER GLU VAL LEU ARG VAL LEU HIS LYS GLU PRO GLY HIS LEU TYR VAL CYS GLY ASP VAL ARG MET ALA ARG ASP VAL ALA HIS THR LEU LYS GLN LEU LEU VAL ALA ALA LYS LEU LYS LEU ASN GLU GLU GLN VAL GLU ASP TYR PHE PHE GLN LEU

LYS SER GLN LYS ARG TYR HIS GLU ASP ILE PHE GLY ALA VAL PHE PRO TYR GLU ALA LYS LYS ASP ARG VAL ALA VAL GLN PRO SER SER LEU GLU MET SER ALA LEU

- Molecule 2: Calmodulin-1

Chain C:






# 4 Experimental information (i)

| Property | Value | Source |
|---|---|---|
| EM reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, Not provided | |
| Number of particles used | 56656 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | TFS KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/Å^2$) | 66 | Depositor |
| Minimum defocus (nm) | 1000 | Depositor |
| Maximum defocus (nm) | 2000 | Depositor |
| Magnification | Not provided | |
| Image detector | GATAN K3 BIOQUANTUM (6k x 4k) | Depositor |
| Maximum map value | 1.120 | Depositor |
| Minimum map value | -0.448 | Depositor |
| Average map value | -0.001 | Depositor |
| Map value standard deviation | 0.028 | Depositor |
| Recommended contour level | 0.15 | Depositor |
| Map size (Å) | 283.56, 283.56, 283.56 | wwPDB |
| Map dimensions | 340, 340, 340 | wwPDB |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB |
| Pixel spacing (Å) | 0.834, 0.834, 0.834 | Depositor |

# 5 Model quality

## 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, FAD, H4B, ZN, HEM, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|---|---|---|---|---|---|
| | | RMSZ | $\#\|Z\| > 5$ | RMSZ | $\#\|Z\| > 5$ |
| 1 | A | 0.70 | 0/8590 | 0.96 | 0/11644 |
| 1 | B | 0.72 | 0/3582 | 1.01 | 0/4859 |
| 2 | C | 0.71 | 0/1155 | 0.90 | 0/1551 |
| All | All | 0.71 | 0/13327 | 0.97 | 0/18054 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|---|---|---|---|---|---|---|
| 1 | A | 8370 | 0 | 8273 | 3 | 0 |
| 1 | B | 3481 | 0 | 3390 | 1 | 0 |
| 2 | C | 1143 | 0 | 1071 | 2 | 0 |
| 3 | A | 1 | 0 | 0 | 0 | 0 |
| 4 | A | 43 | 0 | 30 | 0 | 0 |
| 4 | B | 43 | 0 | 30 | 0 | 0 |
| 5 | A | 17 | 0 | 15 | 0 | 0 |
| 5 | B | 17 | 0 | 15 | 0 | 0 |
| 6 | A | 31 | 0 | 18 | 0 | 0 |
| 7 | A | 53 | 0 | 28 | 0 | 0 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|---|---|---|---|---|---|---|
| 8 | C | 4 | 0 | 0 | 0 | 0 |
| All | All | 13203 | 0 | 12870 | 6 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

The worst 5 of 6 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 1:A:1014:PHE:CG | 1:A:1015:GLY:N | 2.75 | 0.53 |
| 2:C:66:PHE:HB3 | 2:C:67:PRO:HD3 | 1.96 | 0.47 |
| 1:A:587:LEU:HD22 | 1:A:622:ALA:HB3 | 1.99 | 0.45 |
| 1:B:194:TRP:CE3 | 1:B:206:TRP:HA | 2.51 | 0.45 |
| 2:C:138:ASN:OD1 | 2:C:138:ASN:C | 2.62 | 0.42 |

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|---|---|---|---|---|---|---|---|
| 1 | A | 1037/1176 (88%) | 1015 (98%) | 22 (2%) | 0 | 100 | 100 |
| 1 | B | 425/1176 (36%) | 417 (98%) | 8 (2%) | 0 | 100 | 100 |
| 2 | C | 143/149 (96%) | 142 (99%) | 1 (1%) | 0 | 100 | 100 |
| All | All | 1605/2501 (64%) | 1574 (98%) | 31 (2%) | 0 | 100 | 100 |

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|---|---|---|---|---|---|
| 1 | A | 915/1033 (89%) | 914 (100%) | 1 (0%) | 88 95 |
| 1 | B | 373/1033 (36%) | 373 (100%) | 0 | 100 100 |
| 2 | C | 124/127 (98%) | 124 (100%) | 0 | 100 100 |
| All | All | 1412/2193 (64%) | 1411 (100%) | 1 (0%) | 87 95 |

All (1) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 796 | THR |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 1001 | GLN |
| 1 | A | 1026 | GLN |
| 2 | C | 50 | GLN |
| 1 | B | 226 | HIS |
| 1 | B | 478 | GLN |

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Counts | RMSZ | $\#\|Z\| > 2$ | Counts | RMSZ | $\#\|Z\| > 2$ |
| 6 | FMN | A | 1204 | - | 33,33,33 | 4.53 | 21 (63%) | 48,50,50 | 2.71 | 21 (43%) |
| 5 | H4B | B | 1202 | - | 17,18,18 | 1.48 | 4 (23%) | 14,26,26 | 1.92 | 2 (14%) |
| 7 | FAD | A | 1205 | - | 58,58,58 | 5.99 | 43 (74%) | 85,89,89 | 3.33 | 31 (36%) |
| 5 | H4B | A | 1203 | - | 17,18,18 | 1.67 | 5 (29%) | 14,26,26 | 1.89 | 3 (21%) |
| 4 | HEM | A | 1202 | 1 | 50,50,50 | 1.50 | 4 (8%) | 67,82,82 | 1.19 | 8 (11%) |
| 4 | HEM | B | 1201 | 1 | 50,50,50 | 1.56 | 7 (14%) | 67,82,82 | 1.32 | 7 (10%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|---|---|---|---|---|---|---|---|
| 6 | FMN | A | 1204 | - | - | 3/18/18/18 | 0/3/3/3 |
| 5 | H4B | B | 1202 | - | - | 0/8/17/17 | 0/2/2/2 |
| 7 | FAD | A | 1205 | - | - | 7/34/50/50 | 0/6/6/6 |
| 5 | H4B | A | 1203 | - | - | 0/8/17/17 | 0/2/2/2 |
| 4 | HEM | A | 1202 | 1 | - | 4/14/54/54 | - |
| 4 | HEM | B | 1201 | 1 | - | 4/14/54/54 | - |

The worst 5 of 84 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|---|---|---|---|---|---|---|---|
| 7 | A | 1205 | FAD | P-O3P | 17.74 | 1.78 | 1.59 |
| 7 | A | 1205 | FAD | PA-O3P | -12.75 | 1.45 | 1.59 |
| 6 | A | 1204 | FMN | C9-C9A | 12.26 | 1.59 | 1.39 |
| 7 | A | 1205 | FAD | C9A-C5X | 11.86 | 1.60 | 1.41 |
| 7 | A | 1205 | FAD | C2A-N1A | 11.85 | 1.55 | 1.33 |

The worst 5 of 72 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|---|---|---|---|---|---|---|---|
| 7 | A | 1205 | FAD | O4B-C1B-N9A | -15.20 | 78.90 | 108.09 |
| 7 | A | 1205 | FAD | O3P-PA-O1A | 11.01 | 143.83 | 110.70 |
| 7 | A | 1205 | FAD | C4A-N9A-C8A | -9.01 | 96.27 | 105.74 |
| 7 | A | 1205 | FAD | C2B-C1B-N9A | 8.00 | 133.19 | 113.30 |
| 7 | A | 1205 | FAD | C5X-C9A-N10 | -7.35 | 111.32 | 117.97 |

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|---|---|---|---|---|
| 4 | A | 1202 | HEM | C2B-C3B-CAB-CBB |
| 4 | A | 1202 | HEM | C4B-C3B-CAB-CBB |
| 4 | B | 1201 | HEM | C2B-C3B-CAB-CBB |
| 7 | A | 1205 | FAD | O4B-C4B-C5B-O5B |
| 7 | A | 1205 | FAD | C3B-C4B-C5B-O5B |

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

**Ligand FMN A 1204**



**Bond lengths**



**Bond angles**



**Torsions**



**Rings**

## Ligand H4B B 1202



Bond lengths



Bond angles



Torsions



Rings

## Ligand FAD A 1205



Bond lengths



Bond angles



Torsions



Rings

**Ligand H4B A 1203**



**Bond lengths**



**Bond angles**



**Torsions**



**Rings**

**Ligand HEM A 1202**



**Bond lengths**



**Bond angles**



**Torsions**



**Rings**

**Ligand HEM B 1201**



**Bond lengths**



**Bond angles**



**Torsions**



**Rings**

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

# 6 Map visualisation

This section contains visualisations of the EMDB entry EMD-76274. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

## 6.1 Orthogonal projections

### 6.1.1 Primary map



X



Y



Z

### 6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

## 6.2 Central slices (i)

### 6.2.1 Primary map



X Index: 170



Y Index: 170



Z Index: 170

### 6.2.2 Raw map



X Index: 170



Y Index: 170



Z Index: 170

The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices (i)

### 6.3.1 Primary map



X Index: 180



Y Index: 158



Z Index: 176

### 6.3.2 Raw map



X Index: 180



Y Index: 158



Z Index: 176

The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) (i)

### 6.4.1 Primary map



X



Y



Z

### 6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views (i)

### 6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.15. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

## 6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

# 7 Map analysis

This section contains the results of statistical analysis of the map.

## 7.1 Map-value distribution




The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate (i)




The volume at the recommended contour level is 133 $nm^3$; this corresponds to an approximate mass of 120 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

## 7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.338 Å$^{-1}$

# 8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

## 8.1 FSC




*Reported resolution corresponds to spatial frequency of 0.338 Å$^{-1}$

## 8.2 Resolution estimates (i)

| Resolution estimate (Å) | Estimation criterion (FSC cut-off) | | |
|---|---|---|---|
| | 0.143 | 0.5 | Half-bit |
| Reported by author | 2.96 | - | - |
| Author-provided FSC curve | - | - | - |
| Unmasked-calculated* | 3.57 | 6.18 | 3.70 |

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.57 differs from the reported value 2.96 by more than 10 %

# 9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-76274 and PDB model 12BB. Per-residue inclusion information can be found in section 3 on page 9.

## 9.1 Map-model overlay



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.15 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model (i)




X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model (i)




X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.15).

## 9.4 Atom inclusion (i)




At the recommended contour level, 96% of all backbone atoms, 86% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.15) and Q-score for the entire model and for each chain.




| Chain | Atom inclusion | Q-score |
|---|---|---|
| All | 0.8610 | 0.3990 |
| A | 0.8500 | 0.3720 |
| B | 0.9410 | 0.5240 |
| C | 0.6960 | 0.2160 |