



wwPDB EM Validation Summary Report ⓘ

Jul 6, 2026 – 02:29 PM EDT

PDB ID : 36IS / pdb_000036is
EMDB ID : EMD-77603
Title : Salmonella Flagellar Export Apparatus (FliPQR/FlhB/FlhA) in the context of the intact basal body
Authors : Johnson, S.; Johnson, M.K.; Lea, S.M.
Deposited on : 2026-06-11
Resolution : 4.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	4585 (3.80 - 4.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A1	245	71% 14% 15%
1	B1	245	71% 14% 15%
1	C1	245	69% 16% 15%
1	D1	245	74% 11% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E1	245	
2	F1	264	
3	G1	89	
3	H1	89	
3	I1	89	
3	J1	89	
4	K1	383	
5	A2	692	
5	B2	692	
5	C2	692	
5	D2	692	
5	E2	692	
5	F2	692	
5	G2	692	
5	H2	692	
5	I2	692	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 35138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flagellar biosynthetic protein FliP.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A1	209	Total	C	N	O	S	0	0
			1623	1084	251	276	12		
1	B1	209	Total	C	N	O	S	0	0
			1623	1084	251	276	12		
1	C1	209	Total	C	N	O	S	0	0
			1623	1084	251	276	12		
1	D1	209	Total	C	N	O	S	0	0
			1623	1084	251	276	12		
1	E1	209	Total	C	N	O	S	0	0
			1623	1084	251	276	12		

- Molecule 2 is a protein called Flagellar biosynthetic protein FliR.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	F1	250	Total	C	N	O	S	0	0
			1925	1293	304	313	15		

- Molecule 3 is a protein called Flagellar biosynthetic protein FliQ.


Mol	Chain	Residues	Atoms					AltConf	Trace
3	G1	89	Total	C	N	O	S	0	0
			670	449	100	114	7		
3	H1	89	Total	C	N	O	S	0	0
			670	449	100	114	7		
3	I1	89	Total	C	N	O	S	0	0
			670	449	100	114	7		
3	J1	89	Total	C	N	O	S	0	0
			670	449	100	114	7		

- Molecule 4 is a protein called Flagellar biosynthetic protein FlhB.



Mol	Chain	Residues	Atoms					AltConf	Trace
4	K1	214	Total	C	N	O	S	0	0
			1637	1076	278	268	15		

- Molecule 5 is a protein called Flagellar biosynthesis protein FlhA.


Mol	Chain	Residues	Atoms					AltConf	Trace
5	A2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		
5	B2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		
5	C2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		
5	D2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		
5	E2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		
5	F2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		
5	G2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		
5	H2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		
5	I2	311	Total	C	N	O	S	0	0
			2309	1516	377	401	15		



- Molecule 1: Flagellar biosynthetic protein FliP




- Molecule 2: Flagellar biosynthetic protein FliR





- Molecule 3: Flagellar biosynthetic protein FliQ




- Molecule 3: Flagellar biosynthetic protein FliQ




- Molecule 3: Flagellar biosynthetic protein FliQ




- Molecule 3: Flagellar biosynthetic protein FliQ








• Molecule 4: Flagellar biosynthetic protein FlhB




• Molecule 5: Flagellar biosynthesis protein FlhA




● Molecule 5: Flagellar biosynthesis protein FlhA




● Molecule 5: Flagellar biosynthesis protein FlhA








Chain G2:



Chain H2:



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	58082	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	58.5	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	TFS FALCON 4i (4k x 4k)	Depositor
Maximum map value	12.235	Depositor
Minimum map value	-0.113	Depositor
Average map value	0.018	Depositor
Map value standard deviation	0.375	Depositor
Recommended contour level	2	Depositor
Map size (Å)	537.6, 537.6, 537.6	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.05, 1.05, 1.05	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A1	0.12	0/1662	0.32	0/2263
1	B1	0.14	0/1662	0.38	0/2263
1	C1	0.14	0/1662	0.30	0/2263
1	D1	0.14	0/1662	0.32	0/2263
1	E1	0.15	0/1662	0.37	0/2263
2	F1	0.14	0/1974	0.30	0/2696
3	G1	0.11	0/681	0.26	0/930
3	H1	0.12	0/681	0.30	0/930
3	I1	0.11	0/681	0.27	0/930
3	J1	0.12	0/681	0.29	0/930
4	K1	0.11	0/1671	0.26	0/2260
5	A2	0.14	0/2346	0.36	0/3187
5	B2	0.14	0/2346	0.38	0/3187
5	C2	0.14	0/2346	0.37	0/3187
5	D2	0.14	0/2346	0.38	0/3187
5	E2	0.14	0/2346	0.40	0/3187
5	F2	0.12	0/2346	0.37	0/3187
5	G2	0.14	0/2346	0.38	0/3187
5	H2	0.14	0/2346	0.36	0/3187
5	I2	0.14	0/2346	0.37	0/3187
All	All	0.13	0/35793	0.35	0/48674

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
5	D2	0	1
5	F2	0	1
5	G2	0	1
5	I2	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
5	D2	125	ILE	Peptide
5	F2	125	ILE	Peptide
5	G2	125	ILE	Peptide
5	I2	125	ILE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A1	1623	0	1715	26	0
1	B1	1623	0	1715	28	0
1	C1	1623	0	1715	28	0
1	D1	1623	0	1715	22	0
1	E1	1623	0	1715	24	0
2	F1	1925	0	2042	29	0
3	G1	670	0	739	9	0
3	H1	670	0	739	17	0
3	I1	670	0	739	15	0
3	J1	670	0	739	14	0
4	K1	1637	0	1761	23	0
5	A2	2309	0	2450	65	0
5	B2	2309	0	2450	53	0
5	C2	2309	0	2450	54	0
5	D2	2309	0	2450	70	0
5	E2	2309	0	2450	65	0
5	F2	2309	0	2450	76	0
5	G2	2309	0	2450	61	0
5	H2	2309	0	2450	58	0
5	I2	2309	0	2450	60	0
All	All	35138	0	37384	667	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 667 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D2:235:ILE:O	5:D2:239:ALA:HB3	1.72	0.89
5:D2:86:LEU:HD13	5:E2:136:ILE:HD11	1.61	0.82
5:F2:34:MET:HA	5:F2:39:LEU:HD11	1.62	0.82
5:F2:96:ILE:HD11	5:F2:245:LEU:HB3	1.63	0.79
5:I2:85:ARG:HH22	5:I2:260:ILE:HD11	1.47	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A1	207/245 (84%)	199 (96%)	7 (3%)	1 (0%)	24	62
1	B1	207/245 (84%)	200 (97%)	7 (3%)	0	100	100
1	C1	207/245 (84%)	200 (97%)	7 (3%)	0	100	100
1	D1	207/245 (84%)	202 (98%)	5 (2%)	0	100	100
1	E1	207/245 (84%)	196 (95%)	11 (5%)	0	100	100
2	F1	248/264 (94%)	237 (96%)	10 (4%)	1 (0%)	30	66
3	G1	87/89 (98%)	87 (100%)	0	0	100	100
3	H1	87/89 (98%)	87 (100%)	0	0	100	100
3	I1	87/89 (98%)	87 (100%)	0	0	100	100
3	J1	87/89 (98%)	87 (100%)	0	0	100	100
4	K1	212/383 (55%)	210 (99%)	2 (1%)	0	100	100
5	A2	309/692 (45%)	288 (93%)	21 (7%)	0	100	100
5	B2	309/692 (45%)	287 (93%)	22 (7%)	0	100	100
5	C2	309/692 (45%)	286 (93%)	23 (7%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	D2	309/692 (45%)	288 (93%)	21 (7%)	0	100	100
5	E2	309/692 (45%)	287 (93%)	22 (7%)	0	100	100
5	F2	309/692 (45%)	289 (94%)	20 (6%)	0	100	100
5	G2	309/692 (45%)	286 (93%)	23 (7%)	0	100	100
5	H2	309/692 (45%)	288 (93%)	21 (7%)	0	100	100
5	I2	309/692 (45%)	286 (93%)	23 (7%)	0	100	100
All	All	4624/8456 (55%)	4377 (95%)	245 (5%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A1	167	MET
2	F1	164	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A1	179/204 (88%)	179 (100%)	0	100	100
1	B1	179/204 (88%)	179 (100%)	0	100	100
1	C1	179/204 (88%)	179 (100%)	0	100	100
1	D1	179/204 (88%)	179 (100%)	0	100	100
1	E1	179/204 (88%)	179 (100%)	0	100	100
2	F1	207/221 (94%)	207 (100%)	0	100	100
3	G1	74/74 (100%)	73 (99%)	1 (1%)	59	71
3	H1	74/74 (100%)	74 (100%)	0	100	100
3	I1	74/74 (100%)	74 (100%)	0	100	100
3	J1	74/74 (100%)	74 (100%)	0	100	100
4	K1	178/316 (56%)	178 (100%)	0	100	100
5	A2	241/561 (43%)	240 (100%)	1 (0%)	84	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	B2	241/561 (43%)	241 (100%)	0	100	100
5	C2	241/561 (43%)	241 (100%)	0	100	100
5	D2	241/561 (43%)	240 (100%)	1 (0%)	84	82
5	E2	241/561 (43%)	241 (100%)	0	100	100
5	F2	241/561 (43%)	241 (100%)	0	100	100
5	G2	241/561 (43%)	241 (100%)	0	100	100
5	H2	241/561 (43%)	241 (100%)	0	100	100
5	I2	241/561 (43%)	241 (100%)	0	100	100
All	All	3745/6902 (54%)	3742 (100%)	3 (0%)	87	88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G1	63	ILE
5	A2	210	ILE
5	D2	96	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
5	B2	172	ASN
5	D2	288	ASN
5	B2	230	GLN
5	D2	137	ASN
5	F2	137	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-77603. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256




Y Index: 256




Z Index: 256


6.2.2 Raw map



X Index: 256



Y Index: 256



Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 242




Y Index: 246




Z Index: 198


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) ⓘ


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 2.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1102 nm^3 ; this corresponds to an approximate mass of 995 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.233 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.233 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.30	-	-
Author-provided FSC curve	4.30	6.34	4.38
Unmasked-calculated*	8.42	18.05	8.62

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.42 differs from the reported value 4.3 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-77603 and PDB model 36IS. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 2.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2).

9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8750	 0.3210
A1	 0.9270	 0.4230
A2	 0.8270	 0.2090
B1	 0.9410	 0.4350
B2	 0.8100	 0.2030
C1	 0.9390	 0.4390
C2	 0.8050	 0.2070
D1	 0.9360	 0.4530
D2	 0.8410	 0.2580
E1	 0.9410	 0.4360
E2	 0.8500	 0.2770
F1	 0.9380	 0.4420
F2	 0.8590	 0.2850
G1	 0.9350	 0.4590
G2	 0.8710	 0.2750
H1	 0.9220	 0.4640
H2	 0.8430	 0.2450
I1	 0.9220	 0.4760
I2	 0.8270	 0.2060
J1	 0.9340	 0.4540
K1	 0.8870	 0.3950

