



wwPDB EM Validation Summary Report ⓘ

Jun 25, 2025 – 06:58 am BST

PDB ID : 7ARB / pdb_00007arb
EMDB ID : EMD-11878
Title : Cryo-EM structure of Arabidopsis thaliana Complex-I (complete composition)
Authors : Klusch, N.; Kuelbrandt, W.; Yildiz, O.
Deposited on : 2020-10-23
Resolution : 3.41 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0rc1
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	119	10% 57% 20% 23%
2	B	218	55% 17% 28%
3	C	190	74% 23% .
4	D	394	76% 21% .
5	E	255	10% 58% 16% 25%
6	F	486	5% 69% 20% 11%
7	G	748	5% 75% 17% 8%
8	H	325	6% 70% 26% .








Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	I	222	
10	J	205	
11	K	100	
12	L	669	
13	M	495	
14	N	499	
15	O	159	
16	P	402	
17	Q	154	
18	R	110	
19	S	97	
20	T	122	
21	U	126	
22	V	169	
23	W	133	
24	X	106	
25	Z	143	
26	a	65	
27	b	65	
28	c	88	
29	d	81	
30	e	83	
31	f	106	
32	g	114	
33	i	98	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
34	j	69	
35	k	72	
36	l	125	
37	m	71	
38	n	117	
39	o	103	
40	p	106	
41	q	159	
42	r	131	
43	u	30	
44	v	113	
45	x	256	
46	y	278	
47	z	275	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
48	SF4	F	501	-	-	X	-
50	FES	E	500	-	-	X	-

2 Entry composition

There are 61 unique types of molecules in this entry. The entry contains 61765 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-ubiquinone oxidoreductase chain 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	92	Total	C	N	O	S	0	0
			779	553	105	117	4		

- Molecule 2 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 7, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	157	Total	C	N	O	S	0	0
			1244	797	218	215	14		

- Molecule 3 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	185	Total	C	N	O	S	0	0
			1581	1021	271	283	6		

- Molecule 4 is a protein called NADH dehydrogenase subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	385	Total	C	N	O	S	0	0
			3077	1954	542	557	24		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	363	SER	LEU	variant	UNP A0A2P2CLH2

- Molecule 5 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	192	Total	C	N	O	S	0	0
			1500	954	248	287	11		

- Molecule 6 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	434	Total	C	N	O	S	0	0
			3368	2125	600	618	25		

- Molecule 7 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	688	Total	C	N	O	S	0	0
			5252	3291	921	1001	39		

- Molecule 8 is a protein called NADH-ubiquinone oxidoreductase chain 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	311	Total	C	N	O	S	0	0
			2439	1659	372	393	15		

- Molecule 9 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 8-A, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	165	Total	C	N	O	S	0	0
			1349	849	229	261	10		

- Molecule 10 is a protein called NADH-ubiquinone oxidoreductase chain 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	147	Total	C	N	O	S	0	0
			1162	784	179	191	8		

- Molecule 11 is a protein called NADH dehydrogenase subunit 4L.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	90	Total	C	N	O	S	0	0
			707	476	109	115	7		

- Molecule 12 is a protein called NADH-ubiquinone oxidoreductase chain 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L	615	Total	C	N	O	S	0	0
			4807	3191	748	832	36		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	91	PHE	SER	conflict	UNP B5TM94

- Molecule 13 is a protein called NADH-ubiquinone oxidoreductase chain 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	M	487	Total	C	N	O	S	0	0
			3887	2627	601	636	23		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	326	LEU	PRO	conflict	UNP B5TM93

- Molecule 14 is a protein called NADH-ubiquinone oxidoreductase chain 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	488	Total	C	N	O	S	0	0
			3820	2573	577	642	28		

- Molecule 15 is a protein called AT3G07480.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	122	Total	C	N	O	S	0	0
			956	598	169	185	4		

- Molecule 16 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 9, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	P	332	Total	C	N	O	S	0	0
			2560	1643	439	463	15		

- Molecule 17 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Q	119	Total	C	N	O	S	0	0
			939	600	163	175	1		

- Molecule 18 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 6, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	R	62	Total	C	N	O	S	0	0
			482	304	84	89	5		

- Molecule 19 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	S	93	Total	C	N	O	S	0	0
			727	459	129	133	6		

- Molecule 20 is a protein called Acyl carrier protein 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	T	84	Total	C	N	O	S	0	0
			667	421	105	138	3		

- Molecule 21 is a protein called Acyl carrier protein 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	U	83	Total	C	N	O	S	0	0
			650	411	103	135	1		

- Molecule 22 is a protein called Probable NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 5, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	V	140	Total	C	N	O	S	0	0
			1123	712	187	219	5		

- Molecule 23 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	W	112	Total	C	N	O	S	0	0
			904	578	161	162	3		

- Molecule 24 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8-B.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	X	97	Total	C	N	O	S	0	0
			767	480	132	143	12		

- Molecule 25 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	Z	125	Total	C	N	O	S	0	0
			997	640	175	177	5		

- Molecule 26 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	a	58	Total	C	N	O	S	0	0
			469	302	84	78	5		

- Molecule 27 is a protein called At2g46540/F11C10.23.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	b	40	Total	C	N	O	S	0	0
			295	195	48	49	3		

- Molecule 28 is a protein called Transmembrane protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	c	76	Total	C	N	O	S	0	0
			617	396	115	100	6		

- Molecule 29 is a protein called Excitatory amino acid transporter.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	d	75	Total	C	N	O	S	0	0
			592	382	106	99	5		

- Molecule 30 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 5-B.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	e	65	Total	C	N	O	S	0	0
			557	344	106	100	7		

- Molecule 31 is a protein called At4g16450.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	f	101	Total	C	N	O	S	0	0
			763	490	126	142	5		

- Molecule 32 is a protein called ESSS subunit of NADH:ubiquinone oxidoreductase (Complex I) protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	g	75	Total	C	N	O	S	0	0
			615	396	107	109	3		

- Molecule 33 is a protein called P1.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	i	85	Total	C	N	O	S	0	0
			737	466	135	131	5		

- Molecule 34 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	j	51	Total	C	N	O	S	0	0
			415	275	73	64	3		

- Molecule 35 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	k	47	Total	C	N	O	S	0	0
			374	238	71	62	3		

- Molecule 36 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
36	l	66	Total	C	N	O	S	0	0
			502	328	80	93	1		

- Molecule 37 is a protein called B15 – 1 beta subcomplex subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
37	m	70	Total	C	N	O	S	0	0
			577	370	107	98	2		

- Molecule 38 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
38	n	109	Total	C	N	O	S	0	0
			911	580	170	160	1		

- Molecule 39 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
39	o	80	Total	C	N	O	S	0	0
			657	413	115	119	10		

- Molecule 40 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10-B.

Mol	Chain	Residues	Atoms					AltConf	Trace
40	p	93	Total	C	N	O	S	0	0
			778	493	144	137	4		

- Molecule 41 is a protein called Probable NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 12.

Mol	Chain	Residues	Atoms				AltConf	Trace
41	q	65	Total	C	N	O	0	0
			536	342	95	99		

- Molecule 42 is a protein called Furry.

Mol	Chain	Residues	Atoms				AltConf	Trace
42	r	10	Total	C	N	O	0	0
			89	56	18	15		

- Molecule 43 is a protein called unknown.

Mol	Chain	Residues	Atoms				AltConf	Trace
43	u	30	Total	C	N	O	0	0
			150	90	30	30		

- Molecule 44 is a protein called Uncharacterized protein At2g27730, mitochondrial.

Mol	Chain	Residues	Atoms				AltConf	Trace
44	v	30	Total	C	N	O	0	0
			226	147	39	40		

- Molecule 45 is a protein called Gamma carbonic anhydrase-like 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
45	x	214	Total	C	N	O	S	0	0
			1659	1063	285	306	5		


- Molecule 46 is a protein called Gamma carbonic anhydrase 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
46	y	268	Total	C	N	O	S	0	0
			2032	1271	363	391	7		

- Molecule 47 is a protein called Gamma carbonic anhydrase 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
47	z	233	Total	C	N	O	S	0	0
			1772	1111	325	330	6		

- Molecule 48 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			AltConf
48	B	1	Total	Fe	S	0
			8	4	4	
48	F	1	Total	Fe	S	0
			8	4	4	
48	G	1	Total	Fe	S	0
			8	4	4	
48	G	1	Total	Fe	S	0
			8	4	4	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			AltConf
48	I	1	Total	Fe	S	0
			8	4	4	
48	I	1	Total	Fe	S	0
			8	4	4	

- Molecule 49 is PHOSPHATIDYLETHANOLAMINE (CCD ID: PTY) (formula: C₄₀H₈₀NO₈P).




Mol	Chain	Residues	Atoms					AltConf
49	D	1	Total	C	N	O	P	0
			50	40	1	8	1	
49	M	1	Total	C	N	O	P	0
			50	40	1	8	1	
49	N	1	Total	C	N	O	P	0
			50	40	1	8	1	

- Molecule 50 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			AltConf
50	E	1	Total	Fe	S	0
			4	2	2	
50	G	1	Total	Fe	S	0
			4	2	2	

- Molecule 51 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: C₁₇H₂₁N₄O₉P).




Mol	Chain	Residues	Atoms					AltConf
51	F	1	Total	C	N	O	P	0
			31	17	4	9	1	

- Molecule 52 is Ubiquinone-9 (CCD ID: UQ9) (formula: C₅₄H₈₂O₄).



Mol	Chain	Residues	Atoms			AltConf
52	H	1	Total	C	O	0
			35	31	4	

- Molecule 53 is Lauryl Maltose Neopentyl Glycol (CCD ID: LMN) (formula: $C_{47}H_{88}O_{22}$).




Mol	Chain	Residues	Atoms			AltConf
53	M	1	Total	C	O	0
			69	47	22	

- Molecule 54 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		AltConf
54	O	1	Total	Fe	0
			1	1	

- Molecule 55 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					AltConf
55	P	1	Total	C	N	O	P	0
			48	21	7	17	3	

- Molecule 56 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
56	R	1	Total	Zn	0
			1	1	
56	y	1	Total	Zn	0
			1	1	

- Molecule 57 is S-[2-({N-[(2R)-2-hydroxy-3,3-dimethyl-4-(phosphonooxy)butanoyl]-beta-alanyl}amino)ethyl] dodecanethioate (CCD ID: 8Q1) (formula: $C_{23}H_{45}N_2O_8PS$).



- Molecule 58 is (7S)-4-HYDROXY-N,N,N-TRIMETHYL-9-OXO-7-[(PALMITOYLOXY) METHYL]-3,5,8-TRIOXA-4-PHOSPHAHEXACOSAN-1-AMINIUM 4-OXIDE (CCD ID: PC7) (formula: $C_{42}H_{85}NO_8P$).




Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
58	v	1	52	42	1	8	1	0

- Molecule 59 is (1S)-2-{{[(2R)-2,3-DIHYDROXYPROPYL]OXY}(HYDROXY)PHOSPHORYL]OXY}-1-[(PALMITOYLOXY)METHYL]ETHYL STEARATE (CCD ID: PGT) (formula: C₄₀H₇₉O₁₀P).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	O	P	
59	f	1	41	30	10	1	0

- Molecule 60 is Phosphatidylinositol (CCD ID: T7X) (formula: C₄₇H₈₃O₁₃P).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	O	P	
60	f	1	61	47	13	1	0


- Molecule 61 is 1,2-DICAPROYL-SN-PHOSPHATIDYL-L-SERINE (CCD ID: PSF) (formula: C₁₈H₃₄NO₁₀P).




3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NADH-ubiquinone oxidoreductase chain 3




- Molecule 2: NADH dehydrogenase [ubiquinone] iron-sulfur protein 7, mitochondrial




- Molecule 3: NADH dehydrogenase [ubiquinone] iron-sulfur protein 3




- Molecule 4: NADH dehydrogenase subunit 7







- Molecule 5: NADH dehydrogenase [ubiquinone] flavoprotein 2, mitochondrial




- Molecule 6: NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial




- Molecule 7: NADH dehydrogenase [ubiquinone] iron-sulfur protein 1, mitochondrial




• Molecule 8: NADH-ubiquinone oxidoreductase chain 1




• Molecule 9: NADH dehydrogenase [ubiquinone] iron-sulfur protein 8-A, mitochondrial







• Molecule 10: NADH-ubiquinone oxidoreductase chain 6



• Molecule 11: NADH dehydrogenase subunit 4L




• Molecule 12: NADH-ubiquinone oxidoreductase chain 5




• Molecule 13: NADH-ubiquinone oxidoreductase chain 4

Chain M: 




• Molecule 14: NADH-ubiquinone oxidoreductase chain 2

Chain N: 




• Molecule 15: AT3G07480.1

Chain O: 




• Molecule 16: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 9, mitochondrial



Chain P: 



- Molecule 17: NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial



- Molecule 18: NADH dehydrogenase [ubiquinone] iron-sulfur protein 6, mitochondrial




- Molecule 19: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 2




- Molecule 20: Acyl carrier protein 1, mitochondrial







- Molecule 21: Acyl carrier protein 2, mitochondrial




- Molecule 22: Probable NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 5, mitochondrial




- Molecule 23: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6





- Molecule 24: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8-B






- Molecule 25: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13-A



- Molecule 26: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1




- Molecule 27: At2g46540/F11C10.23



- Molecule 28: Transmembrane protein




- Molecule 29: Excitatory amino acid transporter





- Molecule 30: NADH dehydrogenase [ubiquinone] iron-sulfur protein 5-B








- Molecule 31: At4g16450



- Molecule 32: ESSS subunit of NADH:ubiquinone oxidoreductase (Complex I) protein



- Molecule 33: P1




- Molecule 34: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 2




- Molecule 35: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3-A





- Molecule 36: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial






- Molecule 37: B15 – 1 beta subcomplex subunit 4




- Molecule 38: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 9




- Molecule 39: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 7




- Molecule 40: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10-B




- Molecule 41: Probable NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 12




- Molecule 42: Furry




- Molecule 43: unknown




- Molecule 44: Uncharacterized protein At2g27730, mitochondrial




- Molecule 45: Gamma carbonic anhydrase-like 2, mitochondrial




- Molecule 46: Gamma carbonic anhydrase 2, mitochondrial





• Molecule 47: Gamma carbonic anhydrase 1, mitochondrial



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	459177	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	43	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.083	Depositor
Minimum map value	-0.043	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.011	Depositor
Map size (Å)	502.2, 502.2, 502.2	wwPDB
Map dimensions	600, 600, 600	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.837, 0.837, 0.837	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, NDP, PGT, LMN, ZN, 8Q1, FES, PC7, T7X, PSF, PTY, UQ9, FE, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/807	0.28	0/1096
2	B	0.18	0/1279	0.33	0/1734
3	C	0.15	0/1629	0.29	0/2207
4	D	0.17	0/3147	0.32	0/4256
5	E	0.14	0/1535	0.35	0/2084
6	F	0.14	0/3441	0.31	1/4641 (0.0%)
7	G	0.15	0/5347	0.29	0/7242
8	H	0.19	0/2510	0.37	0/3416
9	I	0.16	0/1378	0.29	0/1862
10	J	0.15	0/1187	0.30	0/1617
11	K	0.16	0/717	0.30	0/969
12	L	0.15	0/4938	0.32	0/6706
13	M	0.14	0/3995	0.27	0/5428
14	N	0.15	0/3924	0.31	0/5327
15	O	0.11	0/971	0.26	0/1314
16	P	0.14	0/2617	0.34	1/3544 (0.0%)
17	Q	0.12	0/966	0.24	0/1305
18	R	0.12	0/493	0.25	0/668
19	S	0.16	0/739	0.32	0/996
20	T	0.14	0/679	0.37	0/922
21	U	0.11	0/660	0.29	0/892
22	V	0.15	0/1146	0.39	0/1555
23	W	0.12	0/923	0.27	0/1249
24	X	0.12	0/781	0.27	0/1049
25	Z	0.16	0/1027	0.36	0/1392
26	a	0.15	0/481	0.35	0/646
27	b	0.13	0/300	0.25	0/407
28	c	0.12	0/637	0.25	0/860
29	d	0.15	0/605	0.28	0/815
30	e	0.13	0/570	0.20	0/759
31	f	0.14	0/779	0.26	0/1052
32	g	0.12	0/635	0.28	0/863

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
33	i	0.13	0/757	0.26	0/1019
34	j	0.10	0/433	0.20	0/592
35	k	0.09	0/384	0.23	0/515
36	l	0.12	0/513	0.26	0/696
37	m	0.13	0/592	0.28	0/793
38	n	0.10	0/938	0.28	0/1273
39	o	0.14	0/666	0.30	0/886
40	p	0.14	0/799	0.31	0/1074
41	q	0.11	0/553	0.32	0/750
42	r	0.11	0/89	0.35	0/118
44	v	0.10	0/230	0.20	0/311
45	x	0.14	0/1700	0.29	0/2320
46	y	0.13	0/2066	0.26	0/2800
47	z	0.13	0/1804	0.23	0/2441
All	All	0.15	0/62367	0.30	2/84461 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	368	VAL	N-CA-C	-6.39	106.22	111.91
16	P	338	ILE	N-CA-C	-5.59	108.40	113.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	779	0	796	23	0
2	B	1244	0	1231	36	0
3	C	1581	0	1529	36	0
4	D	3077	0	3043	61	0
5	E	1500	0	1473	42	0
6	F	3368	0	3352	65	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	G	5252	0	5241	79	0
8	H	2439	0	2554	70	0
9	I	1349	0	1293	23	0
10	J	1162	0	1238	29	0
11	K	707	0	771	20	0
12	L	4807	0	4847	108	0
13	M	3887	0	4036	83	0
14	N	3820	0	3926	70	0
15	O	956	0	968	19	0
16	P	2560	0	2605	68	0
17	Q	939	0	928	8	0
18	R	482	0	478	8	0
19	S	727	0	759	18	0
20	T	667	0	648	16	0
21	U	650	0	635	13	0
22	V	1123	0	1112	22	0
23	W	904	0	901	15	0
24	X	767	0	766	9	0
25	Z	997	0	979	14	0
26	a	469	0	472	11	0
27	b	295	0	322	8	0
28	c	617	0	619	15	0
29	d	592	0	610	7	0
30	e	557	0	525	7	0
31	f	763	0	767	15	0
32	g	615	0	604	10	0
33	i	737	0	705	7	0
34	j	415	0	406	6	0
35	k	374	0	369	6	0
36	l	502	0	504	10	0
37	m	577	0	564	12	0
38	n	911	0	879	22	0
39	o	657	0	671	14	0
40	p	778	0	766	9	0
41	q	536	0	495	13	0
42	r	89	0	95	4	0
43	u	150	0	34	0	0
44	v	226	0	235	1	0
45	x	1659	0	1673	21	0
46	y	2032	0	2044	30	0
47	z	1772	0	1771	35	0
48	B	8	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
48	F	8	0	0	2	0
48	G	16	0	0	0	0
48	I	16	0	0	0	0
49	D	50	0	79	3	0
49	M	50	0	79	2	0
49	N	50	0	79	6	0
50	E	4	0	0	6	0
50	G	4	0	0	0	0
51	F	31	0	19	1	0
52	H	35	0	43	2	0
53	M	69	0	88	1	0
54	O	1	0	0	0	0
55	P	48	0	26	1	0
56	R	1	0	0	0	0
56	y	1	0	0	0	0
57	W	35	0	0	0	0
57	n	35	0	0	1	0
58	c	52	0	84	13	0
58	v	52	0	84	1	0
59	f	41	0	52	3	0
60	f	61	0	0	0	0
61	z	30	0	32	3	0
All	All	61765	0	61904	1023	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 1023 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:180:MET:HE3	50:E:500:FES:S2	1.77	1.25
5:E:169:MET:HE1	50:E:500:FES:S2	1.98	1.03
16:P:322:ARG:NH1	16:P:333:LEU:HD23	1.78	0.98
3:C:60:CYS:HB3	4:D:354:LYS:HD2	1.54	0.88
2:B:80:ARG:HA	8:H:57:LYS:HD3	1.56	0.88

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	88/119 (74%)	86 (98%)	2 (2%)	0	100	100
2	B	155/218 (71%)	147 (95%)	8 (5%)	0	100	100
3	C	183/190 (96%)	179 (98%)	4 (2%)	0	100	100
4	D	383/394 (97%)	368 (96%)	15 (4%)	0	100	100
5	E	190/255 (74%)	182 (96%)	8 (4%)	0	100	100
6	F	432/486 (89%)	421 (98%)	11 (2%)	0	100	100
7	G	686/748 (92%)	654 (95%)	31 (4%)	1 (0%)	48	78
8	H	307/325 (94%)	289 (94%)	17 (6%)	1 (0%)	37	67
9	I	163/222 (73%)	161 (99%)	2 (1%)	0	100	100
10	J	143/205 (70%)	136 (95%)	7 (5%)	0	100	100
11	K	88/100 (88%)	84 (96%)	4 (4%)	0	100	100
12	L	613/669 (92%)	582 (95%)	31 (5%)	0	100	100
13	M	483/495 (98%)	474 (98%)	8 (2%)	1 (0%)	44	73
14	N	486/499 (97%)	474 (98%)	12 (2%)	0	100	100
15	O	120/159 (76%)	115 (96%)	5 (4%)	0	100	100
16	P	330/402 (82%)	318 (96%)	12 (4%)	0	100	100
17	Q	117/154 (76%)	111 (95%)	6 (5%)	0	100	100
18	R	60/110 (54%)	57 (95%)	3 (5%)	0	100	100
19	S	91/97 (94%)	81 (89%)	10 (11%)	0	100	100
20	T	82/122 (67%)	77 (94%)	5 (6%)	0	100	100
21	U	81/126 (64%)	75 (93%)	6 (7%)	0	100	100
22	V	138/169 (82%)	135 (98%)	3 (2%)	0	100	100
23	W	108/133 (81%)	101 (94%)	7 (6%)	0	100	100
24	X	95/106 (90%)	92 (97%)	3 (3%)	0	100	100
25	Z	123/143 (86%)	118 (96%)	4 (3%)	1 (1%)	16	45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
26	a	56/65 (86%)	56 (100%)	0	0	100	100
27	b	38/65 (58%)	38 (100%)	0	0	100	100
28	c	74/88 (84%)	74 (100%)	0	0	100	100
29	d	73/81 (90%)	71 (97%)	2 (3%)	0	100	100
30	e	63/83 (76%)	63 (100%)	0	0	100	100
31	f	99/106 (93%)	96 (97%)	3 (3%)	0	100	100
32	g	73/114 (64%)	71 (97%)	2 (3%)	0	100	100
33	i	83/98 (85%)	80 (96%)	3 (4%)	0	100	100
34	j	49/69 (71%)	44 (90%)	5 (10%)	0	100	100
35	k	45/72 (62%)	42 (93%)	3 (7%)	0	100	100
36	l	62/125 (50%)	57 (92%)	5 (8%)	0	100	100
37	m	68/71 (96%)	63 (93%)	5 (7%)	0	100	100
38	n	107/117 (92%)	105 (98%)	2 (2%)	0	100	100
39	o	78/103 (76%)	76 (97%)	2 (3%)	0	100	100
40	p	91/106 (86%)	87 (96%)	4 (4%)	0	100	100
41	q	63/159 (40%)	56 (89%)	7 (11%)	0	100	100
42	r	8/131 (6%)	7 (88%)	1 (12%)	0	100	100
44	v	28/113 (25%)	27 (96%)	1 (4%)	0	100	100
45	x	212/256 (83%)	206 (97%)	6 (3%)	0	100	100
46	y	266/278 (96%)	255 (96%)	11 (4%)	0	100	100
47	z	231/275 (84%)	227 (98%)	4 (2%)	0	100	100
All	All	7612/9221 (83%)	7318 (96%)	290 (4%)	4 (0%)	50	78

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
8	H	201	ALA
7	G	253	GLU
13	M	231	VAL
25	Z	120	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/106 (78%)	83 (100%)	0	100	100
2	B	132/184 (72%)	132 (100%)	0	100	100
3	C	175/179 (98%)	175 (100%)	0	100	100
4	D	331/340 (97%)	331 (100%)	0	100	100
5	E	166/220 (76%)	165 (99%)	1 (1%)	84	90
6	F	353/396 (89%)	353 (100%)	0	100	100
7	G	571/625 (91%)	571 (100%)	0	100	100
8	H	262/272 (96%)	262 (100%)	0	100	100
9	I	147/195 (75%)	147 (100%)	0	100	100
10	J	131/186 (70%)	131 (100%)	0	100	100
11	K	78/86 (91%)	78 (100%)	0	100	100
12	L	518/568 (91%)	518 (100%)	0	100	100
13	M	426/434 (98%)	426 (100%)	0	100	100
14	N	406/416 (98%)	406 (100%)	0	100	100
15	O	107/141 (76%)	107 (100%)	0	100	100
16	P	273/334 (82%)	273 (100%)	0	100	100
17	Q	100/128 (78%)	100 (100%)	0	100	100
18	R	55/97 (57%)	55 (100%)	0	100	100
19	S	82/85 (96%)	82 (100%)	0	100	100
20	T	79/112 (70%)	79 (100%)	0	100	100
21	U	75/113 (66%)	75 (100%)	0	100	100
22	V	123/148 (83%)	123 (100%)	0	100	100
23	W	98/114 (86%)	98 (100%)	0	100	100
24	X	87/94 (93%)	87 (100%)	0	100	100
25	Z	100/115 (87%)	100 (100%)	0	100	100
26	a	48/53 (91%)	48 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
27	b	33/53 (62%)	33 (100%)	0	100	100
28	c	66/71 (93%)	66 (100%)	0	100	100
29	d	60/66 (91%)	60 (100%)	0	100	100
30	e	60/73 (82%)	60 (100%)	0	100	100
31	f	81/84 (96%)	81 (100%)	0	100	100
32	g	65/96 (68%)	65 (100%)	0	100	100
33	i	77/90 (86%)	77 (100%)	0	100	100
34	j	42/51 (82%)	42 (100%)	0	100	100
35	k	38/60 (63%)	38 (100%)	0	100	100
36	l	53/97 (55%)	53 (100%)	0	100	100
37	m	58/59 (98%)	58 (100%)	0	100	100
38	n	92/99 (93%)	92 (100%)	0	100	100
39	o	70/87 (80%)	70 (100%)	0	100	100
40	p	83/93 (89%)	83 (100%)	0	100	100
41	q	54/133 (41%)	54 (100%)	0	100	100
42	r	10/118 (8%)	10 (100%)	0	100	100
44	v	23/84 (27%)	23 (100%)	0	100	100
45	x	183/216 (85%)	183 (100%)	0	100	100
46	y	223/232 (96%)	223 (100%)	0	100	100
47	z	188/228 (82%)	188 (100%)	0	100	100
All	All	6565/7831 (84%)	6564 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	E	135	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 71 such sidechains are listed below:

Mol	Chain	Res	Type
45	x	116	ASN
46	y	42	HIS
47	z	101	GLN
12	L	363	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
12	L	340	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 3 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
48	SF4	G	802	7	0,12,12	-	-	-		
57	8Q1	n	200	-	31,34,34	1.68	5 (16%)	40,43,43	1.56	4 (10%)
48	SF4	B	500	2	0,12,12	-	-	-		
59	PGT	f	201	-	40,40,50	1.16	3 (7%)	43,46,56	1.11	2 (4%)
58	PC7	v	201	-	51,51,51	0.96	4 (7%)	57,59,59	1.04	2 (3%)
55	NDP	P	500	-	45,52,52	2.26	5 (11%)	53,80,80	1.74	11 (20%)
48	SF4	G	803	7	0,12,12	-	-	-		
57	8Q1	W	200	-	31,34,34	1.71	6 (19%)	40,43,43	1.56	4 (10%)
48	SF4	I	501	9	0,12,12	-	-	-		
48	SF4	F	501	6	0,12,12	-	-	-		
61	PSF	z	301	-	28,29,29	1.18	4 (14%)	32,36,36	1.22	2 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
49	PTY	M	501	-	49,49,49	0.88	4 (8%)	52,54,54	1.10	2 (3%)
50	FES	G	801	7	0,4,4	-	-	-	-	-
60	T7X	f	202	-	61,61,61	0.84	4 (6%)	71,73,73	1.06	2 (2%)
48	SF4	I	500	9	0,12,12	-	-	-	-	-
53	LMN	M	502	-	72,72,72	1.65	14 (19%)	96,98,98	1.02	3 (3%)
49	PTY	D	401	-	49,49,49	0.87	4 (8%)	52,54,54	1.10	2 (3%)
51	FMN	F	500	-	33,33,33	1.09	2 (6%)	48,50,50	1.22	8 (16%)
49	PTY	N	501	-	49,49,49	0.86	4 (8%)	52,54,54	1.11	2 (3%)
50	FES	E	500	5	0,4,4	-	-	-	-	-
52	UQ9	H	500	-	35,35,58	2.49	13 (37%)	42,45,73	1.94	12 (28%)
58	PC7	c	101	-	51,51,51	0.97	4 (7%)	57,59,59	1.03	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
48	SF4	G	802	7	-	-	0/6/5/5
57	8Q1	n	200	-	-	19/41/41/41	-
48	SF4	B	500	2	-	-	0/6/5/5
59	PGT	f	201	-	-	22/45/45/55	-
58	PC7	v	201	-	-	20/55/55/55	-
55	NDP	P	500	-	-	8/30/77/77	0/5/5/5
57	8Q1	W	200	-	-	25/41/41/41	-
48	SF4	G	803	7	-	-	0/6/5/5
48	SF4	I	501	9	-	-	0/6/5/5
48	SF4	F	501	6	-	-	0/6/5/5
61	PSF	z	301	-	-	12/35/35/35	-
49	PTY	M	501	-	-	25/53/53/53	-
50	FES	G	801	7	-	-	0/1/1/1
60	T7X	f	202	-	-	29/56/80/80	0/1/1/1
48	SF4	I	500	9	-	-	0/6/5/5
53	LMN	M	502	-	-	38/50/130/130	0/4/4/4
49	PTY	D	401	-	-	19/53/53/53	-
51	FMN	F	500	-	-	8/18/18/18	0/3/3/3
49	PTY	N	501	-	-	23/53/53/53	-
50	FES	E	500	5	-	-	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
52	UQ9	H	500	-	-	7/30/54/81	0/1/1/1
58	PC7	c	101	-	-	29/55/55/55	-

The worst 5 of 76 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
55	P	500	NDP	P2B-O2B	12.70	1.83	1.59
52	H	500	UQ9	C6-C1	9.77	1.53	1.35
57	W	200	8Q1	C34-N36	5.54	1.45	1.33
57	n	200	8Q1	C34-N36	5.45	1.45	1.33
57	W	200	8Q1	C39-N41	5.37	1.45	1.33

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
55	P	500	NDP	PN-O3-PA	-7.01	108.78	132.83
57	n	200	8Q1	C6-C1-S44	5.65	120.03	113.46
57	W	200	8Q1	C6-C1-S44	5.61	119.98	113.46
52	H	500	UQ9	C7-C6-C5	5.45	125.03	118.48
60	f	202	T7X	O16-C10-C12	4.30	120.76	111.50

There are no chirality outliers.

5 of 284 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
49	D	401	PTY	N1-C2-C3-O11
49	D	401	PTY	C5-O14-P1-O13
49	M	501	PTY	N1-C2-C3-O11
49	M	501	PTY	O10-C8-O7-C6
49	M	501	PTY	C11-C8-O7-C6

There are no ring outliers.

14 monomers are involved in 45 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
57	n	200	8Q1	1	0
59	f	201	PGT	3	0
58	v	201	PC7	1	0
55	P	500	NDP	1	0
48	F	501	SF4	2	0



Continued on next page...


Continued from previous page...



Mol	Chain	Res	Type	Clashes	Symm-Clashes
61	z	301	PSF	3	0
49	M	501	PTY	2	0
53	M	502	LMN	1	0
49	D	401	PTY	3	0
51	F	500	FMN	1	0
49	N	501	PTY	6	0
50	E	500	FES	6	0
52	H	500	UQ9	2	0
58	c	101	PC7	13	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
13	M	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	M	270:PRO	C	271:MET	N	3.13


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-11878. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 300



Y Index: 300




Z Index: 300


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 222



Y Index: 268




Z Index: 334


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.011. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 303 nm³; this corresponds to an approximate mass of 274 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.293 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.293 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.41	-	-
Author-provided FSC curve	3.40	4.07	3.45
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-11878 and PDB model 7ARB. Per-residue inclusion information can be found in section [3](#) on page [20](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.011 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.011).

9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.011) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7370	 0.5440
A	 0.6970	 0.5420
B	 0.9140	 0.5780
C	 0.8390	 0.5960
D	 0.8460	 0.5850
E	 0.7070	 0.5280
F	 0.7550	 0.5610
G	 0.7960	 0.5810
H	 0.7350	 0.5350
I	 0.8800	 0.5820
J	 0.6910	 0.5410
K	 0.6770	 0.5450
L	 0.7070	 0.5370
M	 0.7450	 0.5480
N	 0.7580	 0.5680
O	 0.4740	 0.5190
P	 0.7480	 0.5180
Q	 0.7550	 0.5930
R	 0.7680	 0.5860
S	 0.7690	 0.5600
T	 0.6220	 0.4570
U	 0.1300	 0.2530
V	 0.7480	 0.5390
W	 0.5240	 0.4890
X	 0.6700	 0.5170
Z	 0.7280	 0.5220
a	 0.7010	 0.5150
b	 0.6600	 0.5300
c	 0.7600	 0.5430
d	 0.7660	 0.5520
e	 0.7930	 0.5590
f	 0.7830	 0.5690
g	 0.7160	 0.5160
i	 0.7260	 0.5190
j	 0.7400	 0.5170



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
k	 0.7020	 0.5040
l	 0.5850	 0.4970
m	 0.6070	 0.4680
n	 0.6590	 0.4890
o	 0.7430	 0.5130
p	 0.7480	 0.4980
q	 0.4920	 0.4870
r	 0.3810	 0.4860
u	 0.7270	 0.4840
v	 0.6780	 0.5540
x	 0.7900	 0.5640
y	 0.7300	 0.5450
z	 0.7780	 0.5580