

Mar 5, 2025 – 12:26 AM EST

PDB ID	:	9B4J
EMDB ID	:	EMD-44182
Title	:	Filament of D-TLKIVWS, a D-peptide that disaggregates Alzheimer's Paired
		Helical Filaments, determined by Cryo-EM
Authors	:	Hou, K.; Ge, P.; Sawaya, M.R.; Eisenberg, D.S.
Deposited on	:	2024-03-20
Resolution	:	3.50  Å(reported)
This is	s a Ì	Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev117
MolProbity	:	4.02b-467
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 3.50 Å.

There are no overall percentile quality scores available for this entry.

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq=3, 2, 1$  and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq=5\%$  The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	0	7	100%
1	1	7	100%
1	1	•	100%
1	2	7	100%
1	3	7	100%
1	4	7	100%
1	5	7	100%
1	6	7	100%
1	7	7	100%
1	8	7	100%
1	9	7	100%
-			14%
1	А	7	100%
1	В	7	14%
1	С	7	100%
1	D	7	100%
1	Е	7	100%
1	F	7	100%
1	G	7	100%



Continued from previous page...

Mol	Chain	Length	Quality of chain
1	Н	7	100%
1	Ι	7	100%
1	J	7	100%
1	K	7	100%
1	L	7	100%
1	М	7	100%
1	N	7	100%
1	О	7	100%
1	P	7	100%
1	Q	7	100%
1	R	7	100%
1	S	7	14%
1	T	7	100%
			100%
1	U	7	100%
1	V	7	100%
1	W	7	100%
1	Х	7	100%
1	Y	7	100%
1	Z	7	100%
1	a	7	100%
1	b	7	100%
1	с	7	100%
1	d	7	100%
1	е	7	100%
1	f	7	100%



Continued from previous page...

Mol	Chain	Length	Quality of chain
1	g	7	100%
1	h	7	100%
1	i	7	100%
1	j	7	100%
1	j k	7	
			100%
1	1	7	100%
1	m	7	100%
1	n	7	100%
1	0	7	100%
1	р	7	100%
1	q	7	100%
1	r	7	100%
1	S	7	100%
1	t	7	100%
1	u	7	100%
1	V	7	100%
1	W	7	14%
1	x	7	14%
	<u> </u>	•	100 /0



# 2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 3600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein (with D amino acids) called DTH-DLE-DLY-DIL-DVA-DTR-DSN.

Mol	Chain	Residues	Atoms		AltConf	Trace
1	А	7	Total C N	V O	0	0
1	A	1	60 41 9	) 10	0	0
1	В	7	Total C N	V O	0	0
1	D	4	60 41 9		0	0
1	С	7	Total C N		0	0
1	U	4	60 41 9		0	0
1	D	7	Total C N		0	0
	D	1	60 41 9		0	0
1	Е	7	Total C N	-	0	0
		•	60 41 9		0	0
1	F	7	Total C N	-	0	0
-	1	•	60 41 9		0	
1	G	7	Total C N		0	0
		•	60 41 9		0	
1	Н	7	Total C N		0	0
-		•	60 41 9		Ŭ	<u> </u>
1	Ι	7	Total C N		0	0
	-	•	60 41 9		Ŭ	
1	J	7	Total C N		0	0
		-	60 41 9		_	
1	Κ	7	Total C N		0	0
			60 41 9			
1	L	7	Total C N		0	0
			60 41 9			
1	М	7	Total C N		0	0
			60  41  9			
1	Ν	7	Total C N	-	0	0
			60  41  9			
1	Ο	7	Total C N	-	0	0
			60  41  9			
1	Р	7	Total C N	-	0	0
			60  41  9			
1	Q	7	Total C N		0	0
	L Č		60 41 9	) 10		



Continued from previous page...

Mol	Chain	Residues	<u> </u>	on	ıs		AltConf	Trace
1	D	7	Total (	С	Ν	0	0	0
1	R	(	60 4	41	9	10	0	0
1	S	7	Total	С	Ν	0	0	0
	G	1	60 4	41	9	10	0	0
1	Т	7	Total (	С	Ν	0	0	0
	1	1	60 4	41	9	10	0	0
1	U	7	Total (	С	Ν	0	0	0
T	0	1		41	9	10	0	0
1	V	7		С	Ν	0	0	0
T	v	1	60 4	41	9	10	0	0
1	W	7		С	Ν	Ο	0	0
1	**	1		41	9	10	0	0
1	Х	7		С	Ν	Ο	0	0
-		•		41	9	10	0	0
1	Y	7		С	Ν	Ο	0	0
	-	•		41	9	10	0	0
1	Z	7		$\mathbf{C}$	Ν	Ο	0	0
		•		41	9	10		0
1	0	7		С	Ν	Ο	0	0
	, , , , , , , , , , , , , , , , , , ,	•		41	9	10		
1	1	7		С	Ν	0	0	0
		-		41 G	9	10	_	
1	2	7		C	N	0	0	0
				41 C	9	10		
1	3	7		C	N	0	0	0
				$\frac{41}{C}$	9 N	10		
1	4	7				0 10	0	0
				$\frac{41}{C}$	9 N	10 0		
1	5	7		U 41	N 9	10	0	0
				$\frac{1}{C}$	9 N	$\frac{10}{0}$		
1	6	7		11 41	1N 9	10	0	0
				$\frac{1}{C}$	N	$\frac{10}{0}$		
1	7	7		U 41	1N 9	10	0	0
				$\frac{1}{C}$	N	$\frac{10}{0}$		
1	8	7		41	9	10	0	0
				$\frac{1}{C}$	N	$\frac{10}{0}$		
1	9	7		41	9	10	0	0
				C	N	0		
1	a	7		41	9	10	0	0
				$\frac{\Gamma}{C}$	N	$\frac{10}{0}$		
1	b	7		41	9	10	0	0
l			00		0		_	



Continued from previous page...

Mol	Chain	Residues	At	on	ıs		AltConf	Trace
1		7	Total (	С	Ν	0	0	0
1	С	7	60 4	11	9	10	0	0
1	d	7	Total (	С	Ν	0	0	0
1	d	(	60 4	11	9	10	0	0
1		7	Total (	С	Ν	0	0	0
	е	1	60 4	11	9	10	0	0
1	f	7	Total (	С	Ν	0	0	0
1	1	1	60 4	11	9	10	0	0
1	ď	7	Total (	С	Ν	0	0	0
T	g	1	60 4	11	9	10	0	0
1	h	7	Total (	С	Ν	0	0	0
T	11	1		11	9	10	0	0
1	i	7		С	Ν	Ο	0	0
-	1	•		11	9	10	0	0
1	j	7		С	Ν	Ο	0	0
	J	•		11	9	10	•	0
1	k	7		С	Ν	Ο	0	0
		•		11	9	10		Ŭ
1	1	7		С	Ν	Ο	0	0
	-	•		11	9	10		
1	m	7		С	Ν	0	0	0
				11	9	10		
1	n	7		С	Ν	0	0	0
		-		11	9	10		
1	0	7		С	N	0	0	0
				11 	9 N	10		
1	р	7		С	N	0	0	0
	-			11 G	9	10		
1	q	7		C	N	0	0	0
	_			11 	$\frac{9}{N}$	10		
1	r	7		C	N 0	0	0	0
				11 C	$\frac{9}{N}$	10 0		
1	S	7		U 11	N 9		0	0
				$\frac{1}{C}$	9 N	10 0		
1	t	7		U 11	N 9	10	0	0
				$\frac{1}{C}$	9 N	$\frac{10}{0}$		
1	u	7		11	1N 9	10	0	0
				$\frac{1}{C}$	9 N	$\frac{10}{0}$		
1	v	7		11	1N 9	10	0	0
				$\frac{1}{C}$	9 N	$\frac{10}{0}$		
1	W	7		11	9	10	0	0
			00 4	τ⊥	3	10		



Continued from previous page...


Mol	Chain	Residues	A	4ton	ns		AltConf	Trace
1	x	7	Total 60	C 41	N 9	O 10	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN



Chain G: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain H: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain I: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain J: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain K: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain L: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain M: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain N: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain O: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain P: 100%



There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain Q: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain R: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain S: 100%
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain T: 100%
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain U: 100%
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain V: 100%
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain W: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain X: 100%



There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain Y: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain Z: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain 0: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain 1: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain 2: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain 3: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain 4: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain 5: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain 6: 100%
There are no outlier residues recorded for this chain

There are no outlier residues recorded for this chain.



• Molecule 1: DTH-DLE-I	)LY-DIL-DVA-DTR-DSN	
Chain 7:	100%	
There are no outlier residu	es recorded for this chain.	
• Molecule 1: DTH-DLE-I	DLY-DIL-DVA-DTR-DSN	
Chain 8:	100%	
There are no outlier residu	es recorded for this chain.	
• Molecule 1: DTH-DLE-I	DLY-DIL-DVA-DTR-DSN	
Chain 9:	100%	
There are no outlier residu	es recorded for this chain.	
• Molecule 1: DTH-DLE-I	DLY-DIL-DVA-DTR-DSN	
Chain a:	100%	
There are no outlier residu	es recorded for this chain.	
• Molecule 1: DTH-DLE-I	DLY-DIL-DVA-DTR-DSN	
Chain b:	100%	
There are no outlier residu	es recorded for this chain.	
• Molecule 1: DTH-DLE-I	DLY-DIL-DVA-DTR-DSN	
Chain c:	100%	
•		
1 5		
• Molecule 1: DTH-DLE-I	DLY-DIL-DVA-DTR-DSN	
Chain d:		
Cham d:	100%	
₽ Fi Ka		
• Molecule 1: DTH-DLE-I	DLY-DIL-DVA-DTR-DSN	
Chain e:	100%	
• •		
2 <mark>7 1</mark> 1		

• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN




Chain f: 100%
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain g: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain h: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain i: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain j: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain k: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain I: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain m: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain n: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN



Chain o: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain p: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain q: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain r: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain s: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain t: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain u: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain v: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN
Chain w: 100%

• Molecule 1: DTH-DLE-DLY-DIL-DVA-DTR-DSN







# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=3.509°, rise=9.558 Å, axial	Depositor
	sym=C1	
Number of segments used	32435	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	50	Depositor
Minimum defocus (nm)	1800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	24.899	Depositor
Minimum map value	-11.872	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	6.5	Depositor
Map size (Å)	341.44, 341.44, 341.44	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ( $^{\circ}$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.067, 1.067, 1.067	Depositor



# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DIL, DVA, DLE, DTH, DSN, DLY, DTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

There are no protein molecules in this entry.

#### 5.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

420 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




# 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-44182. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

# 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.



## 6.2 Central slices (i)

#### 6.2.1 Primary map



X Index: 160

Y Index: 160





The images above show central slices of the map in three orthogonal directions.



## 6.3 Largest variance slices (i)

#### 6.3.1 Primary map





X Index: 165

Y Index: 152




The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) (i)

#### 6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




#### 6.5 Orthogonal surface views (i)

#### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 6.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




## 6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

#### 6.6.1 emd\_44182\_msk\_1.map (i)






# 7 Map analysis (i)

This section contains the results of statistical analysis of the map.


# 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 7.2 Volume estimate (i)




The volume at the recommended contour level is 227  $\rm nm^3;$  this corresponds to an approximate mass of 205 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum (i)




\*Reported resolution corresponds to spatial frequency of 0.286  ${\rm \AA^{-1}}$ 



# 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

#### 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.286  $\text{\AA}^{-1}$ 



# 8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
Resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.46	4.04	3.49
Unmasked-calculated*	3.81	4.33	3.92


\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.



# 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-44182 and PDB model 9B4J. Per-residue inclusion information can be found in section 3 on page 9.


## 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 6.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




#### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


#### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (6.5).



### 9.4 Atom inclusion (i)



At the recommended contour level, 86% of all backbone atoms, 70% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

## 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (6.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7020	0.4750
0	0.6440	0.4920
1	0.7290	0.5110
2	0.7290	0.4770
3	0.7120	0.5100
4	0.7290	0.5050
5	0.7120	0.4780
6	0.7630	0.5000
7	0.7120	0.5050
8	0.8140	0.5210
9	0.7970	0.5110
А	0.7120	0.5170
В	0.6610	0.5220
С	0.7630	0.5080
D	0.7460	0.4850
Е	0.8140	0.5340
F	0.6950	0.5100
G	0.6440	0.4900
Н	0.7290	0.5110
Ι	0.6950	0.4760
J	0.7290	0.5100
Κ	0.7460	0.5140
$\mathbf{L}$	0.7290	0.4820
М	0.7630	0.4980
Ν	0.6950	0.5100
О	0.7970	0.5210
Р	0.7800	0.5080
Q	0.6610	0.4450
R	0.7120	0.4840
S	0.4580	0.2340
Т	0.4750	0.2400
U	0.6610	0.5220
V	0.6780	0.5160
W	0.7630	0.5110
Х	0.7460	0.4880



Continued from previous page...

Chain	Atom inclusion	Q-score
Y	0.7970	0.5390
Z	0.7120	0.5100
a	0.6950	0.4520
b	0.6950	0.4780
с	0.4580	0.2370
d	0.4750	0.2230
е	0.6780	0.5280
f	0.6440	0.5220
g	0.7970	0.5100
h	0.7460	0.4890
i	0.8140	0.5280
j	0.7630	0.5200
k	0.6440	0.4870
1	0.7290	0.5120
m	0.7120	0.4770
n	0.7290	0.5100
0	0.7630	0.5170
р	0.7120	0.4800
q	0.7630	0.4920
r	0.7120	0.5110
s	0.8140	0.5210
t	0.7630	0.5030
u	0.6610	0.4440
V	0.7120	0.4800
W	0.4580	0.2260
Х	0.4750	0.2460

