

wwPDB X-ray Structure Validation Summary Report (i)

Apr 28, 2025 – 11:02 PM EDT

PDB ID	:	$3 \mathrm{BZU} \ / \ \mathrm{pdb} \ 00003 \mathrm{bzu}$
Title	:	Crystal structure of human 11-beta-hydroxysteroid dehydrogenase(HSD1) in
		complex with NADP and thiazolone inhibitor
Authors	:	Min, X.; Sudom, A.; Xu, H.; Wang, Z.; Walker, N.P.
Deposited on	:	2008-01-18
Resolution	:	2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	286	^{2%} 72%	18%	•• 8%
1	В	286	~ 74%	17%	• 8%
1	С	286	70%	18%	11%
1	D	286	<mark>6%</mark> 71%	15%	•• 11%



3BZU

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 8581 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	٨	263	Total	С	Ν	0	\mathbf{S}	0	C	0
	А	205	2039	1299	341	383	16	0	6	0
1	В	262	Total	С	Ν	0	S	0	ົງ	0
	D	202	2011	1283	339	374	15	0	3	0
1	С	254	Total	С	Ν	0	S	0	9	0
	C	204	1964	1254	330	365	15	0	3	0
1	1 D	254	Total	С	Ν	0	S	0	1	0
		254	1946	1244	329	358	15	0		0

• Molecule 1 is a protein called Corticosteroid 11-beta-dehydrogenase isozyme 1.

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Actual Comment	
А	7	MET	-	expression tag	UNP P28845
А	8	LYS	-	expression tag	UNP P28845
A	9	HIS	-	expression tag	UNP P28845
А	10	GLN	-	expression tag	UNP P28845
А	11	HIS	-	expression tag	UNP P28845
А	12	GLN	-	expression tag	UNP P28845
А	13	HIS	-	expression tag	UNP P28845
А	14	GLN	-	expression tag	UNP P28845
А	15	HIS	-	expression tag	UNP P28845
А	16	GLN	-	expression tag	UNP P28845
А	17	HIS	-	expression tag	UNP P28845
А	18	GLN	-	expression tag	UNP P28845
A	19	HIS	-	expression tag	UNP P28845
А	20	GLN	-	expression tag	UNP P28845
A	21	GLN	-	expression tag	UNP P28845
А	22	PRO	-	expression tag	UNP P28845
А	23	LEU	-	expression tag	UNP P28845
А	272	SER	CYS	engineered mutation	UNP P28845
В	7	MET	-	expression tag	UNP P28845
В	8	LYS	-	expression tag	UNP P28845
В	9	HIS	-	expression tag	UNP P28845



Chain	Residue	Modelled	Actual	Comment	Reference
В	10	GLN	-	expression tag	UNP P28845
В	11	HIS	-	expression tag	UNP P28845
В	12	GLN	-	expression tag	UNP P28845
В	13	HIS	-	expression tag	UNP P28845
В	14	GLN	-	expression tag	UNP P28845
В	15	HIS	-	expression tag	UNP P28845
В	16	GLN	-	expression tag	UNP P28845
В	17	HIS	-	expression tag	UNP P28845
В	18	GLN	-	expression tag	UNP P28845
В	19	HIS	-	expression tag	UNP P28845
В	20	GLN	-	expression tag	UNP P28845
В	21	GLN	-	expression tag	UNP P28845
В	22	PRO	-	expression tag	UNP P28845
В	23	LEU	-	expression tag	UNP P28845
В	272	SER	CYS	engineered mutation	UNP P28845
С	7	MET	-	expression tag	UNP P28845
С	8	LYS	-	expression tag	UNP P28845
С	9	HIS	-	expression tag	UNP P28845
С	10	GLN	-	expression tag	UNP P28845
С	11	HIS	-	expression tag	UNP P28845
С	12	GLN	-	expression tag	UNP P28845
С	13	HIS	-	expression tag	UNP P28845
С	14	GLN	-	expression tag	UNP P28845
С	15	HIS	-	expression tag	UNP P28845
С	16	GLN	-	expression tag	UNP P28845
С	17	HIS	-	expression tag	UNP P28845
С	18	GLN	-	expression tag	UNP P28845
С	19	HIS	-	expression tag	UNP P28845
С	20	GLN	-	expression tag	UNP P28845
С	21	GLN	-	expression tag	UNP P28845
С	22	PRO	-	expression tag	UNP P28845
С	23	LEU	-	expression tag	UNP P28845
С	272	SER	CYS	engineered mutation	UNP P28845
D	7	MET	-	expression tag	UNP P28845
D	8	LYS	-	expression tag	UNP P28845
D	9	HIS	-	expression tag	UNP P28845
D	10	GLN	-	expression tag	UNP P28845
D	11	HIS	-	expression tag	UNP P28845
D	12	GLN	-	expression tag	UNP P28845
D	13	HIS	-	expression tag	UNP P28845
D	14	GLN	-	expression tag	UNP P28845
D	15	HIS	-	expression tag	UNP P28845

Continued from previous page...
Chain Residue Modelled Actual




3BZU
00

Chain	Residue	Modelled	Actual	Comment	Reference
D	16	GLN	-	expression tag	UNP P28845
D	17	HIS	-	expression tag	UNP P28845
D	18	GLN	-	expression tag	UNP P28845
D	19	HIS	-	expression tag	UNP P28845
D	20	GLN	-	expression tag	UNP P28845
D	21	GLN	-	expression tag	UNP P28845
D	22	PRO	-	expression tag	UNP P28845
D	23	LEU	-	expression tag	UNP P28845
D	272	SER	CYS	engineered mutation	UNP P28845

Continued from previous page...


• Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues		Ate	oms		ZeroOcc	AltConf	
2	Λ	1	Total	С	Ν	Ο	Р	0	0
	А	1	48	21	7	17	3	0	0
2	В	1	Total	С	Ν	Ο	Р	0	0
	2 Б		48	21	7	17	3	0	0
2	С	1	Total	С	Ν	Ο	Р	0	0
			48	21	7	17	3	0	0
2	2 D	D 1	Total	С	Ν	Ο	Р	0	0
		1	48	21	$\overline{7}$	17	3	0	0

• Molecule 3 is (5S)-2-{[(1S)-1-(2-fluorophenyl)ethyl]amino}-5-methyl-5-(trifluoromethyl)-1,3 -thiazol-4(5H)-one (CCD ID: A21) (formula: $C_{13}H_{12}F_4N_2OS$).





Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	Δ	1	Total	С	F	Ν	0	S	0	0
5	A	1	21	13	4	2	1	1	0	0
3	р	1	Total	С	F	Ν	0	S	0	0
5	3 D	1	21	13	4	2	1	1	0	
2	С	1	Total	С	F	Ν	0	S	0	0
5	3 C		21	13	4	2	1	1	0	0
2	3 D	1	Total	С	F	Ν	0	S	0	0
5		1	21	13	4	2	1	1	0	0


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	104	Total O 104 104	0	0
4	В	99	Total O 99 99	0	0
4	С	97	Total O 97 97	0	0
4	D	45	Total O 45 45	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Corticosteroid 11-beta-dehydrogenase isozyme 1









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Depositor
Resolution (Å)	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	Depositor EDS
% Data completeness (in resolution range)	96.5 (31.33-2.25) 96.5 (31.33-2.25)	Depositor EDS
R _{merge}	0.05	Depositor
R _{sym}	0.05	Depositor
$< I/\sigma(I) > 1$	$3.39 (at 2.24 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.3.0026	Depositor
R, R_{free}	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	Depositor DCC
R_{free} test set	2862 reflections $(5.06%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	45.5	Xtriage
Anisotropy	0.337	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, 41.0	EDS
L-test for twinning ²	$< L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.036 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8581	wwPDB-VP
Average B, all atoms $(Å^2)$	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, $\mathrm{A21}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.80	0/2099	0.93	1/2835~(0.0%)	
1	В	0.78	0/2058	0.99	0/2779	
1	С	0.78	0/2008	0.93	0/2710	
1	D	0.70	0/1982	0.99	2/2675~(0.1%)	
All	All	0.77	0/8147	0.96	3/10999~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	232	HIS	N-CA-C	6.72	118.60	108.46
1	D	279	LEU	CA-CB-CG	5.20	134.50	116.30
1	D	37	VAL	N-CA-C	5.12	115.27	108.11

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group		
1	А	231	VAL	Peptide		
Continued on out or a						



Continued from previous page...

Mol	Chain	Res	Type	Group
1	D	278	PHE	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2039	0	2069	34	0
1	В	2011	0	2063	35	0
1	С	1964	0	2006	31	0
1	D	1946	0	1992	42	0
2	А	48	0	25	0	0
2	В	48	0	25	1	0
2	С	48	0	25	1	0
2	D	48	0	25	2	0
3	А	21	0	12	0	0
3	В	21	0	12	0	0
3	С	21	0	12	0	0
3	D	21	0	12	0	0
4	А	104	0	0	1	0
4	В	99	0	0	8	0
4	С	97	0	0	4	0
4	D	45	0	0	1	0
All	All	8581	0	8278	132	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 132 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:227:VAL:HB	1:B:231:VAL:CG1	1.90	1.02
1:D:140:MET:HE1	1:D:186:SER:HA	1.57	0.85
1:D:243:LEU:O	1:D:247:LYS:HG3	1.81	0.79
1:A:178:PRO:O	1:A:179:MET:HB2	1.83	0.78
1:B:227:VAL:HB	1:B:231:VAL:HG12	1.64	0.77



There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	267/286~(93%)	248~(93%)	16~(6%)	3~(1%)	12 8
1	В	263/286~(92%)	246 (94%)	16~(6%)	1 (0%)	30 32
1	С	253/286~(88%)	238~(94%)	13~(5%)	2(1%)	16 14
1	D	251/286~(88%)	232 (92%)	16~(6%)	3~(1%)	11 7
All	All	1034/1144~(90%)	964 (93%)	61~(6%)	9~(1%)	14 12

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	230	ILE
1	D	263	TRP
1	А	228	SER
1	D	280	TYR
1	С	202	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	hain Analysed Rotameric Outliers		Percentiles		
1	А	226/243~(93%)	211 (93%)	15 (7%)	14 12	
1	В	222/243~(91%)	210~(95%)	12 (5%)	18 18	
1	С	215/243~(88%)	202~(94%)	13~(6%)	16 15	



Continuea from previous page							
Mol	Chain	Analysed	ysed Rotameric Outliers		Percentiles		
1	D	211/243~(87%)	196~(93%)	15 (7%)	12	10	
All	All	874/972~(90%)	819 (94%)	55~(6%)	14	14	

Continued from previous page...

5 of 55 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	С	28	ARG
1	С	222	THR
1	D	279	LEU
1	D	232	HIS
1	С	44	LYS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such side chains are listed below:

Mol	Chain	Res	Type
1	С	270	ASN
1	D	135	HIS
1	D	270	ASN
1	D	232	HIS
1	D	120	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Aol Type Chain Res Lin		Link	Bo	ond leng	ths	Bond angles			
NIOI	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	NAP	В	2	-	46,52,52	1.85	7 (15%)	61,80,80	1.66	5 (8%)
3	A21	В	293	-	22,22,22	2.05	6 (27%)	27,34,34	1.81	8 (29%)
2	NAP	D	4	-	46,52,52	1.78	4 (8%)	61,80,80	1.32	4 (6%)
2	NAP	С	3	-	46,52,52	1.77	4 (8%)	61,80,80	1.41	8 (13%)
3	A21	А	293	-	22,22,22	1.69	4 (18%)	27,34,34	1.85	9 (33%)
2	NAP	А	1	-	46,52,52	1.77	3 (6%)	61,80,80	1.57	<mark>6 (9%)</mark>
3	A21	С	293	-	22,22,22	1.72	5 (22%)	27,34,34	1.96	9 (33%)
3	A21	D	293	-	22,22,22	2.28	6 (27%)	27,34,34	1.99	11 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	В	2	-	-	8/31/67/67	0/5/5/5
3	A21	В	293	-	-	0/17/32/32	0/2/2/2
2	NAP	D	4	-	-	4/31/67/67	0/5/5/5
2	NAP	С	3	-	-	4/31/67/67	0/5/5/5
3	A21	А	293	-	-	0/17/32/32	0/2/2/2
2	NAP	А	1	-	-	6/31/67/67	0/5/5/5
3	A21	С	293	-	-	0/17/32/32	0/2/2/2
3	A21	D	293	-	-	0/17/32/32	0/2/2/2

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	А	1	NAP	O7N-C7N	9.16	1.41	1.24
2	D	4	NAP	O7N-C7N	9.10	1.41	1.24
2	С	3	NAP	O7N-C7N	8.66	1.40	1.24
3	D	293	A21	C11-S1	-8.06	1.67	1.75
2	В	2	NAP	O7N-C7N	7.87	1.38	1.24

The worst 5 of 60 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	В	2	NAP	N3A-C2A-N1A	-8.84	116.67	128.67
2	D	4	NAP	N3A-C2A-N1A	-6.59	119.73	128.67
2	А	1	NAP	O4B-C1B-N9A	6.45	117.30	108.75
2	С	3	NAP	N3A-C2A-N1A	-6.19	120.28	128.67
2	А	1	NAP	N3A-C2A-N1A	-5.92	120.64	128.67

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	1	NAP	C5D-O5D-PN-O1N
2	В	2	NAP	C5D-O5D-PN-O3
2	В	2	NAP	C5D-O5D-PN-O1N
2	С	3	NAP	C5D-O5D-PN-O3
2	С	3	NAP	C5D-O5D-PN-O1N


There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	2	NAP	1	0
2	D	4	NAP	2	0
2	С	3	NAP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(A^2)$	Q < 0.9
1	А	263/286~(91%)	0.03	6 (2%) 61 61	24, 48, 76, 87	6(2%)
1	В	262/286~(91%)	0.10	3 (1%) 77 79	28, 47, 69, 84	3 (1%)
1	С	254/286~(88%)	0.02	6 (2%) 59 60	26, 46, 73, 84	3 (1%)
1	D	254/286~(88%)	0.64	16 (6%) 27 26	29, 62, 85, 89	1 (0%)
All	All	1033/1144~(90%)	0.20	31 (3%) 52 53	24, 50, 79, 89	13 (1%)

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	281	SER	4.5
1	В	278	PHE	3.8
1	D	235	ALA	3.3
1	С	263	TRP	3.0
1	А	263	TRP	2.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q < 0.9
3	A21	А	293	21/21	0.93	0.08	$39,\!43,\!45,\!47$	0
3	A21	D	293	21/21	0.94	0.09	51,54,58,61	0
2	NAP	D	4	48/48	0.95	0.08	47,52,57,60	0
3	A21	С	293	21/21	0.96	0.06	39,45,49,52	0
3	A21	В	293	21/21	0.96	0.07	42,46,50,52	0
2	NAP	С	3	48/48	0.97	0.07	31,38,41,44	0
2	NAP	А	1	48/48	0.97	0.06	31,36,41,45	0
2	NAP	В	2	48/48	0.97	0.06	30,33,43,46	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





















6.5 Other polymers (i)

There are no such residues in this entry.

