



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 25, 2025 – 04:15 PM EDT

PDB ID : 9CBP / pdb_00009cbp
Title : Methionine synthase from *Thermus thermophilus* HB8, Homocysteine, Folate, and Cobalamin Domains, Pre-Catalytic State (Pre-Hcy-on)
Authors : Yamada, K.; Mendoza, J.; Koutmos, M.
Deposited on : 2024-06-19
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1096 (2.46-2.46)
Clashscore	180529	1178 (2.46-2.46)
Ramachandran outliers	177936	1170 (2.46-2.46)
Sidechain outliers	177891	1170 (2.46-2.46)
RSRZ outliers	164620	1096 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	874	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	901	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6971 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methionine synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	862	6624	4213	1159	1223	29	0	0	0

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	110	ALA	PHE	engineered mutation	UNP Q5SKM5
A	123	ALA	GLU	engineered mutation	UNP Q5SKM5
A	124	ALA	GLU	engineered mutation	UNP Q5SKM5
A	296	ALA	TYR	engineered mutation	UNP Q5SKM5
A	651	ALA	ASP	engineered mutation	UNP Q5SKM5
A	652	ALA	PRO	engineered mutation	UNP Q5SKM5
A	653	ALA	GLY	engineered mutation	UNP Q5SKM5

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 2-AMINO-4-MERCAPTO-BUTYRIC ACID (CCD ID: HCS) (formula: $C_4H_9NO_2S$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			8	4	1	2	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

- Molecule 5 is COBALAMIN (CCD ID: B12) (formula: $C_{62}H_{89}CoN_{13}O_{14}P$) (labeled as "Lig- and of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		

- Molecule 6 is PROPANE (CCD ID: TME) (formula: C_3H_8) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C		
6	A	1	3	3	0	0

- Molecule 7 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total 1	K 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	200	Total 200	O 200	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methionine synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.95Å 83.82Å 163.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.99 – 2.45 81.86 – 2.45	Depositor EDS
% Data completeness (in resolution range)	98.1 (81.99-2.45) 98.4 (81.86-2.45)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.162 , 0.237 0.176 , 0.241	Depositor DCC
R_{free} test set	1790 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.564	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 65.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6971	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, GOL, TME, B12, HCS, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	2/6744 (0.0%)	1.49	48/9124 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	16

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	114	ARG	NE-CZ	5.70	1.39	1.33
1	A	521	HIS	CE1-NE2	5.36	1.38	1.32

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	345	ARG	NE-CZ-NH2	-8.51	111.54	119.20
1	A	465	SER	N-CA-CB	7.91	123.85	110.49
1	A	283	ASN	CA-CB-CG	7.70	120.30	112.60
1	A	573	ASN	CA-CB-CG	-7.33	105.27	112.60
1	A	532	LEU	N-CA-C	-7.10	99.03	110.32
1	A	633	GLU	CB-CG-CD	6.95	124.42	112.60
1	A	48	ARG	O-C-N	-6.88	116.63	121.65
1	A	801	HIS	CA-CB-CG	-6.82	106.98	113.80
1	A	345	ARG	NE-CZ-NH1	6.75	128.25	121.50
1	A	543	GLU	CB-CA-C	6.54	121.15	110.88
1	A	533	LEU	N-CA-CB	-6.50	100.56	110.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	839	LEU	N-CA-CB	-6.48	100.40	110.57
1	A	337	VAL	N-CA-CB	-6.26	103.67	112.12
1	A	617	GLU	CB-CG-CD	6.25	123.23	112.60
1	A	477	ARG	CG-CD-NE	-6.08	98.62	112.00
1	A	474	ARG	NE-CZ-NH1	-5.92	115.58	121.50
1	A	635	PHE	N-CA-CB	-5.90	100.60	110.57
1	A	526	GLU	CB-CG-CD	5.87	122.57	112.60
1	A	313	GLU	CB-CG-CD	-5.84	102.67	112.60
1	A	845	GLU	CB-CG-CD	5.84	122.53	112.60
1	A	138	ARG	CB-CG-CD	5.78	124.60	111.30
1	A	443	ASP	CA-CB-CG	5.73	118.33	112.60
1	A	858	GLU	CB-CG-CD	5.70	122.28	112.60
1	A	469	GLU	CA-C-O	-5.66	114.88	120.82
1	A	559	ARG	N-CA-CB	5.51	118.23	110.12
1	A	859	ASP	CA-CB-CG	5.51	118.11	112.60
1	A	477	ARG	NE-CZ-NH1	-5.48	116.02	121.50
1	A	742	GLU	CB-CA-C	5.46	118.75	109.84
1	A	464	ASN	CA-C-N	5.39	131.84	121.54
1	A	464	ASN	C-N-CA	5.39	131.84	121.54
1	A	206	VAL	CA-CB-CG1	5.33	119.45	110.40
1	A	560	GLU	N-CA-CB	5.24	118.01	110.20
1	A	824	ARG	NE-CZ-NH2	-5.24	114.49	119.20
1	A	648	HIS	CA-CB-CG	5.20	119.00	113.80
1	A	270	PHE	CA-CB-CG	5.19	118.99	113.80
1	A	796	GLU	CB-CG-CD	5.18	121.40	112.60
1	A	738	GLU	CB-CG-CD	5.15	121.36	112.60
1	A	674	GLU	CB-CG-CD	5.15	121.35	112.60
1	A	283	ASN	CB-CA-C	5.13	120.20	110.51
1	A	696	ASP	CA-CB-CG	5.11	117.71	112.60
1	A	421	GLU	CB-CG-CD	5.11	121.28	112.60
1	A	349	PHE	CB-CA-C	5.07	116.26	109.42
1	A	811	LYS	CD-CE-NZ	5.06	128.09	111.90
1	A	281	LEU	N-CA-CB	5.05	117.36	111.09
1	A	138	ARG	CA-CB-CG	5.04	124.17	114.10
1	A	474	ARG	NH1-CZ-NH2	5.02	125.83	119.30
1	A	31	ASN	CA-C-N	5.00	127.25	120.65
1	A	31	ASN	C-N-CA	5.00	127.25	120.65

There are no chirality outliers.

All (16) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	135	ARG	Sidechain
1	A	138	ARG	Sidechain
1	A	171	ARG	Sidechain
1	A	178	ARG	Sidechain
1	A	27	ARG	Sidechain
1	A	35	ARG	Sidechain
1	A	374	ARG	Sidechain
1	A	400	ARG	Sidechain
1	A	464	ASN	Peptide
1	A	465	SER	Peptide
1	A	477	ARG	Sidechain
1	A	524	ARG	Sidechain
1	A	559	ARG	Sidechain
1	A	732	ARG	Sidechain
1	A	778	ARG	Sidechain
1	A	824	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6624	0	6743	51	0
2	A	42	0	56	13	0
3	A	8	0	7	1	0
4	A	2	0	0	0	0
5	A	91	0	88	15	0
6	A	3	0	8	1	0
7	A	1	0	0	0	0
8	A	200	0	0	9	0
All	All	6971	0	6902	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:910:B12:H351	5:A:910:B12:H362	1.46	0.97
5:A:910:B12:H552	5:A:910:B12:H531	1.48	0.93
1:A:465:SER:OG	1:A:492:LEU:HB2	1.76	0.85
5:A:910:B12:H531	5:A:910:B12:C55	2.17	0.74
5:A:910:B12:H351	5:A:910:B12:C36	2.21	0.67
1:A:370:LEU:H	1:A:408:HIS:HD1	1.44	0.66
1:A:545:ARG:HD2	1:A:631:ARG:HH21	1.65	0.62
1:A:503:ARG:HB2	8:A:1167:HOH:O	2.00	0.60
1:A:606:ALA:N	2:A:904:GOL:O1	2.36	0.59
1:A:669:LYS:HE3	8:A:1184:HOH:O	2.05	0.56
1:A:192:THR:HA	1:A:224:THR:OG1	2.07	0.55
1:A:834:LEU:HB3	1:A:839:LEU:HD21	1.88	0.54
1:A:762:ASP:H	5:A:910:B12:H522	1.53	0.54
1:A:276:ARG:O	1:A:345:ARG:HD3	2.08	0.54
1:A:626:LEU:HD11	2:A:901:GOL:H11	1.90	0.54
1:A:353:VAL:HG12	1:A:460:ARG:HG2	1.91	0.53
1:A:605:ASP:HA	2:A:904:GOL:O1	2.09	0.53
1:A:252:VAL:O	1:A:278:VAL:HA	2.09	0.52
5:A:910:B12:H533	5:A:910:B12:H482	1.92	0.51
5:A:910:B12:N24	6:A:912:TME:H31	2.25	0.51
1:A:670:ARG:HD2	8:A:1114:HOH:O	2.11	0.50
1:A:233:VAL:HG23	1:A:356:LEU:HD11	1.94	0.49
1:A:455:LYS:HE2	2:A:906:GOL:H12	1.94	0.49
1:A:629:ASP:OD1	2:A:901:GOL:O3	2.29	0.49
1:A:159:VAL:HG22	2:A:908:GOL:H31	1.94	0.48
1:A:759:ASP:HA	5:A:910:B12:H452	1.79	0.48
1:A:177:ALA:HB2	1:A:206:VAL:HG22	1.95	0.48
3:A:902:HCS:HG2	8:A:1109:HOH:O	2.13	0.48
1:A:537:ILE:HB	2:A:901:GOL:H12	1.95	0.47
1:A:626:LEU:HD12	2:A:901:GOL:H32	1.96	0.47
1:A:159:VAL:O	2:A:908:GOL:H12	2.15	0.47
1:A:233:VAL:CG2	1:A:356:LEU:HD11	2.44	0.47
1:A:263:MET:O	1:A:267:VAL:HG23	2.14	0.47
1:A:798:HIS:O	1:A:799:LYS:C	2.58	0.47
1:A:614:GLN:NE2	8:A:1009:HOH:O	2.47	0.47
1:A:277:PHE:HA	1:A:345:ARG:HD2	1.98	0.46
5:A:910:B12:H3	5:A:910:B12:N29	2.31	0.46
1:A:604:VAL:O	2:A:904:GOL:H32	2.15	0.46
1:A:135:ARG:NH2	1:A:138:ARG:HG2	2.31	0.45
1:A:503:ARG:HD3	8:A:1167:HOH:O	2.16	0.45
1:A:803:VAL:HG23	1:A:830:LEU:HD23	1.97	0.45
1:A:400:ARG:HG3	8:A:1174:HOH:O	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:503:ARG:NH1	1:A:553:LEU:HD13	2.32	0.45
1:A:857:ALA:CB	5:A:910:B12:HM62	2.47	0.44
1:A:384:ARG:HG3	1:A:388:PHE:CZ	2.52	0.44
1:A:233:VAL:HG22	8:A:1035:HOH:O	2.18	0.43
5:A:910:B12:H3	5:A:910:B12:H291	1.82	0.43
5:A:910:B12:H492	5:A:910:B12:C47	2.49	0.43
1:A:263:MET:HE2	8:A:1104:HOH:O	2.18	0.43
5:A:910:B12:H492	5:A:910:B12:H471	2.01	0.42
1:A:456:TYR:CD2	2:A:908:GOL:H2	2.54	0.42
1:A:873:HIS:O	1:A:874:ALA:C	2.62	0.42
1:A:492:LEU:HD12	1:A:531:ASP:HB3	2.02	0.42
1:A:767:LEU:HA	1:A:767:LEU:HD23	1.89	0.42
1:A:376:ASN:OD1	1:A:378:THR:HB	2.20	0.41
1:A:632:LYS:HB2	1:A:632:LYS:HE3	1.84	0.41
1:A:1:MET:HB3	1:A:4:VAL:HG22	2.03	0.41
1:A:760:VAL:HG22	5:A:910:B12:C43	2.51	0.41
1:A:830:LEU:HG	1:A:831:PRO:HD2	2.02	0.41
1:A:166:TRP:HB3	2:A:907:GOL:H12	2.03	0.40
1:A:626:LEU:CD1	2:A:901:GOL:H11	2.50	0.40
5:A:910:B12:H262	5:A:910:B12:H91	1.86	0.40
1:A:1:MET:HG2	1:A:4:VAL:HG13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	858/874 (98%)	835 (97%)	21 (2%)	2 (0%)	44 54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	465	SER
1	A	747	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	679/687 (99%)	658 (97%)	21 (3%)	35	50

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	ASP
1	A	61	GLN
1	A	62	LYS
1	A	86	ARG
1	A	138	ARG
1	A	187	LEU
1	A	192	THR
1	A	206	VAL
1	A	213	VAL
1	A	283	ASN
1	A	337	VAL
1	A	378	THR
1	A	461	VAL
1	A	465	SER
1	A	524	ARG
1	A	539	GLN
1	A	573	ASN
1	A	580	PRO
1	A	610	LEU
1	A	743	LYS
1	A	811	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	GLN
1	A	273	HIS
1	A	352	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 3 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	907	-	5,5,5	0.33	0	5,5,5	0.53	0
6	TME	A	912	-	2,2,2	0.39	0	1,1,1	0.37	0
3	HCS	A	902	4	6,7,7	1.51	1 (16%)	5,8,8	2.12	3 (60%)
2	GOL	A	909	-	5,5,5	0.24	0	5,5,5	0.51	0
5	B12	A	910	1	91,101,101	0.79	2 (2%)	140,166,166	1.59	16 (11%)
2	GOL	A	905	-	5,5,5	0.30	0	5,5,5	0.60	0
2	GOL	A	901	-	5,5,5	0.33	0	5,5,5	0.69	0
2	GOL	A	908	-	5,5,5	0.50	0	5,5,5	1.02	0
2	GOL	A	904	-	5,5,5	0.22	0	5,5,5	0.79	0
2	GOL	A	906	-	5,5,5	0.58	0	5,5,5	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	907	-	-	2/4/4/4	-
3	HCS	A	902	4	-	2/7/7/7	-
2	GOL	A	909	-	-	2/4/4/4	-
5	B12	A	910	1	-	10/52/223/223	0/3/11/11
2	GOL	A	905	-	-	2/4/4/4	-
2	GOL	A	901	-	-	2/4/4/4	-
2	GOL	A	908	-	-	2/4/4/4	-
2	GOL	A	904	-	-	1/4/4/4	-
2	GOL	A	906	-	-	4/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	910	B12	C14-N23	3.80	1.40	1.35
3	A	902	HCS	O-C	3.24	1.31	1.22
5	A	910	B12	C16-C15	-2.66	1.37	1.44

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	910	B12	C19-N24-C16	9.88	118.09	107.29
5	A	910	B12	C19-C1-N21	5.54	107.85	102.14
5	A	910	B12	C18-C19-N24	-3.72	96.73	102.33
5	A	910	B12	C20-C1-N21	-3.42	104.61	110.26
5	A	910	B12	C41-C8-C9	3.36	117.04	111.19
5	A	910	B12	C9-N22-C6	3.36	109.32	105.28
5	A	910	B12	O6R-C1R-N1B	-3.29	104.38	108.75
3	A	902	HCS	CG-CB-CA	2.88	117.84	113.14
5	A	910	B12	O2-C3R-C4R	-2.73	100.39	110.03
5	A	910	B12	O5-P-O4	2.72	125.12	112.44
5	A	910	B12	C26-C27-N29	2.70	124.86	116.49
5	A	910	B12	C7-C8-C9	-2.56	97.65	100.89
3	A	902	HCS	OXT-C-O	-2.53	118.35	124.08
5	A	910	B12	C3P-C2P-C1P	-2.52	106.55	111.42
5	A	910	B12	C5-C4-N21	2.46	128.14	124.19
5	A	910	B12	C26-C2-C1	2.45	113.78	110.00
5	A	910	B12	C17-C18-C19	2.44	106.07	102.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	910	B12	O3-C2P-C1P	2.17	111.24	106.94
3	A	902	HCS	CB-CG-SD	2.14	115.97	113.74

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	901	GOL	C1-C2-C3-O3
2	A	905	GOL	C1-C2-C3-O3
2	A	908	GOL	O1-C1-C2-C3
2	A	909	GOL	C1-C2-C3-O3
3	A	902	HCS	N-CA-CB-CG
5	A	910	B12	C3-C30-C31-C32
5	A	910	B12	C18-C60-C61-N62
5	A	910	B12	C18-C60-C61-O63
2	A	906	GOL	O1-C1-C2-C3
2	A	907	GOL	C1-C2-C3-O3
2	A	905	GOL	O2-C2-C3-O3
2	A	906	GOL	O1-C1-C2-O2
2	A	907	GOL	O2-C2-C3-O3
5	A	910	B12	O6R-C4R-C5R-O8R
2	A	901	GOL	O2-C2-C3-O3
2	A	908	GOL	O1-C1-C2-O2
3	A	902	HCS	C-CA-CB-CG
2	A	909	GOL	O2-C2-C3-O3
5	A	910	B12	C48-C49-C50-N52
5	A	910	B12	C38-C37-C7-C36
5	A	910	B12	C2-C26-C27-N29
2	A	904	GOL	O2-C2-C3-O3
5	A	910	B12	C2-C26-C27-O28
2	A	906	GOL	C1-C2-C3-O3
5	A	910	B12	C48-C49-C50-O51
2	A	906	GOL	O2-C2-C3-O3
5	A	910	B12	C42-C41-C8-C9

There are no ring outliers.

8 monomers are involved in 29 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	907	GOL	1	0
6	A	912	TME	1	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	902	HCS	1	0
5	A	910	B12	15	0
2	A	901	GOL	5	0
2	A	908	GOL	3	0
2	A	904	GOL	3	0
2	A	906	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand HCS A 902



Ligand B12 A 910



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	862/874 (98%)	0.08	19 (2%) 62 65	49, 65, 93, 149	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	539	GLN	4.2
1	A	2	VAL	3.7
1	A	4	VAL	3.5
1	A	634	GLY	3.4
1	A	743	LYS	3.1
1	A	541	ASP	2.9
1	A	767	LEU	2.9
1	A	654	LYS	2.8
1	A	61	GLN	2.5
1	A	873	HIS	2.5
1	A	874	ALA	2.4
1	A	1	MET	2.4
1	A	540	GLY	2.4
1	A	744	LYS	2.4
1	A	830	LEU	2.3
1	A	543	GLU	2.3
1	A	283	ASN	2.2
1	A	872	GLY	2.1
1	A	650	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	901	6/6	0.85	0.19	75,86,111,123	0
3	HCS	A	902	8/8	0.85	0.20	92,106,116,138	0
2	GOL	A	906	6/6	0.90	0.17	76,86,98,99	0
2	GOL	A	907	6/6	0.91	0.18	57,92,99,127	0
2	GOL	A	909	6/6	0.92	0.15	65,75,85,88	0
2	GOL	A	908	6/6	0.92	0.14	64,79,100,110	0
2	GOL	A	905	6/6	0.94	0.11	53,60,75,85	0
2	GOL	A	904	6/6	0.94	0.14	64,71,86,99	0
6	TME	A	912	3/3	0.95	0.20	66,66,67,76	0
7	K	A	913	1/1	0.95	0.11	71,71,71,71	0
5	B12	A	910	91/91	0.96	0.09	43,62,76,88	0
4	ZN	A	903	1/1	0.97	0.07	77,77,77,77	0
4	ZN	A	911	1/1	0.99	0.04	58,58,58,58	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HCS A 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TME A 912:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around B12 A 910:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around ZN A 903:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 911:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.