

Full wwPDB EM Validation Report (i)

Mar 6, 2025 – 04:23 PM EST

PDB ID : 9CCJ

EMDB ID : EMD-45444

Title: Dissecting human monoclonal antibody responses from mRNA and protein-

based booster vaccinations against XBB1.5 SARS-CoV-2

Authors: Bajic, G.; Civljak, A.

Deposited on : 2024-06-21

Resolution : 2.37 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev117

Mogul : 2022.3.0, CSD as543be (2022)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : 1.9.13

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.41.4

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	Н	222	79%	15% 6%					
2	L	212	85%	10% •					
3	A	1229	21% • 78%						
4	В	3	67% 67%	33%					
5	С	2	100%						



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called M2 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	Н	209	Total 1577	C 1003	N 259	O 309	S 6	0	0

• Molecule 2 is a protein called M2 Fab Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	т	203	Total	С	N	О	S	0	0
2	Ъ	203	1512	943	247	316	6	U	U

• Molecule 3 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	A	276	Total 2222	C 1438	N 364	O 413	S 7	0	0

There are 93 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	ILE	THR	variant	UNP P0DTC2
A	?	-	LEU	variant	UNP P0DTC2
A	?	-	PRO	variant	UNP P0DTC2
A	?	-	PRO	variant	UNP P0DTC2
A	24	SER	ALA	variant	UNP P0DTC2
A	80	ALA	VAL	variant	UNP P0DTC2
A	139	ASP	GLY	variant	UNP P0DTC2
A	?	-	TYR	variant	UNP P0DTC2
A	142	GLN	HIS	variant	UNP P0DTC2
A	179	GLU	GLN	variant	UNP P0DTC2
A	209	GLU	VAL	variant	UNP P0DTC2
A	248	VAL	GLY	variant	UNP P0DTC2
A	303	ILE	THR	conflict	UNP P0DTC2
A	335	HIS	GLY	variant	UNP P0DTC2
A	342	THR	ARG	variant	UNP P0DTC2

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
A	364	ILE	LEU	variant	UNP P0DTC2
A	367	PHE	SER	variant	UNP P0DTC2
A	369	PRO	SER	variant	UNP P0DTC2
A	371	PHE	SER	variant	UNP P0DTC2
A	372	ALA	THR	variant	UNP P0DTC2
A	401	ASN	ASP	variant	UNP P0DTC2
A	404	SER	ARG	variant	UNP P0DTC2
A	413	ASN	LYS	variant	UNP P0DTC2
A	436	LYS	ASN	variant	UNP P0DTC2
A	441	PRO	VAL	variant	UNP P0DTC2
A	442	SER	GLY	variant	UNP P0DTC2
A	451	PHE	LEU	conflict	UNP P0DTC2
A	452	LEU	PHE	conflict	UNP P0DTC2
A	456	LYS	ASN	variant	UNP P0DTC2
A	471	VAL	ALA	conflict	UNP P0DTC2
A	473	ASN	SER	variant	UNP P0DTC2
A	474	LYS	THR	variant	UNP P0DTC2
A	480	ALA	GLU	variant	UNP P0DTC2
A	482	PRO	PHE	variant	UNP P0DTC2
A	486	SER	PHE	variant	UNP P0DTC2
A	494	ARG	GLN	variant	UNP P0DTC2
A	497	TYR	ASN	variant	UNP P0DTC2
A	501	HIS	TYR	variant	UNP P0DTC2
A	610	GLY	ASP	variant	UNP P0DTC2
A	651	TYR	HIS	variant	UNP P0DTC2
A	675	LYS	ASN	variant	UNP P0DTC2
A	677	HIS	PRO	variant	UNP P0DTC2
A	678	GLY	ARG	engineered mutation	UNP P0DTC2
A	679	SER	ARG	engineered mutation	UNP P0DTC2
A	681	SER	ARG	engineered mutation	UNP P0DTC2
A	760	LYS	ASN	variant	UNP P0DTC2
A	792	TYR	ASP	variant	UNP P0DTC2
A	813	PRO	PHE	engineered mutation	UNP P0DTC2
A	888	PRO	ALA	engineered mutation	UNP P0DTC2
A	895	PRO	ALA	engineered mutation	UNP P0DTC2
A	938	PRO	ALA	engineered mutation	UNP P0DTC2
A	950	HIS	GLN	variant	UNP P0DTC2
A	965	LYS	ASN	variant	UNP P0DTC2
A	982	PRO	LYS	engineered mutation	UNP P0DTC2
A	983	PRO	VAL	engineered mutation	UNP P0DTC2
A	1205	GLY	-	expression tag	UNP P0DTC2
A	1206	SER	-	expression tag	UNP P0DTC2

Continued on next page...



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
A	1207	GLY	-	expression tag	UNP P0DTC2
A	1208	TYR	-	expression tag	UNP P0DTC2
A	1209	ILE	-	expression tag	UNP P0DTC2
A	1210	PRO	-	expression tag	UNP P0DTC2
A	1211	GLU	-	expression tag	UNP P0DTC2
A	1212	ALA	-	expression tag	UNP P0DTC2
A	1213	PRO	-	expression tag	UNP P0DTC2
A	1214	ARG	-	expression tag	UNP P0DTC2
A	1215	ASP	-	expression tag	UNP P0DTC2
A	1216	GLY	-	expression tag	UNP P0DTC2
A	1217	GLN	-	expression tag	UNP P0DTC2
A	1218	ALA	-	expression tag	UNP P0DTC2
A	1219	TYR	-	expression tag	UNP P0DTC2
A	1220	VAL	-	expression tag	UNP P0DTC2
A	1221	ARG	-	expression tag	UNP P0DTC2
A	1222	LYS	-	expression tag	UNP P0DTC2
A	1223	ASP	-	expression tag	UNP P0DTC2
A	1224	GLY	-	expression tag	UNP P0DTC2
A	1225	GLU	-	expression tag	UNP P0DTC2
A	1226	TRP	-	expression tag	UNP P0DTC2
A	1227	VAL	-	expression tag	UNP P0DTC2
A	1228	LEU	-	expression tag	UNP P0DTC2
A	1229	LEU	-	expression tag	UNP P0DTC2
A	1230	SER	-	expression tag	UNP P0DTC2
A	1231	THR	-	expression tag	UNP P0DTC2
A	1232	PHE	-	expression tag	UNP P0DTC2
A	1233	LEU	-	expression tag	UNP P0DTC2
A	1234	GLY	-	expression tag	UNP P0DTC2
A	1235	ARG	-	expression tag	UNP P0DTC2
A	1236	SER	-	expression tag	UNP P0DTC2
A	1237	LEU	-	expression tag	UNP P0DTC2
A	1238	GLU	-	expression tag	UNP P0DTC2
A	1239	VAL	-	expression tag	UNP P0DTC2
A	1240	LEU	-	expression tag	UNP P0DTC2
A	1241	PHE	-	expression tag	UNP P0DTC2
A	1242	GLN	-	expression tag	UNP P0DTC2

 \bullet Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.






Mol	Chain	Residues	Atoms				AltConf	Trace
4	В	3	Total 39	C 22	N 2	O 15	0	0

• Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			AltConf	Trace	
5	С	2	Total 28	C 16	N 2	O 10	0	0

• Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	A	AltConf			
6	Λ	1	Total	С	N	О	0
0	A	1	14	8	1	5	U
6	Λ	1	Total	С	N	О	0
0	A	1	14	8	1	5	U

Continued on next page...




Mol	Chain	Residues	Atoms				AltConf
6	Λ	1	Total	С	N	О	0
0	A	1	14	8	1	5	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: M2 Fab Heavy Chain





PHE	ASN	LEU	THR	GLY	GLY	LEU	THR	SER	ASN	LYS	LYS	LEG	PRO	PHE	G GLN	PHE	GLY	ARG	ILE	ALA	ASP	THR	ASP	VAL	ARG	ASP	GLN	THR	OTO	ILE	ASP	ILE	THR	CYS	SER	GLY	GLY	VAL	VAL	ILE	THR	GLY	IHK
ASN	用品	ASN	GLN	VAL ALA	VAL	TYR	GLN	GLY VAL	ASN	CYS	E E	VAL	PRO	VAL	ALA II.E	HIS	ALA	GLN	LEU	THR	25 15 15 15 15 15 15 15 15 15 15 15 15 15	TE TE	ARG	TYR	SER	E 2	SER	ASN	VAL	GLN	ARG	ALA	GLY	LEU	ILE GIV	ALA	GLU	TYR	VAL	ASN	SER	GLU	CYS
ASP	ILE	ILE	GLY	ALA GLY	ILE	ALA	SER	TYK	THR	GLN	THR	SER	HIS	GLY	AI.A	SER	SER	VAL	SER	GLN	NEK T.F.	ILE	ALA	THR	MET	SER	GLY	ALA	ASN	SER	ALA	TYR	SER	ASN	SER	ALA	ILE	PRO	ASN	PHE	THR	SER	VAL
THR	THR	ILE	LEU	PRO VAT.	SER	THR	LYS	SER	VAL	ASP	CYS	MET	TYR	ILE	GI.Y	ASP	SER	GLU	CYS	SER	ASN	PE	LEU	TYR	GLY	SER	CYS	THR	LEU	LYS	ALA	LEU	THR GL.Y	ILE	ALA	GLU	GLN	ASP	ASN	THR	GLN	VAL	PHE
ALA	GLN	LYS	GLN	ILE	LYS	PRO	PRO	LYS	TYR	PHE	GLY	PHE	ASN	PHE	GLN	ILE	LEU	ASP	PRO	SER	PRO	SER	LYS	SER	PRO	ILE	ASP	LEU	PHE	ASN	VAL	THR	LEU	ASP	ALA	PHE	ILE	LYS	TYR	GLY	ASP	LEU	GLY
ASP	ILE	ALA	ARG	ASP 1.EII	ITE	ALA	GLN	LYS	ASN	GLY	LEU	VAL	LEU	PRO	LEU	LEU	THE	GLU	MET	ILE	ALA	TYR	THR	ALA	LEU	LEU	GLY	THR	H H	SER	TRP	THR	PHE GLY	ALA	GLY	ALA	LEU	GLN	PRO	PHE	PRO	GLN	MEI
ALA	TYR	PHE	ASN	GLY TI.E	GLY	THR	GLN	VAL	LEU	TYR	GLU	GLN	LYS	LEU	AT.A	ASN	GLN	ASN	SER	ALA	크 ^ - - -	LYS	ILE	ASP	SER	LEU	SER	THR	SER	ALA	GLY	LYS	CILN	ASP	VAL	ASN	HIS	ASN	GLN	ALA	LEU	THR	רבּח
VAL	LYS	LEU	SER	SER 1.YS	PHE	ALA	ILE	SER	VAL	LEU	ASN	ILE	LEU	SER	LEU	ASP	PRO	GLU	ALA	GLU	VAL GI.N	ILE	ASP	LEU	ILE	THR	ARG	LEU	SER	LEU	THR	TYR	VAL	GLN	GLN	ILE	ARG	ALA	GLU	ILE	ARG	SER	ALA
ASN	LEU	ALA	THR	LYS	SER	CYS	VAL	CEU	GLN	SER	LYS	VAL	ASP	PHE	GI.Y	LYS	GLY	HIS	LEU	MET	DHE	PRO	GLN	ALA	PRO	HIS	VAL	VAL	LEU	HIS	THR	TYR	VAL	ALA	GLN	LYS	ASN	PHE	H H	ALA	PRO	ILE	CIN
HIS	ASP	LYS	ALA	HIS	PRO	GLU	GLY	VAL	VAL	SER	ASN	THR	HIS	TRP	VAT.	THR	GLN	ASN	PHE	TYR	GLO	GLN	ILE	THE	TH.	ASP	THR	PHE	SER	GLY	CYS	ASP	VAL VAI.	111	GLY	VAL	ASN	ASN	VAL	TYR	ASP	LEU	STIN
PRO	GLU	ASP	SER	PHE	GLU	LEU	ASP	LYS	PHE	LYS	ASN	THR	SER	PRO	VAI.	ASP	LEU	GLY	ILE	SER	7 T. T.	ASN	ALA	VAL	VAL	ASN	GLN	LYS	ILE	ASP	LEU	ASN	GLU VAI.	ALA	LYS	LEU	ASN	GLU	LEU	ILE	ASP	GLN	GLU
LEU	GLY	TYR	GLU	GLN GLY	SER	TYR	ILE	GLU	ALA	PRO	ARG	GLY	GLN	ALA	VAT.	ARG	LYS	GLY	GLU	TRP	VAL	LEU	SER	PHE	LEU	GLY	SER	LEU	VAL	LEU	GLN												

 $\bullet \ \, \text{Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose} \\$

Chain B: 67% 33%



 \bullet Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 100%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	298206	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	49.883	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.829	Depositor
Minimum map value	-0.002	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.010	Depositor
Recommended contour level	0.1	Depositor
Map size (Å)	422.4, 422.4, 422.4	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.825, 0.825, 0.825	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles				
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5			
1	Н	0.29	0/1617	0.51	0/2210			
2	L	0.29	0/1547	0.47	0/2114			
3	A	0.29	0/2275	0.50	0/3097			
All	All	0.29	0/5439	0.50	0/7421			

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Н	1577	0	1558	20	0
2	L	1512	0	1452	14	0
3	A	2222	0	2149	12	0
4	В	39	0	34	0	0
5	С	28	0	25	0	0
6	A	42	0	39	0	0
All	All	5420	0	5257	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.



All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

		Interatomic	Clash
Atom-1	Atom-2	distance (\mathring{A})	overlap(A)
2:L:110:LYS:HZ1	2:L:198:GLU:HG3	1.53	0.73
2:L:38:LYS:NZ	2:L:80:MET:O	2.25	0.69
1:H:156:VAL:HG22	1:H:202:VAL:HG22	1.84	0.60
3:A:191:LYS:HD3	3:A:193:ILE:HD12	1.86	0.58
3:A:90:ALA:HB3	3:A:262:TYR:HB2	1.86	0.57
1:H:146:VAL:HB	1:H:182:LEU:HB3	1.88	0.54
3:A:119:ASN:O	3:A:121:THR:N	2.41	0.53
1:H:163:LEU:HD21	1:H:186:VAL:HG11	1.91	0.52
2:L:110:LYS:NZ	2:L:198:GLU:HG3	2.24	0.50
1:H:99:PRO:HG2	1:H:103:ARG:HE	1.75	0.50
1:H:9:LEU:HB2	1:H:151:PRO:HG3	1.94	0.50
2:L:132:LEU:HB2	2:L:178:LEU:HB3	1.93	0.50
1:H:143:GLY:HA2	1:H:158:TRP:CZ2	2.49	0.48
2:L:1:SER:OG	2:L:2:SER:N	2.44	0.47
2:L:151:ASP:OD1	2:L:190:SER:N	2.48	0.47
2:L:132:LEU:HD12	2:L:178:LEU:HD23	1.97	0.47
3:A:30:THR:HG22	3:A:30:THR:O	2.15	0.46
1:H:167:VAL:HG22	1:H:186:VAL:HG22	1.96	0.46
3:A:127:VAL:HG21	3:A:227:ILE:HG21	1.98	0.46
2:L:82:GLU:HG3	2:L:103:LEU:O	2.16	0.46
1:H:201:ASN:ND2	1:H:212:ASP:OD1	2.43	0.46
2:L:21:THR:HG23	2:L:69:THR:HG23	1.97	0.45
1:H:97:ARG:CG	1:H:106:TYR:HB2	2.47	0.45
3:A:26:THR:HG23	3:A:59:VAL:HG23	1.98	0.45
3:A:30:THR:HG21	3:A:216:PHE:CZ	2.52	0.45
3:A:212:LEU:HA	3:A:213:PRO:HD3	1.85	0.44
3:A:41:ARG:O	3:A:279:GLY:HA2	2.17	0.44
1:H:143:GLY:HA3	1:H:184:SER:O	2.18	0.44
2:L:47:ILE:HD12	2:L:72:LEU:HD13	1.99	0.44
1:H:204:HIS:CD2	1:H:206:PRO:HD2	2.53	0.43
2:L:181:THR:HG23	2:L:184:GLN:H	1.82	0.43
3:A:102:ILE:O	3:A:234:PHE:HA	2.18	0.43
2:L:117:LEU:HD21	2:L:132:LEU:HD22	2.01	0.42
1:H:97:ARG:HD3	1:H:106:TYR:HB2	2.02	0.42
2:L:93:SER:OG	3:A:157:SER:O	2.36	0.42
1:H:148:ASP:HB3	1:H:179:LEU:HD13	2.02	0.42
1:H:155:THR:OG1	1:H:203:ASN:HB3	2.19	0.42
1:H:190:SER:O	1:H:193:LEU:HG	2.20	0.42
1:H:123:PRO:HD2	1:H:209:THR:HG21	2.02	0.41
1:H:34:TRP:CZ3	1:H:97:ARG:HB3	2.56	0.41

Continued on next page...



Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:H:51:ILE:HB	1:H:69:MET:HE2	2.00	0.41
1:H:27:ILE:HA	1:H:34:TRP:CZ2	2.54	0.41
2:L:138:ASP:OD1	2:L:167:GLN:NE2	2.54	0.41
3:A:144:ASN:O	3:A:146:LYS:HG3	2.21	0.40
1:H:97:ARG:HG3	1:H:105:ASP:OD1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	Н	205/222~(92%)	196 (96%)	9 (4%)	0	100	100
2	L	199/212 (94%)	192 (96%)	7 (4%)	0	100	100
3	A	270/1229 (22%)	260 (96%)	10 (4%)	0	100	100
All	All	674/1663 (40%)	648 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	tiles
1	Н	182/193 (94%)	182 (100%)	0	100	100
2	L	173/182 (95%)	173 (100%)	0	100	100

Continued on next page...



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
3	A	249/1071 (23%)	249 (100%)	0	100	100
All	All	604/1446 (42%)	604 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	B	Bond lengths I				gles
MIOI	Туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	PCA	A	14	3	7,8,9	0.62	0	9,10,12	1.26	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PCA	A	14	3	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	14	PCA	CB-CA-N	2.25	109.44	103.24





Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
3	A	14	PCA	CB-CG-CD	2.08	107.62	104.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trus	Chain	Res	Res Link Bond lengths			Bond angles			
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	В	1	4,3	14,14,15	0.74	0	17,19,21	0.85	0
4	NAG	В	2	4	14,14,15	0.73	0	17,19,21	0.89	0
4	BMA	В	3	4	11,11,12	0.84	0	15,15,17	2.04	3 (20%)
5	NAG	С	1	3,5	14,14,15	0.76	0	17,19,21	0.97	2 (11%)
5	NAG	С	2	5	14,14,15	0.71	0	17,19,21	0.86	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	В	1	4,3	-	0/6/23/26	0/1/1/1
4	NAG	В	2	4	-	0/6/23/26	0/1/1/1
4	BMA	В	3	4	-	0/2/19/22	0/1/1/1
5	NAG	С	1	3,5	-	0/6/23/26	0/1/1/1
5	NAG	С	2	5	-	2/6/23/26	0/1/1/1

There are no bond length outliers.



All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	В	3	BMA	C1-O5-C5	5.98	120.20	112.19
4	В	3	BMA	C3-C4-C5	2.61	114.96	110.23
5	С	1	NAG	O5-C1-C2	-2.31	107.71	111.29
4	В	3	BMA	C2-C3-C4	2.23	114.78	110.86
5	С	1	NAG	O4-C4-C3	-2.22	105.15	110.38
5	С	2	NAG	C2-N2-C7	2.14	125.76	122.90

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	С	2	NAG	C8-C7-N2-C2
5	С	2	NAG	O7-C7-N2-C2


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	nd leng	ths	Bond angles		
WIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	A	1302	3	14,14,15	0.70	0	17,19,21	0.90	0
6	NAG	A	1303	3	14,14,15	0.73	0	17,19,21	0.88	1 (5%)
6	NAG	A	1301	3	14,14,15	0.81	0	17,19,21	0.94	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	1302	3	-	0/6/23/26	0/1/1/1
6	NAG	A	1303	3	-	3/6/23/26	0/1/1/1
6	NAG	A	1301	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$	
6	A	1303	NAG	C2-N2-C7	2.11	125.73	122.90	

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	1303	NAG	C8-C7-N2-C2
6	A	1303	NAG	O7-C7-N2-C2
6	A	1303	NAG	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-45444. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map



The images above show the map projected in three orthogonal directions.



6.2 Central slices (i)

6.2.1 Primary map








Y Index: 256




Z Index: 256


6.2.2 Raw map



X Index: 256



Y Index: 256




Z Index: 256

The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map



6.3.2 Raw map





The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map









6.4.2 Raw map








The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_45444_msk_1.map (i)






7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 69 nm^3 ; this corresponds to an approximate mass of 62 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.422 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.422 $\rm \AA^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)				
resolution estimate (A)	0.143	0.5	Half-bit		
Reported by author	2.37	-	-		
Author-provided FSC curve	-	-	-		
Unmasked-calculated*	3.72	7.91	3.85		

^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.72 differs from the reported value 2.37 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-45444 and PDB model 9CCJ. Per-residue inclusion information can be found in section 3 on page 8.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).



9.4 Atom inclusion (i)



At the recommended contour level, 95% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8870	0.6230
A	0.9040	0.6510
В	0.3330	0.4590
С	0.7140	0.5260
Н	0.8720	0.6060
L	0.8960	0.6050



