



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 11, 2025 – 04:22 PM EDT

PDB ID : 9D57 / pdb_00009d57
Title : iGABASnFR2 fluorescent GABA sensor in complex with GABA
Authors : Zhang, Y.; Looger, L.L.
Deposited on : 2024-08-13
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	564	<div> <div>85%</div> <div>10% . .</div> </div>
1	B	564	<div> <div>%</div> <div>85%</div> <div>9% . .</div> </div>
1	C	564	<div> <div>%</div> <div>84%</div> <div>9% . 6%</div> </div>
1	D	564	<div> <div>86%</div> <div>9% . .</div> </div>
1	E	564	<div> <div>2%</div> <div>86%</div> <div>8% . .</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	564	<div><div><div>%</div><div><div></div></div><div>84%</div><div>10% • 5%</div></div></div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 24592 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC transporter substrate-binding protein, Circular permuted superfolder GFP fusion.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	544	Total	C	N	O	S	0	0	0
			4216	2692	712	800	12			
1	B	539	Total	C	N	O	S	0	0	0
			4063	2590	687	775	11			
1	D	540	Total	C	N	O	S	0	0	0
			4114	2623	695	785	11			
1	F	538	Total	C	N	O	S	0	0	0
			4059	2596	680	773	10			
1	C	532	Total	C	N	O	S	0	0	0
			4021	2566	677	769	9			
1	E	539	Total	C	N	O	S	0	0	0
			4045	2583	688	763	11			

- Molecule 2 is GAMMA-AMINO-BUTANOIC ACID (CCD ID: ABU) (formula: $C_4H_9NO_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			7	4	1	2		
2	B	1	Total	C	N	O	0	0
			7	4	1	2		
2	D	1	Total	C	N	O	0	0
			7	4	1	2		
2	F	1	Total	C	N	O	0	0
			7	4	1	2		
2	C	1	Total	C	N	O	0	0
			7	4	1	2		
2	E	1	Total	C	N	O	0	0
			7	4	1	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	O	0	0
			7	7		
3	B	8	Total	O	0	0
			8	8		
3	D	2	Total	O	0	0
			2	2		
3	F	4	Total	O	0	0
			4	4		
3	C	7	Total	O	0	0
			7	7		
3	E	4	Total	O	0	0
			4	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ABC transporter substrate-binding protein, Circular permuted superfolder GFP fusion



- Molecule 1: ABC transporter substrate-binding protein, Circular permuted superfolder GFP fusion





- Molecule 1: ABC transporter substrate-binding protein, Circular permuted superfolder GFP fusion









- Molecule 1: ABC transporter substrate-binding protein, Circular permuted superfolder GFP fusion




- Molecule 1: ABC transporter substrate-binding protein, Circular permuted superfolder GFP fusion



- Molecule 1: ABC transporter substrate-binding protein, Circular permuted superfolder GFP fusion





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.50Å 109.92Å 214.31Å 90.00° 100.25° 90.00°	Depositor
Resolution (Å)	47.39 – 2.60 47.39 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.3 (47.39-2.60) 93.7 (47.39-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.206 , 0.249 0.202 , 0.244	Depositor DCC
R_{free} test set	5396 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	52.7	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 29.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.020 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	24592	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CRO, ABU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/4294	1.14	15/5836 (0.3%)
1	B	0.59	0/4140	1.12	12/5649 (0.2%)
1	C	0.59	0/4100	1.13	18/5593 (0.3%)
1	D	0.59	0/4193	1.13	18/5716 (0.3%)
1	E	0.59	0/4123	1.13	15/5625 (0.3%)
1	F	0.60	0/4136	1.14	18/5640 (0.3%)
All	All	0.59	0/24986	1.13	96/34059 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	2
All	All	0	3

There are no bond length outliers.

The worst 5 of 96 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	351	ARG	CD-NE-CZ	9.21	137.30	124.40
1	C	201	GLU	CB-CG-CD	7.78	125.82	112.60
1	F	126	ASP	CA-CB-CG	7.58	120.18	112.60
1	A	42	ASP	CA-CB-CG	7.26	119.86	112.60
1	C	386	GLU	CB-CG-CD	7.25	124.92	112.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	140	ARG	Sidechain
1	D	140	ARG	Sidechain
1	D	351	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4216	0	4082	31	0
1	B	4063	0	3790	35	0
1	C	4021	0	3745	34	0
1	D	4114	0	3880	25	0
1	E	4045	0	3781	31	0
1	F	4059	0	3821	31	0
2	A	7	0	0	0	0
2	B	7	0	0	1	0
2	C	7	0	0	0	0
2	D	7	0	0	1	0
2	E	7	0	0	0	0
2	F	7	0	0	0	0
3	A	7	0	0	0	0
3	B	8	0	0	0	0
3	C	7	0	0	1	0
3	D	2	0	0	0	0
3	E	4	0	0	0	0
3	F	4	0	0	0	0
All	All	24592	0	23099	187	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 187 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:93:ILE:HD13	1:C:103:VAL:CG2	1.82	1.09
1:C:230:LEU:HB3	1:C:544:LEU:HD11	1.46	0.96
1:B:112:LEU:HD13	1:B:130:LEU:HD11	1.50	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:346:ASP:H	1:F:353:HIS:HE1	1.15	0.92
1:E:141:ALA:O	1:E:142:LEU:HD12	1.75	0.86

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	535/564 (95%)	520 (97%)	15 (3%)	0	100	100
1	B	530/564 (94%)	513 (97%)	17 (3%)	0	100	100
1	C	523/564 (93%)	510 (98%)	13 (2%)	0	100	100
1	D	531/564 (94%)	516 (97%)	15 (3%)	0	100	100
1	E	530/564 (94%)	517 (98%)	13 (2%)	0	100	100
1	F	529/564 (94%)	515 (97%)	14 (3%)	0	100	100
All	All	3178/3384 (94%)	3091 (97%)	87 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	439/464 (95%)	431 (98%)	8 (2%)	54	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	402/464 (87%)	394 (98%)	8 (2%)	50	74
1	C	398/464 (86%)	388 (98%)	10 (2%)	42	68
1	D	416/464 (90%)	407 (98%)	9 (2%)	47	72
1	E	397/464 (86%)	392 (99%)	5 (1%)	65	84
1	F	404/464 (87%)	394 (98%)	10 (2%)	42	68
All	All	2456/2784 (88%)	2406 (98%)	50 (2%)	50	74

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	134	LYS
1	C	135	THR
1	E	460	GLN
1	F	361	THR
1	F	423	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	279	GLN
1	F	419	ASN
1	E	228	GLN
1	F	353	HIS
1	F	552	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CRO	C	447	1	22,23,24	0.57	0	30,32,34	1.06	2 (6%)
1	CRO	F	447	1	22,23,24	0.63	0	30,32,34	1.02	2 (6%)
1	CRO	E	447	1	22,23,24	0.60	0	30,32,34	1.08	3 (10%)
1	CRO	D	447	1	22,23,24	0.55	0	30,32,34	1.01	3 (10%)
1	CRO	A	447	1	22,23,24	0.56	0	30,32,34	1.08	3 (10%)
1	CRO	B	447	1	22,23,24	0.53	0	30,32,34	1.05	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	C	447	1	-	2/12/31/32	0/2/2/2
1	CRO	F	447	1	-	2/12/31/32	0/2/2/2
1	CRO	E	447	1	-	2/12/31/32	0/2/2/2
1	CRO	D	447	1	-	2/12/31/32	0/2/2/2
1	CRO	A	447	1	-	2/12/31/32	0/2/2/2
1	CRO	B	447	1	-	2/12/31/32	0/2/2/2

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	447	CRO	C1-CA1-N1	-2.69	104.98	109.78
1	A	447	CRO	C1-CA1-N1	-2.64	105.06	109.78
1	E	447	CRO	C1-CA1-N1	-2.55	105.21	109.78
1	D	447	CRO	C1-CA1-N1	-2.54	105.23	109.78
1	B	447	CRO	C1-CA1-N1	-2.54	105.24	109.78

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	447	CRO	C3-CA3-N3-C2
1	D	447	CRO	C3-CA3-N3-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	C	447	CRO	C3-CA3-N3-C2
1	E	447	CRO	C3-CA3-N3-C2
1	B	447	CRO	C3-CA3-N3-C2

There are no ring outliers.

6 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	447	CRO	2	0
1	F	447	CRO	3	0
1	E	447	CRO	2	0
1	D	447	CRO	2	0
1	A	447	CRO	2	0
1	B	447	CRO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ABU	C	600	-	6,6,6	0.84	0	6,6,6	1.36	1 (16%)
2	ABU	F	600	-	6,6,6	0.62	0	6,6,6	1.24	2 (33%)
2	ABU	E	600	-	6,6,6	0.81	0	6,6,6	1.85	1 (16%)
2	ABU	D	600	-	6,6,6	0.60	0	6,6,6	1.39	2 (33%)
2	ABU	A	600	-	6,6,6	0.87	0	6,6,6	0.97	0
2	ABU	B	600	-	6,6,6	0.97	0	6,6,6	1.33	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ABU	C	600	-	-	0/4/4/4	-
2	ABU	F	600	-	-	1/4/4/4	-
2	ABU	E	600	-	-	2/4/4/4	-
2	ABU	D	600	-	-	2/4/4/4	-
2	ABU	A	600	-	-	2/4/4/4	-
2	ABU	B	600	-	-	2/4/4/4	-

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	600	ABU	CB-CG-C	3.99	124.91	114.51
2	C	600	ABU	CB-CG-C	2.40	120.78	114.51
2	B	600	ABU	CB-CG-C	2.27	120.43	114.51
2	D	600	ABU	O-C-CG	-2.20	116.11	123.09
2	F	600	ABU	O-C-CG	-2.13	116.34	123.09

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	600	ABU	CD-CB-CG-C
2	A	600	ABU	O-C-CG-CB
2	A	600	ABU	OXT-C-CG-CB
2	B	600	ABU	O-C-CG-CB
2	B	600	ABU	OXT-C-CG-CB


There are no ring outliers.



2 monomers are involved in 2 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	600	ABU	1	0
2	B	600	ABU	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	1
1	E	1
1	A	1
1	B	1
1	C	1
1	F	1

The worst 5 of 6 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	447:CRO	C3	448:VAL	N	1.76
1	E	447:CRO	C3	448:VAL	N	1.76
1	A	447:CRO	C3	448:VAL	N	1.75
1	B	447:CRO	C3	448:VAL	N	1.75

Continued on next page...

Continued from previous page...

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	447:CRO	C3	448:VAL	N	1.75

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	543/564 (96%)	-0.40	2 (0%) 89 86	27, 44, 69, 106	0
1	B	538/564 (95%)	-0.35	3 (0%) 85 83	26, 47, 78, 101	0
1	C	531/564 (94%)	-0.24	7 (1%) 74 70	30, 48, 82, 120	0
1	D	539/564 (95%)	-0.35	2 (0%) 89 86	31, 47, 71, 110	0
1	E	538/564 (95%)	-0.11	9 (1%) 69 64	25, 53, 86, 117	0
1	F	537/564 (95%)	-0.22	6 (1%) 77 74	28, 50, 76, 122	0
All	All	3226/3384 (95%)	-0.28	29 (0%) 81 77	25, 48, 79, 122	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	326	ASP	4.9
1	C	214	ALA	4.0
1	D	370	ASP	3.8
1	E	370	ASP	3.7
1	F	122	SER	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRO	D	447	22/23	0.94	0.08	38,43,50,67	0
1	CRO	F	447	22/23	0.94	0.10	40,47,52,53	0
1	CRO	E	447	22/23	0.94	0.09	37,47,55,62	0
1	CRO	B	447	22/23	0.95	0.07	26,30,40,41	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	CRO	C	447	22/23	0.95	0.07	33,40,48,60	0
1	CRO	A	447	22/23	0.95	0.07	30,35,42,52	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ABU	E	600	7/7	0.91	0.11	31,32,37,37	0
2	ABU	C	600	7/7	0.93	0.10	37,43,45,48	0
2	ABU	B	600	7/7	0.94	0.10	40,47,50,60	0
2	ABU	F	600	7/7	0.94	0.10	33,35,38,39	0
2	ABU	A	600	7/7	0.95	0.09	30,39,41,44	0
2	ABU	D	600	7/7	0.97	0.07	27,33,34,34	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ABU E 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ABU C 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ABU B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ABU F 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ABU A 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ABU D 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.