



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 20, 2023 – 10:31 PM JST

PDB ID : 7DQD
Title : Crystal structure of the AMP-PNP-bound mutant A(S23C)3B(N64C)3 complex from enterococcus hirae V-ATPase
Authors : Maruyama, S.; Suzuki, K.; Mizutani, K.; Imai, F.L.; Ishizuka-Katsura, Y.; Shirouzu, M.; Murata, T.
Deposited on : 2020-12-23
Resolution : 3.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1691 (3.46-3.30)
Clashscore	141614	1762 (3.46-3.30)
Ramachandran outliers	138981	1732 (3.46-3.30)
Sidechain outliers	138945	1731 (3.46-3.30)
RSRZ outliers	127900	1635 (3.46-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	600	
1	B	600	
1	C	600	
1	I	600	
1	J	600	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	600	
2	D	465	
2	E	465	
2	F	465	
2	L	465	
2	M	465	
2	N	465	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ANP	B	601	-	-	X	-
4	ANP	C	601	-	-	X	-
4	ANP	K	601	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 46492 atoms, of which 8 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type sodium ATPase catalytic subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	584	4448	2789	751	883	25	0	0	0
1	B	584	4443	2788	752	877	26	0	0	0
1	C	584	4274	2682	728	840	24	0	0	0
1	I	583	4291	2687	729	853	22	0	0	0
1	J	584	4422	2781	748	866	27	0	0	0
1	K	584	4435	2783	749	876	27	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLY	-	expression tag	UNP Q08636
A	-5	SER	-	expression tag	UNP Q08636
A	-4	SER	-	expression tag	UNP Q08636
A	-3	GLY	-	expression tag	UNP Q08636
A	-2	SER	-	expression tag	UNP Q08636
A	-1	SER	-	expression tag	UNP Q08636
A	0	GLY	-	expression tag	UNP Q08636
A	23	CYS	SER	engineered mutation	UNP Q08636
B	-6	GLY	-	expression tag	UNP Q08636
B	-5	SER	-	expression tag	UNP Q08636
B	-4	SER	-	expression tag	UNP Q08636
B	-3	GLY	-	expression tag	UNP Q08636
B	-2	SER	-	expression tag	UNP Q08636
B	-1	SER	-	expression tag	UNP Q08636
B	0	GLY	-	expression tag	UNP Q08636
B	23	CYS	SER	engineered mutation	UNP Q08636
C	-6	GLY	-	expression tag	UNP Q08636

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	SER	-	expression tag	UNP Q08636
C	-4	SER	-	expression tag	UNP Q08636
C	-3	GLY	-	expression tag	UNP Q08636
C	-2	SER	-	expression tag	UNP Q08636
C	-1	SER	-	expression tag	UNP Q08636
C	0	GLY	-	expression tag	UNP Q08636
C	23	CYS	SER	engineered mutation	UNP Q08636
I	-6	GLY	-	expression tag	UNP Q08636
I	-5	SER	-	expression tag	UNP Q08636
I	-4	SER	-	expression tag	UNP Q08636
I	-3	GLY	-	expression tag	UNP Q08636
I	-2	SER	-	expression tag	UNP Q08636
I	-1	SER	-	expression tag	UNP Q08636
I	0	GLY	-	expression tag	UNP Q08636
I	23	CYS	SER	engineered mutation	UNP Q08636
J	-6	GLY	-	expression tag	UNP Q08636
J	-5	SER	-	expression tag	UNP Q08636
J	-4	SER	-	expression tag	UNP Q08636
J	-3	GLY	-	expression tag	UNP Q08636
J	-2	SER	-	expression tag	UNP Q08636
J	-1	SER	-	expression tag	UNP Q08636
J	0	GLY	-	expression tag	UNP Q08636
J	23	CYS	SER	engineered mutation	UNP Q08636
K	-6	GLY	-	expression tag	UNP Q08636
K	-5	SER	-	expression tag	UNP Q08636
K	-4	SER	-	expression tag	UNP Q08636
K	-3	GLY	-	expression tag	UNP Q08636
K	-2	SER	-	expression tag	UNP Q08636
K	-1	SER	-	expression tag	UNP Q08636
K	0	GLY	-	expression tag	UNP Q08636
K	23	CYS	SER	engineered mutation	UNP Q08636

- Molecule 2 is a protein called V-type sodium ATPase subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	447	Total	C	N	O	S	0	0	0
			3383	2137	589	642	15			
2	E	453	Total	C	N	O	S	0	0	0
			3400	2152	592	643	13			
2	F	445	Total	C	N	O	S	0	0	0
			3295	2080	579	621	15			
2	L	447	Total	C	N	O	S	0	0	0
			3428	2167	591	655	15			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	453	Total	C	N	O	S	0	0	0
			3357	2119	586	639	13			
2	N	429	Total	C	N	O	S	0	0	0
			3162	1999	549	600	14			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	-6	GLY	-	expression tag	UNP Q08637
D	-5	SER	-	expression tag	UNP Q08637
D	-4	SER	-	expression tag	UNP Q08637
D	-3	GLY	-	expression tag	UNP Q08637
D	-2	SER	-	expression tag	UNP Q08637
D	-1	SER	-	expression tag	UNP Q08637
D	0	GLY	-	expression tag	UNP Q08637
D	64	CYS	ASN	engineered mutation	UNP Q08637
E	-6	GLY	-	expression tag	UNP Q08637
E	-5	SER	-	expression tag	UNP Q08637
E	-4	SER	-	expression tag	UNP Q08637
E	-3	GLY	-	expression tag	UNP Q08637
E	-2	SER	-	expression tag	UNP Q08637
E	-1	SER	-	expression tag	UNP Q08637
E	0	GLY	-	expression tag	UNP Q08637
E	64	CYS	ASN	engineered mutation	UNP Q08637
F	-6	GLY	-	expression tag	UNP Q08637
F	-5	SER	-	expression tag	UNP Q08637
F	-4	SER	-	expression tag	UNP Q08637
F	-3	GLY	-	expression tag	UNP Q08637
F	-2	SER	-	expression tag	UNP Q08637
F	-1	SER	-	expression tag	UNP Q08637
F	0	GLY	-	expression tag	UNP Q08637
F	64	CYS	ASN	engineered mutation	UNP Q08637
L	-6	GLY	-	expression tag	UNP Q08637
L	-5	SER	-	expression tag	UNP Q08637
L	-4	SER	-	expression tag	UNP Q08637
L	-3	GLY	-	expression tag	UNP Q08637
L	-2	SER	-	expression tag	UNP Q08637
L	-1	SER	-	expression tag	UNP Q08637
L	0	GLY	-	expression tag	UNP Q08637
L	64	CYS	ASN	engineered mutation	UNP Q08637
M	-6	GLY	-	expression tag	UNP Q08637
M	-5	SER	-	expression tag	UNP Q08637

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
M	-4	SER	-	expression tag	UNP Q08637
M	-3	GLY	-	expression tag	UNP Q08637
M	-2	SER	-	expression tag	UNP Q08637
M	-1	SER	-	expression tag	UNP Q08637
M	0	GLY	-	expression tag	UNP Q08637
M	64	CYS	ASN	engineered mutation	UNP Q08637
N	-6	GLY	-	expression tag	UNP Q08637
N	-5	SER	-	expression tag	UNP Q08637
N	-4	SER	-	expression tag	UNP Q08637
N	-3	GLY	-	expression tag	UNP Q08637
N	-2	SER	-	expression tag	UNP Q08637
N	-1	SER	-	expression tag	UNP Q08637
N	0	GLY	-	expression tag	UNP Q08637
N	64	CYS	ASN	engineered mutation	UNP Q08637

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	14	3	8	3	0	0

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
4	C	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
4	J	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
4	K	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	B	1	Total	Mg	0	0
			1	1		
5	C	1	Total	Mg	0	0
			1	1		
5	J	1	Total	Mg	0	0
			1	1		
5	K	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	1	Total	O	0	0
			1	1		


Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total O 1 1	0	0
6	C	1	Total O 1 1	0	0
6	E	1	Total O 1 1	0	0
6	I	2	Total O 2 2	0	0
6	L	1	Total O 1 1	0	0
6	M	1	Total O 1 1	0	0
6	N	4	Total O 4 4	0	0




• Molecule 1: V-type sodium ATPase catalytic subunit A







● Molecule 1: V-type sodium ATPase catalytic subunit A




● Molecule 1: V-type sodium ATPase catalytic subunit A







● Molecule 1: V-type sodium ATPase catalytic subunit A







• Molecule 2: V-type sodium ATPase subunit B




• Molecule 2: V-type sodium ATPase subunit B







• Molecule 2: V-type sodium ATPase subunit B




• Molecule 2: V-type sodium ATPase subunit B







• Molecule 2: V-type sodium ATPase subunit B



• Molecule 2: V-type sodium ATPase subunit B





4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	117.57Å 123.38Å 230.28Å 90.00° 90.07° 90.00°	Depositor
Resolution (Å)	49.37 – 3.38 49.36 – 3.38	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.37-3.38) 99.3 (49.36-3.38)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 3.40Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.261 , 0.286 0.261 , 0.288	Depositor DCC
R_{free} test set	4725 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	77.5	Xtrriage
Anisotropy	0.202	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.22$	Xtrriage
Estimated twinning fraction	0.069 for -k,-h,-l 0.075 for k,h,-l 0.248 for h,-k,-l	Xtrriage
Reported twinning fraction	0.350 for -h,-k,l	Depositor
Outliers	0 of 91940 reflections	Xtrriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	46492	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.20	0/4524	0.37	0/6132
1	B	0.22	0/4519	0.39	0/6125
1	C	0.21	0/4346	0.38	0/5903
1	I	0.21	0/4361	0.40	1/5929 (0.0%)
1	J	0.25	0/4498	0.41	0/6096
1	K	0.21	0/4511	0.38	0/6115
2	D	0.24	0/3443	0.43	1/4660 (0.0%)
2	E	0.21	0/3461	0.38	0/4693
2	F	0.21	0/3351	0.39	0/4548
2	L	0.21	0/3488	0.38	0/4717
2	M	0.20	0/3413	0.38	0/4626
2	N	0.22	0/3218	0.40	0/4367
All	All	0.22	0/47133	0.39	2/63911 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	339	GLN	C-N-CD	-8.10	102.78	120.60
1	I	501	LEU	CA-CB-CG	6.77	130.86	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4448	0	4297	295	0
1	B	4443	0	4280	328	0
1	C	4274	0	3998	290	0
1	I	4291	0	4030	278	0
1	J	4422	0	4271	349	0
1	K	4435	0	4281	344	0
2	D	3383	0	3279	298	0
2	E	3400	0	3255	261	0
2	F	3295	0	3157	263	0
2	L	3428	0	3377	312	0
2	M	3357	0	3207	281	0
2	N	3162	0	2993	307	0
3	A	6	8	8	0	0
4	B	31	0	13	9	0
4	C	31	0	13	14	0
4	J	31	0	13	7	0
4	K	31	0	13	16	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	J	1	0	0	0	0
5	K	1	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	E	1	0	0	0	0
6	I	2	0	0	3	0
6	L	1	0	0	0	0
6	M	1	0	0	1	0
6	N	4	0	0	0	0
All	All	46484	8	44485	3401	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

The worst 5 of 3401 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:20:SER:HB3	1:I:45:ASP:HB2	1.25	1.16
1:K:497:ARG:HA	1:K:501:LEU:HG	1.28	1.16

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:9:VAL:HG21	1:J:58:ILE:HB	1.27	1.16
1:J:215:VAL:HB	1:J:501:LEU:HA	1.23	1.15
1:B:497:ARG:HA	1:B:501:LEU:HB2	1.29	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	582/600 (97%)	579 (100%)	3 (0%)	0	100	100
1	B	582/600 (97%)	570 (98%)	12 (2%)	0	100	100
1	C	582/600 (97%)	569 (98%)	13 (2%)	0	100	100
1	I	579/600 (96%)	570 (98%)	9 (2%)	0	100	100
1	J	582/600 (97%)	572 (98%)	10 (2%)	0	100	100
1	K	582/600 (97%)	572 (98%)	10 (2%)	0	100	100
2	D	443/465 (95%)	441 (100%)	1 (0%)	1 (0%)	47	78
2	E	451/465 (97%)	447 (99%)	4 (1%)	0	100	100
2	F	443/465 (95%)	438 (99%)	5 (1%)	0	100	100
2	L	443/465 (95%)	435 (98%)	8 (2%)	0	100	100
2	M	451/465 (97%)	441 (98%)	10 (2%)	0	100	100
2	N	427/465 (92%)	411 (96%)	16 (4%)	0	100	100
All	All	6147/6390 (96%)	6045 (98%)	101 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	105	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	471/511 (92%)	430 (91%)	41 (9%)	10	35
1	B	465/511 (91%)	418 (90%)	47 (10%)	7	28
1	C	420/511 (82%)	375 (89%)	45 (11%)	6	25
1	I	430/511 (84%)	391 (91%)	39 (9%)	9	32
1	J	461/511 (90%)	416 (90%)	45 (10%)	8	29
1	K	467/511 (91%)	423 (91%)	44 (9%)	8	31
2	D	337/387 (87%)	301 (89%)	36 (11%)	6	25
2	E	329/387 (85%)	294 (89%)	35 (11%)	6	25
2	F	317/387 (82%)	275 (87%)	42 (13%)	4	16
2	L	354/387 (92%)	309 (87%)	45 (13%)	4	18
2	M	323/387 (84%)	282 (87%)	41 (13%)	4	18
2	N	300/387 (78%)	251 (84%)	49 (16%)	2	10
All	All	4674/5388 (87%)	4165 (89%)	509 (11%)	6	24

5 of 509 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	308	GLU
2	M	215	ASN
1	J	13	LEU
2	M	146	LYS
2	N	98	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	N	365	HIS
2	N	129	GLN
1	J	213	GLN
2	L	163	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	J	50	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ANP	K	601	5	29,33,33	1.09	4 (13%)	31,52,52	1.09	2 (6%)
3	GOL	A	601	-	5,5,5	0.37	0	5,5,5	0.15	0
4	ANP	B	601	5	29,33,33	1.10	4 (13%)	31,52,52	1.24	3 (9%)
4	ANP	J	601	5	29,33,33	1.11	4 (13%)	31,52,52	1.20	4 (12%)
4	ANP	C	601	5	29,33,33	1.07	4 (13%)	31,52,52	1.33	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	K	601	5	-	3/14/38/38	0/3/3/3
3	GOL	A	601	-	-	2/4/4/4	-
4	ANP	B	601	5	-	3/14/38/38	0/3/3/3
4	ANP	J	601	5	-	5/14/38/38	0/3/3/3
4	ANP	C	601	5	-	1/14/38/38	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	J	601	ANP	PG-N3B	2.75	1.70	1.63
4	C	601	ANP	PG-N3B	2.52	1.69	1.63
4	C	601	ANP	PG-O1G	2.51	1.50	1.46
4	B	601	ANP	PB-O3A	-2.48	1.55	1.59
4	K	601	ANP	PG-O1G	2.47	1.50	1.46

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	K	601	ANP	PB-O3A-PA	-3.59	119.98	132.62
4	B	601	ANP	PB-O3A-PA	-3.58	119.99	132.62
4	C	601	ANP	PB-O3A-PA	-3.57	120.03	132.62
4	J	601	ANP	PB-O3A-PA	-3.29	121.04	132.62
4	C	601	ANP	C3'-C2'-C1'	3.12	105.68	100.98

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	GOL	C1-C2-C3-O3
3	A	601	GOL	O2-C2-C3-O3
4	B	601	ANP	PB-N3B-PG-O1G
4	B	601	ANP	PG-N3B-PB-O1B
4	B	601	ANP	PG-N3B-PB-O3A

There are no ring outliers.

4 monomers are involved in 46 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	K	601	ANP	16	0
4	B	601	ANP	9	0
4	J	601	ANP	7	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	601	ANP	14	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	584/600 (97%)	0.52	29 (4%) 28 32	42, 77, 103, 125	0
1	B	584/600 (97%)	0.53	40 (6%) 17 21	37, 72, 98, 116	0
1	C	584/600 (97%)	0.61	49 (8%) 11 13	39, 80, 129, 154	0
1	I	583/600 (97%)	0.42	31 (5%) 26 30	38, 72, 100, 125	0
1	J	584/600 (97%)	0.68	47 (8%) 12 14	66, 92, 117, 151	0
1	K	584/600 (97%)	0.50	44 (7%) 14 17	34, 69, 105, 178	0
2	D	447/465 (96%)	0.64	36 (8%) 12 14	45, 78, 112, 131	0
2	E	453/465 (97%)	0.49	18 (3%) 38 41	49, 73, 107, 135	0
2	F	445/465 (95%)	0.81	47 (10%) 6 8	44, 75, 121, 154	0
2	L	447/465 (96%)	0.62	35 (7%) 13 16	37, 70, 122, 153	0
2	M	453/465 (97%)	0.78	47 (10%) 6 8	52, 90, 130, 150	0
2	N	429/465 (92%)	0.85	56 (13%) 3 4	51, 93, 128, 168	0
All	All	6177/6390 (96%)	0.61	479 (7%) 13 16	34, 78, 117, 178	0

The worst 5 of 479 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	M	389	LEU	8.5
2	F	386	ALA	7.3
2	F	445	ILE	6.7
2	D	102	GLU	6.7
1	K	567	LEU	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	C	602	1/1	0.75	0.26	49,49,49,49	0
3	GOL	A	601	6/6	0.82	0.19	11,13,16,16	0
4	ANP	J	601	31/31	0.94	0.15	29,47,58,71	0
4	ANP	K	601	31/31	0.94	0.18	29,45,69,85	0
4	ANP	C	601	31/31	0.94	0.22	28,34,99,113	0
4	ANP	B	601	31/31	0.95	0.16	19,28,45,45	0
5	MG	J	602	1/1	0.96	0.09	26,26,26,26	0
5	MG	B	602	1/1	0.97	0.14	8,8,8,8	0
5	MG	K	602	1/1	0.97	0.13	27,27,27,27	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP J 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ANP K 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ANP C 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.