



wwPDB EM Validation Summary Report ⓘ

Feb 10, 2026 – 05:38 PM EST

PDB ID : 9E2D / pdb_00009e2d
EMDB ID : EMD-47448
Title : Cryo-EM structure of human DNMT3A2-DNMT3B3 complex bound to a di-nucleosome with a five base pair linker
Authors : Xie, X.; Liu, M.; Zhou, X.E.; Worden, E.; Jones, P.
Deposited on : 2024-10-22
Resolution : 6.08 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev131
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 6.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	489 (5.58 - 6.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	<p>61% 10% 29%</p>
1	E	135	<p>61% 10% 28%</p>
1	M	135	<p>66% 7% 27%</p>
1	Q	135	<p>63% 7% 30%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	103	68% 9% 23%
2	F	103	69% 7% 24%
2	N	103	69% 7% 24%
2	R	103	70% 6% 24%
3	C	129	68% 9% 23%
3	G	129	74% 0% 24%
3	O	129	71% 5% 24%
3	S	129	72% 0% 24%
4	D	123	64% 11% 25%
4	H	123	65% 10% 25%
4	P	123	70% 0% 26%
4	T	123	72% 0% 25%
5	L	773	15% 37% 7% 57%
5	V	773	40% 0% 55%
5	X	773	20% 0% 77%
5	Z	773	20% 0% 77%
6	K	689	33% 72% 9% 20%
6	U	689	54% 8% 38%
6	W	689	18% 51% 10% 39%
6	Y	689	10% 72% 9% 19%
7	I	317	91% 9% 0%
8	J	316	94% 6% 0%

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 48657 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	96	Total 795	C 501	N 154	O 137	S 3	0	0
1	E	97	Total 802	C 506	N 155	O 138	S 3	0	0
1	M	99	Total 820	C 518	N 159	O 140	S 3	0	0
1	Q	95	Total 785	C 495	N 151	O 136	S 3	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	GLY	conflict	UNP Q71DI3
E	102	ALA	GLY	conflict	UNP Q71DI3
M	102	ALA	GLY	conflict	UNP Q71DI3
Q	102	ALA	GLY	conflict	UNP Q71DI3

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	79	Total 633	C 399	N 124	O 109	S 1	0	0
2	F	78	Total 622	C 393	N 120	O 108	S 1	0	0
2	N	78	Total 622	C 393	N 120	O 108	S 1	0	0
2	R	78	Total 622	C 393	N 120	O 108	S 1	0	0

- Molecule 3 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	99	Total	C	N	O	0	0
			764	479	151	134		
3	G	98	Total	C	N	O	0	0
			755	474	149	132		
3	O	98	Total	C	N	O	0	0
			755	474	149	132		
3	S	98	Total	C	N	O	0	0
			752	473	148	131		

- Molecule 4 is a protein called Histone H2B.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	92	Total	C	N	O	S	0	0
			719	453	129	135	2		
4	H	92	Total	C	N	O	S	0	0
			719	453	129	135	2		
4	P	91	Total	C	N	O	S	0	0
			708	447	125	134	2		
4	T	92	Total	C	N	O	S	0	0
			719	453	129	135	2		

- Molecule 5 is a protein called Isoform 3 of DNA (cytosine-5)-methyltransferase 3B.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	V	344	Total	C	N	O	S	0	0
			2753	1751	481	495	26		
5	Z	175	Total	C	N	O	S	0	0
			1421	930	240	245	6		
5	L	335	Total	C	N	O	S	0	0
			2677	1700	464	488	25		
5	X	176	Total	C	N	O	S	0	0
			1434	938	246	245	5		

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
V	391	LYS	-	insertion	UNP Q9UBC3
V	392	ASP	-	insertion	UNP Q9UBC3
V	393	TYR	-	insertion	UNP Q9UBC3
Z	391	LYS	-	insertion	UNP Q9UBC3
Z	392	ASP	-	insertion	UNP Q9UBC3
Z	393	TYR	-	insertion	UNP Q9UBC3
L	391	LYS	-	insertion	UNP Q9UBC3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	392	ASP	-	insertion	UNP Q9UBC3
L	393	TYR	-	insertion	UNP Q9UBC3
X	391	LYS	-	insertion	UNP Q9UBC3
X	392	ASP	-	insertion	UNP Q9UBC3
X	393	TYR	-	insertion	UNP Q9UBC3

- Molecule 6 is a protein called DNA (cytosine-5)-methyltransferase 3A.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	U	424	Total	C	N	O	S	0	0
			3404	2156	604	609	35		
6	Y	555	Total	C	N	O	S	0	0
			4451	2839	784	786	42		
6	W	423	Total	C	N	O	S	0	0
			3400	2154	603	608	35		
6	K	552	Total	C	N	O	S	0	0
			4426	2824	778	782	42		

- Molecule 7 is a DNA chain called DNA (317-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	I	317	Total	C	N	O	P	0	0
			6532	3088	1229	1898	317		

- Molecule 8 is a DNA chain called DNA (316-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
8	J	316	Total	C	N	O	P	0	0
			6445	3057	1173	1899	316		

- Molecule 9 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
9	V	3	Total	Zn	0
			3	3	
9	U	3	Total	Zn	0
			3	3	
9	Y	3	Total	Zn	0
			3	3	
9	L	3	Total	Zn	0
			3	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
9	W	3	Total	Zn	0
			3	3	
9	K	3	Total	Zn	0
			3	3	

- Molecule 10 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).




Mol	Chain	Residues	Atoms					AltConf
10	U	1	Total	C	N	O	S	0
			26	14	6	5	1	
10	Y	1	Total	C	N	O	S	0
			26	14	6	5	1	
10	W	1	Total	C	N	O	S	0
			26	14	6	5	1	
10	K	1	Total	C	N	O	S	0
			26	14	6	5	1	


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H3.2




- Molecule 1: Histone H3.2



- Molecule 1: Histone H3.2



- Molecule 1: Histone H3.2



• Molecule 2: Histone H4



• Molecule 2: Histone H4



• Molecule 2: Histone H4



• Molecule 2: Histone H4



• Molecule 3: Histone H2A




• Molecule 3: Histone H2A




• Molecule 3: Histone H2A




• Molecule 3: Histone H2A




● Molecule 5: Isoform 3 of DNA (cytosine-5)-methyltransferase 3B




● Molecule 5: Isoform 3 of DNA (cytosine-5)-methyltransferase 3B







● Molecule 6: DNA (cytosine-5)-methyltransferase 3A



● Molecule 6: DNA (cytosine-5)-methyltransferase 3A



• Molecule 7: DNA (317-MER)



• Molecule 8: DNA (316-MER)



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	41237	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TALOS ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1100	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.354	Depositor
Minimum map value	-0.085	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.018	Depositor
Recommended contour level	0.051	Depositor
Map size (Å)	371.19998, 371.19998, 371.19998	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.16, 1.16, 1.16	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.98	0/806	1.42	0/1081
1	E	0.98	0/814	1.41	0/1092
1	M	0.97	0/832	1.40	0/1115
1	Q	0.97	0/795	1.43	0/1066
2	B	0.98	0/640	1.39	0/857
2	F	0.99	0/629	1.44	0/843
2	N	0.98	0/629	1.43	0/843
2	R	0.98	0/629	1.44	0/843
3	C	0.98	0/773	1.43	0/1043
3	G	0.99	0/764	1.43	0/1031
3	O	0.99	0/764	1.42	0/1031
3	S	0.98	0/761	1.43	0/1027
4	D	1.00	0/730	1.52	0/983
4	H	1.00	0/730	1.53	0/983
4	P	1.02	0/719	1.52	0/969
4	T	1.01	0/730	1.51	0/983
5	L	0.99	0/2740	1.36	0/3700
5	V	0.99	0/2821	1.36	2/3810 (0.1%)
5	X	0.98	0/1472	1.32	0/1985
5	Z	0.98	0/1459	1.31	0/1970
6	K	0.99	0/4545	1.33	0/6142
6	U	1.00	0/3487	1.34	0/4712
6	W	1.00	0/3482	1.36	0/4703
6	Y	1.00	0/4570	1.31	0/6176
7	I	0.28	0/7336	0.57	2/11329 (0.0%)
8	J	0.28	0/7222	0.56	1/11136 (0.0%)
All	All	0.85	0/50879	1.18	5/71453 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	V	752	SER	CA-C-N	5.79	124.97	120.33
5	V	752	SER	C-N-CA	5.79	124.97	120.33
8	J	8	DG	C4'-C3'-O3'	5.73	118.60	110.00
7	I	3	DG	C4'-C3'-O3'	5.32	117.97	110.00
7	I	313	DG	C4'-C3'-O3'	5.04	117.56	110.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	795	0	833	14	0
1	E	802	0	841	12	0
1	M	820	0	866	11	0
1	Q	785	0	826	14	0
2	B	633	0	673	7	0
2	F	622	0	660	10	0
2	N	622	0	660	7	0
2	R	622	0	660	5	0
3	C	764	0	808	15	0
3	G	755	0	800	9	0
3	O	755	0	800	14	0
3	S	752	0	796	5	0
4	D	719	0	740	17	0
4	H	719	0	740	22	0
4	P	708	0	727	4	0
4	T	719	0	740	3	0
5	L	2677	0	2600	41	0
5	V	2753	0	2673	21	0
5	X	1434	0	1436	16	0
5	Z	1421	0	1422	10	0
6	K	4426	0	4307	40	0
6	U	3404	0	3317	41	0
6	W	3400	0	3314	54	0
6	Y	4451	0	4336	48	0
7	I	6532	0	3552	41	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	J	6445	0	3544	30	0
9	K	3	0	0	0	0
9	L	3	0	0	0	0
9	U	3	0	0	0	0
9	V	3	0	0	0	0
9	W	3	0	0	0	0
9	Y	3	0	0	0	0
10	K	26	0	19	3	0
10	U	26	0	19	4	0
10	W	26	0	19	0	0
10	Y	26	0	19	0	0
All	All	48657	0	42747	413	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 413 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:70:ILE:HA	4:H:98:LEU:CD1	1.76	1.15
4:H:95:VAL:HG13	4:H:99:LEU:HD12	1.16	1.14
3:G:31:HIS:CE1	3:G:35:ARG:HH21	1.67	1.12
3:C:42:ARG:HA	8:J:275:DA:OP1	1.70	0.91
4:H:95:VAL:CG1	4:H:99:LEU:HD12	2.03	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	94/135 (70%)	91 (97%)	3 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	95/135 (70%)	94 (99%)	1 (1%)	0	100	100
1	M	97/135 (72%)	95 (98%)	2 (2%)	0	100	100
1	Q	93/135 (69%)	92 (99%)	1 (1%)	0	100	100
2	B	77/103 (75%)	76 (99%)	1 (1%)	0	100	100
2	F	76/103 (74%)	74 (97%)	2 (3%)	0	100	100
2	N	76/103 (74%)	74 (97%)	2 (3%)	0	100	100
2	R	76/103 (74%)	74 (97%)	2 (3%)	0	100	100
3	C	97/129 (75%)	95 (98%)	2 (2%)	0	100	100
3	G	96/129 (74%)	93 (97%)	3 (3%)	0	100	100
3	O	96/129 (74%)	92 (96%)	4 (4%)	0	100	100
3	S	96/129 (74%)	95 (99%)	1 (1%)	0	100	100
4	D	90/123 (73%)	86 (96%)	4 (4%)	0	100	100
4	H	90/123 (73%)	85 (94%)	5 (6%)	0	100	100
4	P	89/123 (72%)	85 (96%)	4 (4%)	0	100	100
4	T	90/123 (73%)	89 (99%)	1 (1%)	0	100	100
5	L	329/773 (43%)	314 (95%)	15 (5%)	0	100	100
5	V	338/773 (44%)	326 (96%)	12 (4%)	0	100	100
5	X	168/773 (22%)	159 (95%)	9 (5%)	0	100	100
5	Z	167/773 (22%)	161 (96%)	6 (4%)	0	100	100
6	K	544/689 (79%)	521 (96%)	23 (4%)	0	100	100
6	U	420/689 (61%)	400 (95%)	20 (5%)	0	100	100
6	W	417/689 (60%)	399 (96%)	18 (4%)	0	100	100
6	Y	547/689 (79%)	525 (96%)	22 (4%)	0	100	100
All	All	4358/7808 (56%)	4195 (96%)	163 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/110 (76%)	84 (100%)	0	100	100
1	E	85/110 (77%)	85 (100%)	0	100	100
1	M	87/110 (79%)	87 (100%)	0	100	100
1	Q	83/110 (76%)	83 (100%)	0	100	100
2	B	65/79 (82%)	65 (100%)	0	100	100
2	F	64/79 (81%)	64 (100%)	0	100	100
2	N	64/79 (81%)	64 (100%)	0	100	100
2	R	64/79 (81%)	64 (100%)	0	100	100
3	C	77/101 (76%)	77 (100%)	0	100	100
3	G	76/101 (75%)	76 (100%)	0	100	100
3	O	76/101 (75%)	76 (100%)	0	100	100
3	S	75/101 (74%)	75 (100%)	0	100	100
4	D	78/103 (76%)	78 (100%)	0	100	100
4	H	78/103 (76%)	78 (100%)	0	100	100
4	P	77/103 (75%)	77 (100%)	0	100	100
4	T	78/103 (76%)	78 (100%)	0	100	100
5	L	291/660 (44%)	291 (100%)	0	100	100
5	V	299/660 (45%)	299 (100%)	0	100	100
5	X	151/660 (23%)	150 (99%)	1 (1%)	81	87
5	Z	152/660 (23%)	152 (100%)	0	100	100
6	K	475/591 (80%)	475 (100%)	0	100	100
6	U	372/591 (63%)	372 (100%)	0	100	100
6	W	372/591 (63%)	372 (100%)	0	100	100
6	Y	478/591 (81%)	478 (100%)	0	100	100
All	All	3801/6576 (58%)	3800 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	X	612	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
6	Y	82	GLN
5	L	678	ASN
6	Y	129	ASN
6	Y	816	GLN
5	X	601	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 18 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	SAH	U	1004	-	23,28,28	0.73	0	22,40,40	0.65	1 (4%)
10	SAH	Y	1004	-	23,28,28	0.74	0	22,40,40	0.65	1 (4%)
10	SAH	K	1004	-	23,28,28	0.73	0	22,40,40	0.65	1 (4%)
10	SAH	W	1004	-	23,28,28	0.73	0	22,40,40	0.65	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	SAH	U	1004	-	-	4/11/31/31	0/3/3/3
10	SAH	Y	1004	-	-	4/11/31/31	0/3/3/3
10	SAH	K	1004	-	-	3/11/31/31	0/3/3/3
10	SAH	W	1004	-	-	3/11/31/31	0/3/3/3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	Y	1004	SAH	C5-C6-N6	2.31	123.83	120.31
10	U	1004	SAH	C5-C6-N6	2.26	123.75	120.31
10	K	1004	SAH	C5-C6-N6	2.25	123.74	120.31
10	W	1004	SAH	C5-C6-N6	2.24	123.73	120.31

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	U	1004	SAH	N-CA-CB-CG
10	Y	1004	SAH	N-CA-CB-CG
10	W	1004	SAH	N-CA-CB-CG
10	K	1004	SAH	N-CA-CB-CG
10	U	1004	SAH	C-CA-CB-CG


There are no ring outliers.


2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	U	1004	SAH	4	0
10	K	1004	SAH	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-47448. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 167




Y Index: 158




Z Index: 199


6.3.2 Raw map



X Index: 166



Y Index: 159




Z Index: 193


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.051. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1367 nm³; this corresponds to an approximate mass of 1235 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.164 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.164 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	6.08	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	7.48	9.77	7.72

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.48 differs from the reported value 6.08 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-47448 and PDB model 9E2D. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.051 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.051).

9.4 Atom inclusion [i](#)



At the recommended contour level, 87% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.051) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8680	 0.0830
A	 0.9240	 0.1430
B	 0.9190	 0.1380
C	 0.9090	 0.1530
D	 0.9360	 0.1450
E	 0.9120	 0.1350
F	 0.9050	 0.1340
G	 0.9150	 0.1530
H	 0.9290	 0.1440
I	 0.9580	 0.1470
J	 0.9600	 0.1430
K	 0.5700	 0.0200
L	 0.6330	 0.0220
M	 0.9300	 0.0560
N	 0.9380	 0.0360
O	 0.9330	 0.0720
P	 0.9810	 0.0460
Q	 0.9720	 0.0810
R	 0.9830	 0.0940
S	 0.9860	 0.0790
T	 0.9770	 0.0710
U	 0.9420	 0.0570
V	 0.9130	 0.0570
W	 0.6920	 0.0200
X	 0.8590	 0.0390
Y	 0.8590	 0.0590
Z	 0.9830	 0.0510

