



# Full wwPDB X-ray Structure Validation Report i

Dec 23, 2024 – 12:09 PM EST

PDB ID : 9E8M

Title : Covalent inhibitor VVD-442 bound to the RAS binding domain (RBD) of PI3Ka

Authors : Bernard, S.M.; Tamiya, J.

Deposited on : 2024-11-05

Resolution : 2.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.21

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.004 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4293 (2.84-2.80)
Clashscore	180529	4801 (2.84-2.80)
Ramachandran outliers	177936	4739 (2.84-2.80)
Sidechain outliers	177891	4741 (2.84-2.80)
RSRZ outliers	164620	4295 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)


There are 3 unique types of molecules in this entry. The entry contains 2383 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	142	Total	C 1156	N 746	O 192	S 207	0	0	0
1	B	133	Total	C 1089	N 707	O 179	S 192	0	0	0

- Molecule 2 is 1-[(1P)-5-bromo-2'-chloro[1,1'-biphenyl]-2-sulfonyl]-4-fluoro-N-[(2S)-4-(methanesulfonyl)butan-2-yl]piperidine-4-carboxamide (three-letter code: A1BF9) (formula: C<sub>23</sub>H<sub>27</sub>BrClF<sub>2</sub>N<sub>2</sub>O<sub>5</sub>S<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms								ZeroOcc	AltConf
2	A	1	Total	Br 35	C 1	Cl 23	F 1	N 1	O 2	S 5	0	0
2	B	1	Total	Br 35	C 1	Cl 23	F 1	N 1	O 2	S 5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	40	Total O 40 40	0	0
3	B	28	Total O 28 28	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform



- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.39Å 88.30Å 139.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.89 – 2.83 47.89 – 2.83	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.89-2.83) 99.9 (47.89-2.83)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.83 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
$R$ , $R_{free}$	0.191 , 0.268 0.202 , 0.268	Depositor DCC
$R_{free}$ test set	485 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.2	Xtriage
Anisotropy	0.422	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 39.6	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2383	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1BF9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1182	0.88	0/1597
1	B	0.37	0/1112	0.80	0/1498
All	All	0.42	0/2294	0.84	0/3095

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	226	ARG	Sidechain

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1156	0	1193	36	0
1	B	1089	0	1131	15	0
2	A	35	0	0	0	0
2	B	35	0	0	2	0
3	A	40	0	0	1	0
3	B	28	0	0	0	0
All	All	2383	0	2324	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:VAL:HG23	1:A:199:SER:HB3	1.65	0.76
1:A:199:SER:HB2	1:A:200:PRO:HD3	1.71	0.73
1:A:199:SER:CB	1:A:200:PRO:CD	2.72	0.67
1:A:199:SER:HB2	1:A:200:PRO:CD	2.25	0.66
1:A:253:LYS:HD2	1:A:288:MET:HE1	1.76	0.66
1:B:253:LYS:HE2	1:B:257:CYS:O	1.96	0.65
1:A:196:VAL:HG12	1:A:198:VAL:HG12	1.80	0.63
1:A:171:VAL:HG11	1:A:265:TYR:HD2	1.68	0.58
1:A:253:LYS:NZ	1:A:257:CYS:O	2.28	0.57
1:B:250:TYR:CE2	2:B:401:A1BF9:CL	2.96	0.56
1:A:260:TYR:HB3	1:A:262:LEU:HD13	1.88	0.55
1:A:225:ILE:O	1:A:229:THR:HB	2.08	0.54
1:B:228:LYS:HD2	2:B:401:A1BF9:C19	2.39	0.53
1:B:186:ASP:O	1:B:189:GLN:HB2	2.09	0.52
1:B:164:MET:HE3	1:B:169:PRO:CD	2.40	0.52
1:B:160:HIS:CE1	1:B:164:MET:HG3	2.44	0.52
1:A:171:VAL:HG11	1:A:265:TYR:CD2	2.46	0.51
1:A:198:VAL:HG23	1:A:199:SER:N	2.26	0.50
1:A:294:TYR:HA	1:A:297:LEU:HD22	1.93	0.50
1:A:160:HIS:NE2	1:A:164:MET:CE	2.74	0.50
1:A:160:HIS:NE2	1:A:164:MET:HE2	2.27	0.50
1:B:221:ILE:HG12	1:B:287:LEU:HD21	1.95	0.49
1:A:252:LEU:HD13	1:A:285:LEU:HD23	1.97	0.47
1:A:216:VAL:HG13	1:A:217:PRO:HD2	1.97	0.46
1:B:289:ALA:O	1:B:292:SER:HB3	2.16	0.46
1:A:196:VAL:CG1	1:A:198:VAL:HG12	2.46	0.46
1:A:255:CYS:HB2	1:A:284:ASN:O	2.16	0.45
1:A:198:VAL:HG23	1:A:199:SER:CB	2.40	0.45

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:229:THR:O	1:B:230:ARG:C	2.55	0.44
1:A:211:ILE:CD1	1:A:220:VAL:HG22	2.48	0.44
1:B:250:TYR:HB3	1:B:287:LEU:HD22	2.00	0.44
1:A:229:THR:O	1:A:229:THR:CG2	2.66	0.43
1:B:263:GLU:HA	1:B:263:GLU:OE1	2.17	0.43
1:A:268:SER:HA	1:A:273:ILE:CG2	2.49	0.43
1:A:178:PRO:HD2	1:A:181:ILE:HD12	2.00	0.42
1:B:243:VAL:O	1:B:247:GLN:HB2	2.20	0.42
1:A:216:VAL:HB	1:A:219:GLN:OE1	2.19	0.42
1:A:263:GLU:O	1:A:265:TYR:N	2.52	0.42
1:A:207:TYR:CZ	1:A:228:LYS:HG3	2.55	0.42
1:A:267:LEU:HG	1:A:273:ILE:CG1	2.50	0.41
1:A:221:ILE:O	1:A:224:ALA:HB3	2.20	0.41
1:B:223:GLU:HG2	1:B:226:ARG:HH21	1.84	0.41
1:A:297:LEU:HD12	1:A:297:LEU:HA	1.88	0.41
1:B:207:TYR:OH	1:B:228:LYS:HE2	2.20	0.41
1:A:167:TYR:CE2	1:A:293:LEU:HD11	2.55	0.41
1:A:160:HIS:CD2	1:A:164:MET:HE2	2.54	0.41
1:A:166:VAL:HG12	1:A:167:TYR:CD1	2.56	0.41
1:A:267:LEU:HG	1:A:273:ILE:HG12	2.01	0.41
1:A:211:ILE:CG2	1:A:223:GLU:HG3	2.51	0.40
1:A:222:ALA:HB3	3:A:516:HOH:O	2.22	0.40
1:B:287:LEU:HD23	1:B:287:LEU:HA	1.88	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	140/144 (97%)	132 (94%)	5 (4%)	3 (2%)	5 19
1	B	129/144 (90%)	123 (95%)	5 (4%)	1 (1%)	16 42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	269/288 (93%)	255 (95%)	10 (4%)	4 (2%)	8 26

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	SER
1	A	247	GLN
1	A	264	LYS
1	B	230	ARG

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	134/136 (98%)	121 (90%)	13 (10%)	6 20
1	B	125/136 (92%)	115 (92%)	10 (8%)	10 28
All	All	259/272 (95%)	236 (91%)	23 (9%)	8 24

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	SER
1	A	196	VAL
1	A	227	LYS
1	A	229	THR
1	A	237	GLU
1	A	255	CYS
1	A	258	ASP
1	A	259	GLU
1	A	262	LEU
1	A	273	ILE
1	A	275	SER
1	A	286	MET
1	A	287	LEU
1	B	169	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	231	SER
1	B	233	LEU
1	B	239	LEU
1	B	244	LEU
1	B	245	GLU
1	B	247	GLN
1	B	255	CYS
1	B	262	LEU
1	B	275	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	ASN
1	A	205	GLN
1	A	296	GLN
1	B	160	HIS
1	B	183	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1BF9	B	401	1	34,37,37	2.84	9 (26%)	49,56,56	3.18	21 (42%)
2	A1BF9	A	401	1	34,37,37	2.81	9 (26%)	49,56,56	2.96	19 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1BF9	B	401	1	-	12/30/44/44	0/3/3/3
2	A1BF9	A	401	1	-	10/30/44/44	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	A1BF9	C11-S1	-10.84	1.64	1.78
2	B	401	A1BF9	C11-S1	-10.05	1.65	1.78
2	B	401	A1BF9	C1-S	-8.94	1.67	1.78
2	A	401	A1BF9	C1-S	-8.89	1.67	1.78
2	B	401	A1BF9	F-C6	-5.52	1.33	1.42
2	A	401	A1BF9	F-C6	-5.04	1.33	1.42
2	B	401	A1BF9	BR-C14	3.05	1.96	1.90
2	B	401	A1BF9	C8-N1	2.92	1.50	1.47
2	B	401	A1BF9	C9-N1	2.80	1.50	1.47
2	B	401	A1BF9	S1-N1	2.49	1.67	1.63
2	B	401	A1BF9	C22-CL	2.42	1.79	1.73
2	A	401	A1BF9	C22-CL	2.35	1.79	1.73
2	B	401	A1BF9	C-S	-2.30	1.68	1.75
2	A	401	A1BF9	BR-C14	2.25	1.94	1.90
2	A	401	A1BF9	S1-N1	2.22	1.66	1.63
2	A	401	A1BF9	C16-C11	2.18	1.44	1.40
2	A	401	A1BF9	C8-N1	2.08	1.50	1.47
2	A	401	A1BF9	C-S	-2.02	1.68	1.75

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	A1BF9	O4-S1-O3	-8.90	105.69	119.59
2	A	401	A1BF9	O1-S-C1	-8.73	102.03	108.33
2	B	401	A1BF9	O4-S1-N1	8.60	114.79	106.69
2	B	401	A1BF9	O-S-C1	-8.10	102.49	108.33
2	A	401	A1BF9	O-S-C1	7.63	113.83	108.33
2	A	401	A1BF9	O1-S-C	6.95	115.33	108.87
2	B	401	A1BF9	O1-S-C1	-6.22	103.84	108.33
2	A	401	A1BF9	O4-S1-N1	5.78	112.13	106.69
2	A	401	A1BF9	O1-S-O	-5.44	107.02	117.22
2	B	401	A1BF9	C10-C9-N1	5.38	116.14	109.16
2	A	401	A1BF9	O4-S1-O3	-4.99	111.81	119.59
2	B	401	A1BF9	C15-C16-C11	4.75	121.38	117.44
2	B	401	A1BF9	C13-C14-C15	-4.52	115.02	121.53
2	B	401	A1BF9	C7-C8-N1	4.49	114.98	109.16
2	A	401	A1BF9	C7-C8-N1	4.37	114.83	109.16
2	B	401	A1BF9	C-S-C1	4.04	120.29	105.28
2	B	401	A1BF9	C8-C7-C6	-3.92	108.54	112.39
2	A	401	A1BF9	C9-N1-C8	3.87	116.56	112.12
2	A	401	A1BF9	C10-C9-N1	3.78	114.06	109.16
2	A	401	A1BF9	C9-C10-C6	3.78	116.11	112.39
2	A	401	A1BF9	C8-C7-C6	-3.69	108.76	112.39
2	B	401	A1BF9	C12-C11-C16	-3.56	116.72	121.52
2	B	401	A1BF9	O3-S1-C11	3.45	114.04	107.35
2	B	401	A1BF9	C12-C13-C14	3.28	123.11	119.18
2	A	401	A1BF9	O3-S1-N1	3.18	109.68	106.69
2	A	401	A1BF9	C16-C17-C22	3.06	126.31	122.92
2	B	401	A1BF9	C9-N1-C8	3.06	115.63	112.12
2	A	401	A1BF9	C13-C14-C15	-2.82	117.47	121.53
2	B	401	A1BF9	O3-S1-N1	2.75	109.27	106.69
2	B	401	A1BF9	BR-C14-C13	2.63	122.88	119.30
2	A	401	A1BF9	C15-C16-C11	2.58	119.58	117.44
2	A	401	A1BF9	C12-C13-C14	2.56	122.25	119.18
2	B	401	A1BF9	O1-S-O	-2.54	112.45	117.22
2	B	401	A1BF9	C16-C15-C14	2.43	122.64	118.83
2	B	401	A1BF9	C2-C3-N	-2.25	106.55	110.42
2	B	401	A1BF9	C18-C17-C22	2.13	120.00	117.60
2	A	401	A1BF9	C21-C22-CL	-2.09	114.31	118.42
2	A	401	A1BF9	C7-C6-C5	2.04	113.48	109.14
2	B	401	A1BF9	BR-C14-C15	2.03	122.10	119.23
2	A	401	A1BF9	C17-C22-CL	2.02	123.43	120.67

There are no chirality outliers.

All (22) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	A1BF9	C2-C1-S-O
2	A	401	A1BF9	C2-C1-S-O1
2	B	401	A1BF9	C1-C2-C3-C4
2	B	401	A1BF9	C1-C2-C3-N
2	B	401	A1BF9	C2-C1-S-O
2	B	401	A1BF9	C2-C1-S-O1
2	B	401	A1BF9	C2-C1-S-C
2	B	401	A1BF9	C9-N1-S1-O4
2	A	401	A1BF9	C9-N1-S1-O4
2	B	401	A1BF9	C8-N1-S1-O3
2	A	401	A1BF9	C9-N1-S1-C11
2	A	401	A1BF9	C9-N1-S1-O3
2	B	401	A1BF9	C9-N1-S1-C11
2	B	401	A1BF9	C9-N1-S1-O3
2	B	401	A1BF9	C8-N1-S1-O4
2	B	401	A1BF9	C8-N1-S1-C11
2	A	401	A1BF9	C8-N1-S1-O3
2	A	401	A1BF9	C2-C1-S-C
2	A	401	A1BF9	C8-N1-S1-O4
2	A	401	A1BF9	C8-N1-S1-C11
2	A	401	A1BF9	S-C1-C2-C3
2	B	401	A1BF9	C4-C3-N-C5

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	A1BF9	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	142/144 (98%)	-0.44	1 (0%)	84	79	20, 33, 57, 85
1	B	133/144 (92%)	-0.18	2 (1%)	71	65	25, 44, 71, 99
All	All	275/288 (95%)	-0.31	3 (1%)	77	72	20, 40, 67, 99

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	257	CYS	2.4
1	B	180	HIS	2.3
1	B	198	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1BF9	B	401	35/35	0.92	0.10	40,50,74,120	0
2	A1BF9	A	401	35/35	0.94	0.10	42,50,74,120	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.