



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 02:33 PM EDT

PDB ID : 9EE0 / pdb\_00009ee0  
Title : HIV CA - GLFG peptide (9 mM)  
Authors : Melcak, I.; Sarafianos, S.G.  
Deposited on : 2024-11-18  
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.48.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






















Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1351 (3.10-3.10)
Clashscore	180529	1454 (3.10-3.10)
Ramachandran outliers	177936	1391 (3.10-3.10)
Sidechain outliers	177891	1391 (3.10-3.10)
RSRZ outliers	164620	1351 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	 3% 85% 8% 7%
1	B	231	 % 82% 8% 10%
1	C	231	 2% 84% 7% 8%
1	D	231	 3% 79% 10% 12%
1	E	231	 3% 82% 7% 11%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	231	 87% 6% 7%
1	G	231	 86% 8% 6%
1	H	231	 83% 7% 10%
1	I	231	 84% 9% 7%
1	J	231	 82% 11% 6%
1	K	231	 78% 10% 12%
1	L	231	 82% 5% 13%
2	M	9	 11% 11% 78%
2	N	9	 11% 89%
2	O	9	 22% 67% 33%
2	P	9	 11% 78% 22%
2	Q	9	 33% 67%
2	R	9	 22% 44% 11% 44%
2	S	9	 11% 56% 11% 33%
2	T	9	 22% 78%
2	U	9	 33% 67%
2	V	9	 44% 56% 11% 33%
2	W	9	 22% 56% 22% 22%
2	X	9	 22% 78%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 19394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein p24.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	214	Total 1600	C 1011	N 278	O 297	S 14	0	0	0
1	B	208	Total 1587	C 999	N 276	O 298	S 14	0	0	0
1	C	212	Total 1622	C 1018	N 284	O 306	S 14	0	0	0
1	D	204	Total 1534	C 964	N 266	O 290	S 14	0	0	0
1	E	205	Total 1561	C 985	N 272	O 291	S 13	0	0	0
1	F	215	Total 1629	C 1025	N 283	O 307	S 14	0	0	0
1	G	218	Total 1650	C 1036	N 290	O 310	S 14	0	0	0
1	H	208	Total 1583	C 996	N 277	O 296	S 14	0	0	0
1	I	214	Total 1603	C 1014	N 277	O 298	S 14	0	0	0
1	J	216	Total 1645	C 1032	N 290	O 309	S 14	0	0	0
1	K	203	Total 1529	C 963	N 264	O 288	S 14	0	0	0
1	L	202	Total 1515	C 955	N 260	O 287	S 13	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	14	CYS	ALA	engineered mutation	UNP P12493
A	45	CYS	GLU	engineered mutation	UNP P12493
A	184	ALA	TRP	engineered mutation	UNP P12493
A	185	ALA	MET	engineered mutation	UNP P12493
B	14	CYS	ALA	engineered mutation	UNP P12493

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	45	CYS	GLU	engineered mutation	UNP P12493
B	184	ALA	TRP	engineered mutation	UNP P12493
B	185	ALA	MET	engineered mutation	UNP P12493
C	14	CYS	ALA	engineered mutation	UNP P12493
C	45	CYS	GLU	engineered mutation	UNP P12493
C	184	ALA	TRP	engineered mutation	UNP P12493
C	185	ALA	MET	engineered mutation	UNP P12493
D	14	CYS	ALA	engineered mutation	UNP P12493
D	45	CYS	GLU	engineered mutation	UNP P12493
D	184	ALA	TRP	engineered mutation	UNP P12493
D	185	ALA	MET	engineered mutation	UNP P12493
E	14	CYS	ALA	engineered mutation	UNP P12493
E	45	CYS	GLU	engineered mutation	UNP P12493
E	184	ALA	TRP	engineered mutation	UNP P12493
E	185	ALA	MET	engineered mutation	UNP P12493
F	14	CYS	ALA	engineered mutation	UNP P12493
F	45	CYS	GLU	engineered mutation	UNP P12493
F	184	ALA	TRP	engineered mutation	UNP P12493
F	185	ALA	MET	engineered mutation	UNP P12493
G	14	CYS	ALA	engineered mutation	UNP P12493
G	45	CYS	GLU	engineered mutation	UNP P12493
G	184	ALA	TRP	engineered mutation	UNP P12493
G	185	ALA	MET	engineered mutation	UNP P12493
H	14	CYS	ALA	engineered mutation	UNP P12493
H	45	CYS	GLU	engineered mutation	UNP P12493
H	184	ALA	TRP	engineered mutation	UNP P12493
H	185	ALA	MET	engineered mutation	UNP P12493
I	14	CYS	ALA	engineered mutation	UNP P12493
I	45	CYS	GLU	engineered mutation	UNP P12493
I	184	ALA	TRP	engineered mutation	UNP P12493
I	185	ALA	MET	engineered mutation	UNP P12493
J	14	CYS	ALA	engineered mutation	UNP P12493
J	45	CYS	GLU	engineered mutation	UNP P12493
J	184	ALA	TRP	engineered mutation	UNP P12493
J	185	ALA	MET	engineered mutation	UNP P12493
K	14	CYS	ALA	engineered mutation	UNP P12493
K	45	CYS	GLU	engineered mutation	UNP P12493
K	184	ALA	TRP	engineered mutation	UNP P12493
K	185	ALA	MET	engineered mutation	UNP P12493
L	14	CYS	ALA	engineered mutation	UNP P12493
L	45	CYS	GLU	engineered mutation	UNP P12493
L	184	ALA	TRP	engineered mutation	UNP P12493

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
L	185	ALA	MET	engineered mutation	UNP P12493

- Molecule 2 is a protein called GLFG peptide.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	M	2	Total	C	N	O	0	0	0
			19	15	2	2			
2	O	6	Total	C	N	O	0	0	0
			37	24	6	7			
2	P	7	Total	C	N	O	0	0	0
			41	27	7	7			
2	Q	3	Total	C	N	O	0	0	0
			23	17	3	3			
2	R	5	Total	C	N	O	0	0	0
			32	22	5	5			
2	S	6	Total	C	N	O	0	0	0
			37	25	6	6			
2	T	2	Total	C	N	O	0	0	0
			15	11	2	2			
2	U	3	Total	C	N	O	0	0	0
			23	17	3	3			
2	V	6	Total	C	N	O	0	0	0
			36	24	6	6			
2	W	7	Total	C	N	O	0	0	0
			41	27	7	7			
2	X	2	Total	C	N	O	0	0	0
			19	15	2	2			
2	N	1	Total	C	N	O	0	0	0
			11	9	1	1			


- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	H	1	Total	Cl	0	0
			1	1		




- Molecule 1: Capsid protein p24

Chain E: 





- Molecule 1: Capsid protein p24

Chain F: 





- Molecule 1: Capsid protein p24

Chain G: 





- Molecule 1: Capsid protein p24

Chain H: 





- Molecule 1: Capsid protein p24

Chain I: 





- Molecule 1: Capsid protein p24



Chain J: 




- Molecule 1: Capsid protein p24




- Molecule 1: Capsid protein p24





- Molecule 2: GLFG peptide




- Molecule 2: GLFG peptide





- Molecule 2: GLFG peptide




- Molecule 2: GLFG peptide




- Molecule 2: GLFG peptide




- Molecule 2: GLFG peptide





- Molecule 2: GLFG peptide




- Molecule 2: GLFG peptide



- Molecule 2: GLFG peptide



- Molecule 2: GLFG peptide



- Molecule 2: GLFG peptide

Chain X:  22% 78%

ASP	GLY	L5	F6	GLY
SER	GLY			SER
GLY	GLY			LYS
LEU				

- Molecule 2: GLFG peptide

Chain N:  11% 89%

ASP	GLY	F6	GLY
SER	GLY		SER
GLY	LEU		LYS

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.06Å 135.22Å 205.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.90 – 3.10 33.90 – 3.10	Depositor EDS
% Data completeness (in resolution range)	95.5 (33.90-3.10) 95.6 (33.90-3.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.15	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 3.12Å)	Xtrriage
Refinement program	PHENIX 1.21_5207	Depositor
R, $R_{free}$	0.229 , 0.255 0.229 , 0.255	Depositor DCC
$R_{free}$ test set	3346 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.6	Xtrriage
Anisotropy	0.185	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 27.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.038 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	19394	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.09	0/1636	0.24	0/2230
1	B	0.08	0/1618	0.21	0/2197
1	C	0.09	0/1655	0.22	0/2250
1	D	0.08	0/1567	0.21	0/2133
1	E	0.08	0/1593	0.21	0/2166
1	F	0.08	0/1664	0.21	0/2265
1	G	0.08	0/1685	0.20	0/2294
1	H	0.09	0/1619	0.21	0/2206
1	I	0.09	0/1639	0.21	0/2234
1	J	0.09	0/1680	0.23	0/2283
1	K	0.09	0/1562	0.22	0/2126
1	L	0.08	0/1546	0.21	0/2108
2	M	0.23	0/19	0.78	0/24
2	N	0.05	0/11	0.03	0/13
2	O	0.07	0/37	0.12	0/47
2	P	0.09	0/41	0.24	0/53
2	Q	0.12	0/23	0.15	0/29
2	R	0.05	0/32	0.15	0/41
2	S	0.05	0/37	0.09	0/48
2	T	0.05	0/15	0.13	0/18
2	U	0.03	0/23	0.08	0/29
2	V	0.05	0/36	0.12	0/46
2	W	0.05	0/41	0.12	0/53
2	X	0.07	0/19	0.15	0/24
All	All	0.09	0/19798	0.22	0/26917

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1600	0	1550	10	0
1	B	1587	0	1553	13	0
1	C	1622	0	1584	10	0
1	D	1534	0	1454	11	0
1	E	1561	0	1529	9	0
1	F	1629	0	1577	9	0
1	G	1650	0	1605	12	0
1	H	1583	0	1535	10	0
1	I	1603	0	1552	13	0
1	J	1645	0	1603	16	0
1	K	1529	0	1458	14	0
1	L	1515	0	1447	8	0
2	M	19	0	19	2	0
2	N	11	0	8	1	0
2	O	37	0	33	0	0
2	P	41	0	32	0	0
2	Q	23	0	22	0	0
2	R	32	0	27	1	0
2	S	37	0	29	1	0
2	T	15	0	11	0	0
2	U	23	0	22	0	0
2	V	36	0	30	1	0
2	W	41	0	32	2	0
2	X	19	0	19	0	0
3	A	1	0	0	0	0
3	H	1	0	0	0	0
All	All	19394	0	18731	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:70:LYS:HZ1	2:W:9:LYS:HA	1.61	0.65

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:65:ALA:HB1	1:K:141:ILE:HD13	1.80	0.63
1:D:5:ASN:HB3	1:D:9:GLN:H	1.64	0.62
1:J:26:VAL:HG21	1:J:39:MET:HG2	1.81	0.62
1:C:26:VAL:HG21	1:C:39:MET:HG2	1.82	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/231 (91%)	208 (99%)	2 (1%)	0	100	100
1	B	202/231 (87%)	200 (99%)	2 (1%)	0	100	100
1	C	208/231 (90%)	203 (98%)	5 (2%)	0	100	100
1	D	198/231 (86%)	196 (99%)	2 (1%)	0	100	100
1	E	199/231 (86%)	194 (98%)	5 (2%)	0	100	100
1	F	211/231 (91%)	205 (97%)	6 (3%)	0	100	100
1	G	214/231 (93%)	206 (96%)	8 (4%)	0	100	100
1	H	204/231 (88%)	201 (98%)	3 (2%)	0	100	100
1	I	210/231 (91%)	206 (98%)	4 (2%)	0	100	100
1	J	212/231 (92%)	204 (96%)	8 (4%)	0	100	100
1	K	197/231 (85%)	192 (98%)	5 (2%)	0	100	100
1	L	196/231 (85%)	192 (98%)	4 (2%)	0	100	100
2	O	4/9 (44%)	4 (100%)	0	0	100	100
2	P	5/9 (56%)	5 (100%)	0	0	100	100
2	Q	1/9 (11%)	1 (100%)	0	0	100	100
2	R	3/9 (33%)	3 (100%)	0	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	S	4/9 (44%)	3 (75%)	1 (25%)	0	100	100
2	U	1/9 (11%)	1 (100%)	0	0	100	100
2	V	4/9 (44%)	4 (100%)	0	0	100	100
2	W	5/9 (56%)	5 (100%)	0	0	100	100
All	All	2488/2844 (88%)	2433 (98%)	55 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/193 (85%)	163 (99%)	1 (1%)	84	91
1	B	167/193 (86%)	167 (100%)	0	100	100
1	C	172/193 (89%)	171 (99%)	1 (1%)	84	91
1	D	157/193 (81%)	155 (99%)	2 (1%)	65	82
1	E	164/193 (85%)	164 (100%)	0	100	100
1	F	169/193 (88%)	169 (100%)	0	100	100
1	G	172/193 (89%)	170 (99%)	2 (1%)	67	83
1	H	167/193 (86%)	167 (100%)	0	100	100
1	I	165/193 (86%)	165 (100%)	0	100	100
1	J	173/193 (90%)	172 (99%)	1 (1%)	84	91
1	K	158/193 (82%)	158 (100%)	0	100	100
1	L	156/193 (81%)	154 (99%)	2 (1%)	65	82
2	M	2/6 (33%)	1 (50%)	1 (50%)	0	0
2	N	1/6 (17%)	1 (100%)	0	100	100
2	O	3/6 (50%)	3 (100%)	0	100	100
2	P	2/6 (33%)	2 (100%)	0	100	100
2	Q	2/6 (33%)	2 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	R	2/6 (33%)	2 (100%)	0	100	100
2	S	2/6 (33%)	2 (100%)	0	100	100
2	T	1/6 (17%)	1 (100%)	0	100	100
2	U	2/6 (33%)	2 (100%)	0	100	100
2	V	2/6 (33%)	2 (100%)	0	100	100
2	W	2/6 (33%)	2 (100%)	0	100	100
2	X	2/6 (33%)	2 (100%)	0	100	100
All	All	2007/2388 (84%)	1997 (100%)	10 (0%)	86	92

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	136	LEU
1	L	144	MET
2	M	6	PHE
1	D	175	GLU
1	G	70	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	12	HIS
1	L	67	GLN
1	K	121	ASN
1	F	67	GLN
1	J	4	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	214/231 (92%)	-0.12	7 (3%) 49 30	34, 55, 102, 130	0
1	B	208/231 (90%)	-0.12	3 (1%) 73 56	36, 57, 93, 135	0
1	C	212/231 (91%)	-0.12	4 (1%) 66 47	32, 55, 103, 132	0
1	D	204/231 (88%)	0.02	8 (3%) 44 26	34, 56, 128, 157	0
1	E	205/231 (88%)	-0.12	6 (2%) 54 34	34, 55, 97, 127	0
1	F	215/231 (93%)	-0.10	1 (0%) 87 75	34, 59, 115, 123	0
1	G	218/231 (94%)	-0.10	3 (1%) 73 56	42, 63, 112, 144	0
1	H	208/231 (90%)	0.14	9 (4%) 40 24	41, 69, 127, 198	0
1	I	214/231 (92%)	0.15	7 (3%) 49 30	38, 64, 108, 158	0
1	J	216/231 (93%)	-0.01	12 (5%) 31 19	35, 53, 119, 153	0
1	K	203/231 (87%)	0.03	8 (3%) 44 26	37, 59, 131, 170	0
1	L	202/231 (87%)	0.01	8 (3%) 43 25	41, 67, 103, 164	0
2	M	2/9 (22%)	2.55	1 (50%) 0 0	83, 83, 83, 125	0
2	N	1/9 (11%)	1.23	0 100 100	54, 54, 54, 54	1 (100%)
2	O	6/9 (66%)	1.48	2 (33%) 1 0	66, 71, 86, 98	0
2	P	7/9 (77%)	1.39	1 (14%) 7 4	45, 65, 72, 83	7 (100%)
2	Q	3/9 (33%)	1.23	0 100 100	59, 59, 73, 75	0
2	R	5/9 (55%)	1.41	2 (40%) 1 0	60, 60, 75, 84	0
2	S	6/9 (66%)	1.24	1 (16%) 5 3	55, 65, 68, 73	0
2	T	2/9 (22%)	2.28	2 (100%) 0 0	64, 64, 64, 79	2 (100%)
2	U	3/9 (33%)	0.57	0 100 100	66, 66, 68, 72	0
2	V	6/9 (66%)	2.32	4 (66%) 0 0	53, 67, 86, 92	0
2	W	7/9 (77%)	1.45	2 (28%) 1 1	59, 76, 95, 106	0
2	X	2/9 (22%)	0.65	0 100 100	49, 49, 49, 59	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
All	All	2569/2880 (89%)	0.00	91 (3%) 47 28	32, 61, 114, 198	10 (0%)

The worst 5 of 91 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	217	ALA	5.2
1	E	183	ASN	4.9
1	C	219	GLN	4.7
1	I	82	ARG	4.4
2	P	8	SER	4.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	A	301	1/1	0.87	0.15	71,71,71,71	0
3	CL	H	301	1/1	0.89	0.09	70,70,70,70	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.