



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 30, 2025 – 02:55 am BST

PDB ID : 9ESN / pdb\_00009esn  
Title : CDK2-cyclin A in complex with FragLite 28  
Authors : Hope, I.; Martin, M.P.; Waring, M.J.; Noble, M.E.M.; Endicott, J.A.; Tatum, N.J.  
Deposited on : 2024-03-26  
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : ?? (??), CSD ??CSD?? (????)  
Xtrriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	 5% 82% 15% ..
1	C	302	 6% 83% 14% .
2	B	268	 % 86% 11% .
2	D	268	 4% 90% 8% .

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 9374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	298	Total	C	N	O	P	S	0	0	0
			2385	1547	403	426	1	8			
1	C	297	Total	C	N	O	P	S	0	0	0
			2388	1550	404	425	1	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P24941
A	-2	PRO	-	expression tag	UNP P24941
A	-1	GLY	-	expression tag	UNP P24941
A	0	SER	-	expression tag	UNP P24941
C	-3	GLY	-	expression tag	UNP P24941
C	-2	PRO	-	expression tag	UNP P24941
C	-1	GLY	-	expression tag	UNP P24941
C	0	SER	-	expression tag	UNP P24941

- Molecule 2 is a protein called Cyclin-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	262	Total	C	N	O	S	0	1	0
			2118	1370	345	393	10			
2	D	262	Total	C	N	O	S	0	1	0
			2118	1370	345	393	10			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	171	GLY	-	expression tag	UNP P30274
B	433	HIS	-	expression tag	UNP P30274
B	434	HIS	-	expression tag	UNP P30274
B	435	HIS	-	expression tag	UNP P30274

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	436	HIS	-	expression tag	UNP P30274
B	437	HIS	-	expression tag	UNP P30274
B	438	HIS	-	expression tag	UNP P30274
D	171	GLY	-	expression tag	UNP P30274
D	433	HIS	-	expression tag	UNP P30274
D	434	HIS	-	expression tag	UNP P30274
D	435	HIS	-	expression tag	UNP P30274
D	436	HIS	-	expression tag	UNP P30274
D	437	HIS	-	expression tag	UNP P30274
D	438	HIS	-	expression tag	UNP P30274

- Molecule 3 is 4-bromo-1-(2-hydroxyethyl)pyridin-2(1H)-one (CCD ID: UUS) (formula: C<sub>7</sub>H<sub>8</sub>BrNO<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	Br	C	N	O		
3	A	1	Total	Br	C	N	O	0	0
			11	1	7	1	2		
3	B	1	Total	Br	C	N	O	0	0
			11	1	7	1	2		
3	C	1	Total	Br	C	N	O	0	0
			11	1	7	1	2		
3	C	1	Total	Br	C	N	O	0	0
			11	1	7	1	2		
3	D	1	Total	Br	C	N	O	0	0
			11	1	7	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	106	Total 106	O 106	0	0
4	B	113	Total 113	O 113	0	0
4	C	51	Total 51	O 51	0	0
4	D	40	Total 40	O 40	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclin-dependent kinase 2




- Molecule 1: Cyclin-dependent kinase 2



- Molecule 2: Cyclin-A2



- Molecule 2: Cyclin-A2



HIS

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.18Å 134.16Å 147.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	99.58 – 2.39 99.38 – 2.39	Depositor EDS
% Data completeness (in resolution range)	93.5 (99.58-2.39) 93.4 (99.38-2.39)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.91 (at 2.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.209 , 0.213 0.196 , 0.199	Depositor DCC
$R_{free}$ test set	2830 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.7	Xtrriage
Anisotropy	0.130	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9374	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, UUS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/2435	0.84	0/3304
1	C	0.42	0/2438	0.81	0/3308
2	B	0.42	0/2168	0.86	1/2948 (0.0%)
2	D	0.41	0/2168	0.79	0/2948
All	All	0.42	0/9209	0.83	1/12508 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	4
2	B	0	1
2	D	0	2
All	All	0	8

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	346	PRO	N-CA-CB	-10.63	90.54	103.30

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	150	ARG	Sidechain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
2	B	400	ARG	Sidechain
1	C	150	ARG	Sidechain
1	C	217	ARG	Sidechain
1	C	245	ARG	Sidechain
1	C	260	ARG	Sidechain
2	D	293	ARG	Sidechain
2	D	410	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2385	0	2425	52	0
1	C	2388	0	2430	36	0
2	B	2118	0	2131	24	0
2	D	2118	0	2131	16	0
3	A	11	0	0	0	0
3	B	11	0	0	0	0
3	C	22	0	0	0	0
3	D	11	0	0	0	0
4	A	106	0	0	1	0
4	B	113	0	0	4	0
4	C	51	0	0	0	0
4	D	40	0	0	1	0
All	All	9374	0	9117	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:GLU:HG3	1:C:73:GLU:OE1	1.45	1.15
1:A:59:ASN:ND2	1:A:65:LYS:HE3	1.62	1.13
1:C:9:LYS:HE3	1:C:17:VAL:HG23	1.19	1.12
1:A:59:ASN:HD21	1:A:65:LYS:HE3	1.36	0.88

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:9:LYS:HE3	1:C:17:VAL:CG2	2.04	0.87
1:C:15:TYR:HB3	1:C:35:ILE:HG12	1.60	0.83
2:D:223:GLU:OE2	2:D:412:LYS:HD2	1.78	0.83
1:C:9:LYS:CE	1:C:17:VAL:HG23	2.07	0.82
1:C:197:VAL:HG11	1:C:252:VAL:HG13	1.62	0.82
1:C:294:PRO:HB2	1:C:296:LEU:HD13	1.65	0.78
1:A:71:HIS:HE2	2:B:296:HIS:HD1	1.31	0.78
2:D:223:GLU:OE2	2:D:412:LYS:HE3	1.86	0.75
2:D:223:GLU:OE2	2:D:412:LYS:CE	2.38	0.72
1:A:13:GLY:HA3	1:A:15:TYR:CE2	2.25	0.72
1:A:34:LYS:HD2	1:A:75:LYS:HD3	1.73	0.71
1:A:59:ASN:HD22	1:A:65:LYS:HE3	1.55	0.69
1:A:72:THR:HG22	1:A:73:GLU:H	1.58	0.68
1:A:72:THR:HG22	1:A:73:GLU:N	2.08	0.68
2:D:223:GLU:OE2	2:D:412:LYS:CD	2.40	0.68
1:A:177:CYS:SG	1:A:233:MET:CE	2.83	0.66
2:D:327:CYS:HB3	2:D:419:HIS:CE1	2.31	0.66
1:C:252:VAL:HG12	1:C:255:LEU:HB2	1.77	0.65
1:A:175:LEU:CD1	1:A:233:MET:HE1	2.28	0.63
1:C:197:VAL:HG11	1:C:252:VAL:CG1	2.28	0.63
1:A:175:LEU:HD12	1:A:233:MET:HE1	1.81	0.62
1:A:139:GLY:HA2	1:A:294:PRO:HD3	1.80	0.62
1:A:13:GLY:HA3	1:A:15:TYR:CZ	2.34	0.61
1:C:60:HIS:CG	1:C:61:PRO:HD2	2.36	0.60
1:A:72:THR:CG2	1:A:73:GLU:H	2.15	0.60
1:A:121:HIS:O	1:A:122:ARG:HG3	2.03	0.59
1:A:35:ILE:HD12	1:A:78:LEU:HD13	1.85	0.59
2:D:233:HIS:HD2	4:D:819:HOH:O	1.86	0.58
1:A:59:ASN:ND2	1:A:65:LYS:CE	2.53	0.58
2:B:338:GLU:HG3	4:B:808:HOH:O	2.03	0.57
2:B:411:GLU:HA	2:B:414:LYS:HE2	1.85	0.57
1:C:91:MET:HE2	1:C:130:PRO:HB3	1.85	0.57
1:C:177:CYS:SG	1:C:233:MET:SD	3.03	0.56
1:C:139:GLY:HA2	1:C:294:PRO:HD3	1.89	0.55
2:B:395:HIS:HE1	2:B:427:PRO:O	1.90	0.55
1:A:72:THR:CG2	1:A:73:GLU:N	2.71	0.54
1:C:15:TYR:CB	1:C:35:ILE:HG12	2.35	0.53
1:A:36:ARG:O	1:A:40:GLU:HG2	2.09	0.52
1:C:15:TYR:CD2	1:C:33:LYS:HD3	2.45	0.52
1:C:215:ILE:HG23	1:C:219:LEU:HD12	1.91	0.51
1:A:71:HIS:NE2	2:B:296:HIS:ND1	2.45	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:TRP:CD1	1:A:204:PRO:HA	2.47	0.50
1:C:9:LYS:HG3	1:C:17:VAL:CG2	2.42	0.50
1:A:71:HIS:HE2	2:B:296:HIS:CE1	2.30	0.49
2:D:411:GLU:HA	2:D:414:LYS:HD2	1.95	0.48
1:A:177:CYS:SG	1:A:233:MET:HG2	2.53	0.48
2:B:386:THR:HB	4:B:859:HOH:O	2.14	0.48
1:C:88:LYS:HB2	1:C:130:PRO:HB2	1.96	0.48
1:C:121:HIS:O	1:C:122:ARG:HG3	2.14	0.48
1:A:288:ASP:OD1	1:A:288:ASP:N	2.46	0.47
2:D:414:LYS:HG3	2:D:422:SER:OG	2.14	0.47
2:B:183:HIS:HB2	2:B:317:GLN:HE22	1.78	0.47
1:A:73:GLU:OE2	1:C:0:SER:N	2.48	0.47
1:A:9:LYS:HG3	1:A:17:VAL:HG23	1.96	0.47
1:C:137:THR:O	1:C:293:VAL:HG13	2.15	0.47
2:B:407:GLN:O	2:B:411:GLU:HG2	2.15	0.47
1:C:91:MET:CE	1:C:130:PRO:HB3	2.44	0.47
1:A:9:LYS:HG3	1:A:17:VAL:CG2	2.44	0.47
1:A:59:ASN:HD22	1:A:65:LYS:CE	2.23	0.46
1:C:91:MET:HE1	1:C:130:PRO:CG	2.45	0.46
2:B:233:HIS:HE1	4:B:808:HOH:O	1.98	0.46
1:A:56:LYS:HD3	2:B:305[B]:ASP:OD1	2.16	0.46
1:C:252:VAL:CG1	1:C:255:LEU:HB2	2.46	0.46
1:A:121:HIS:C	1:A:122:ARG:HG3	2.36	0.45
1:A:177:CYS:SG	1:A:233:MET:HE2	2.55	0.45
1:A:177:CYS:HB2	4:A:417:HOH:O	2.15	0.45
1:C:60:HIS:CD2	1:C:61:PRO:HD2	2.51	0.45
2:D:412:LYS:HG2	2:D:413:TYR:CD1	2.52	0.45
2:B:383:THR:OG1	2:B:385:GLU:HB3	2.17	0.44
1:A:248:PHE:HE2	1:A:263:LEU:HD23	1.82	0.44
1:A:60:HIS:CG	1:A:61:PRO:HD2	2.52	0.44
2:B:346:PRO:O	2:B:349:LYS:HG2	2.18	0.44
1:A:177:CYS:SG	1:A:233:MET:HE3	2.58	0.43
1:C:175:LEU:HA	1:C:235:ASP:HB2	2.00	0.43
2:D:388:LYS:HB3	2:D:389:PRO:HD3	2.00	0.43
1:C:78:LEU:HD23	1:C:78:LEU:N	2.33	0.43
1:C:273:LYS:HD3	1:C:273:LYS:HA	1.79	0.43
1:A:71:HIS:CD2	1:A:76:LEU:HD13	2.54	0.43
2:D:347:TYR:OH	2:D:394:LEU:HA	2.18	0.43
1:A:88:LYS:HB2	1:A:130:PRO:HB2	2.01	0.43
1:A:71:HIS:CE1	2:B:296:HIS:HD1	2.33	0.43
2:B:215:VAL:O	2:B:219:VAL:HG23	2.19	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:TYR:CD1	1:A:16:GLY:N	2.87	0.42
2:B:183:HIS:HD2	4:B:820:HOH:O	2.02	0.42
2:D:396:GLN:HE21	2:D:400:ARG:NE	2.17	0.42
2:B:401:ALA:N	2:B:402:PRO:CD	2.83	0.42
1:A:161:HIS:CG	1:A:162:GLU:N	2.88	0.42
1:C:231:THR:HA	1:C:236:TYR:CD1	2.55	0.42
1:C:255:LEU:HD12	1:C:255:LEU:HA	1.82	0.42
2:D:329:VAL:HG22	2:D:364:LEU:HA	2.02	0.42
1:A:37:LEU:HD13	2:B:296:HIS:CE1	2.55	0.42
1:C:247:ASP:HB2	1:C:250:LYS:HE2	2.02	0.42
1:C:250:LYS:O	1:C:253:PRO:HD3	2.20	0.42
1:A:101:LEU:N	1:A:102:PRO:CD	2.83	0.41
2:D:335:PHE:HB2	2:D:413:TYR:CD2	2.55	0.41
2:D:384:LEU:HD23	2:D:384:LEU:HA	1.82	0.41
1:C:150:ARG:HD2	1:C:160:TPO:O1P	2.19	0.41
2:B:171:GLY:O	2:B:176:PRO:HD3	2.20	0.41
2:B:388:LYS:HB3	2:B:389:PRO:HD3	2.02	0.41
1:A:37:LEU:O	1:A:40:GLU:HB2	2.20	0.41
1:C:121:HIS:C	1:C:122:ARG:HG3	2.41	0.41
1:A:91:MET:HA	1:A:95:ALA:HB3	2.03	0.41
1:A:37:LEU:O	1:A:40:GLU:N	2.47	0.41
2:B:293:ARG:HB3	1:C:25:LEU:HD11	2.03	0.41
2:B:345:ASP:HA	2:B:346:PRO:HA	1.76	0.41
2:D:223:GLU:CD	2:D:412:LYS:HD2	2.41	0.41
1:A:175:LEU:HD13	1:A:233:MET:HE1	2.03	0.40
1:A:88:LYS:O	1:A:88:LYS:HD3	2.21	0.40
1:A:223:ASP:H	1:A:226:VAL:HG12	1.87	0.40
1:A:230:VAL:HA	1:A:233:MET:SD	2.61	0.40
1:A:202:LEU:HD23	1:A:203:PHE:CE2	2.56	0.40
2:B:223:GLU:HG2	2:B:412:LYS:HD2	2.02	0.40
1:A:177:CYS:SG	1:A:233:MET:SD	3.19	0.40
2:B:235:ALA:O	2:B:239:ILE:HG13	2.21	0.40
1:C:36:ARG:HD3	1:C:36:ARG:HA	1.88	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/302 (98%)	287 (97%)	7 (2%)	1 (0%)	37	51
1	C	294/302 (97%)	286 (97%)	7 (2%)	1 (0%)	37	51
2	B	261/268 (97%)	255 (98%)	6 (2%)	0	100	100
2	D	261/268 (97%)	257 (98%)	4 (2%)	0	100	100
All	All	1111/1140 (98%)	1085 (98%)	24 (2%)	2 (0%)	44	59

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	164	VAL
1	A	164	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/264 (98%)	257 (99%)	3 (1%)	67	82
1	C	261/264 (99%)	257 (98%)	4 (2%)	60	77
2	B	235/240 (98%)	232 (99%)	3 (1%)	65	81
2	D	235/240 (98%)	235 (100%)	0	100	100
All	All	991/1008 (98%)	981 (99%)	10 (1%)	73	86

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	ARG
1	A	150	ARG
1	A	248	PHE
2	B	223	GLU
2	B	346	PRO
2	B	391	LEU
1	C	122	ARG
1	C	150	ARG
1	C	199	ARG
1	C	225	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	74	ASN
1	A	85	GLN
2	B	233	HIS
2	B	254	GLN
2	B	317	GLN
2	B	378	GLN
2	B	395	HIS
2	B	396	GLN
2	B	425	ASN
1	C	59	ASN
1	C	265	GLN
2	D	233	HIS
2	D	254	GLN
2	D	317	GLN
2	D	370	GLN
2	D	396	GLN
2	D	419	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

5 ligands are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.



There are no chirality outliers.



There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	297/302 (98%)	-0.11	16 (5%) 32 30	36, 50, 111, 171	0
1	C	296/302 (98%)	0.27	17 (5%) 30 28	43, 65, 119, 144	0
2	B	262/268 (97%)	-0.25	3 (1%) 77 75	27, 53, 80, 123	1 (0%)
2	D	262/268 (97%)	0.39	11 (4%) 41 38	30, 69, 111, 150	1 (0%)
All	All	1117/1140 (97%)	0.07	47 (4%) 41 38	27, 59, 111, 171	2 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	432	VAL	7.0
2	B	432	VAL	6.8
1	A	-2	PRO	4.3
1	A	-3	GLY	4.1
1	A	96	LEU	3.8
1	C	14	THR	3.7
1	A	15	TYR	3.5
2	D	424	LEU	3.5
1	A	94	SER	3.4
1	A	97	THR	3.1
2	D	430	LEU	3.1
1	C	233	MET	3.0
1	A	292	PRO	3.0
1	A	0	SER	2.9
1	C	296	LEU	2.8
2	D	329	VAL	2.8
1	C	206	ASP	2.6
1	A	293	VAL	2.6
1	C	248	PHE	2.5
1	A	72	THR	2.5
1	C	237	LYS	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	426	PRO	2.5
1	C	239	SER	2.5
1	C	295	HIS	2.5
1	C	217	ARG	2.5
1	C	241	PRO	2.5
1	A	294	PRO	2.4
1	A	98	GLY	2.4
2	D	398	TYR	2.4
1	A	71	HIS	2.4
1	C	238	PRO	2.3
2	D	324	PRO	2.3
1	A	93	ALA	2.2
1	A	41	THR	2.2
2	B	305[A]	ASP	2.2
2	B	223	GLU	2.2
1	C	226	VAL	2.2
1	C	236	TYR	2.2
1	A	290	THR	2.2
2	D	429	THR	2.2
1	C	13	GLY	2.2
2	D	392	LEU	2.2
1	C	15	TYR	2.1
2	D	418	TYR	2.1
2	D	427	PRO	2.1
1	C	242	LYS	2.1
1	C	240	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	C	160	11/12	0.97	0.07	48,54,61,64	0
1	TPO	A	160	11/12	0.98	0.05	42,48,52,53	0

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	UUS	A	301	11/11	0.84	0.19	67,79,100,108	0
3	UUS	C	502	11/11	0.85	0.15	119,126,143,178	0
3	UUS	C	501	11/11	0.94	0.14	68,87,103,107	0
3	UUS	D	701	11/11	0.94	0.13	76,87,105,106	0
3	UUS	B	701	11/11	0.96	0.12	68,82,99,107	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around UUS C 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UUS C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UUS D 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around UUS B 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.