



wwPDB EM Validation Summary Report ⓘ

Mar 10, 2026 – 03:47 pm GMT

PDB ID : 9EVT / pdb_00009evt
EMDB ID : EMD-50014
Title : Structure of the small subunit of the flowering plant mitoribosome with the maturation factor RsgA
Authors : Waltz, F.; Skaltsogiannis, V.; Giege, P.
Deposited on : 2024-04-02
Resolution : 2.27 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
RNA backbone	6643	2191	-
Q-score	-	25397	3593 (1.77 - 2.77)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	88% (green), 8% (yellow), 4% (grey)
2	B	554	68% (green), 15% (yellow), 16% (grey), 1% (red)
3	C	362	81% (green), 11% (yellow), 8% (grey), 1% (red)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	501	
5	E	138	
6	F	157	
7	G	129	
8	H	383	
9	I	228	
10	J	304	
11	K	125	
12	L	154	
13	M	155	
14	N	414	
15	O	136	
16	P	110	
17	Q	237	
18	R	212	
19	S	100	
20	T	94	
21	U	192	
22	V	193	
23	W	483	
24	X	496	
25	Y	102	
26	Z	153	
27	7	496	
28	a	424	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
29	b	80	
30	c	128	
31	d	110	
32	e	383	
33	f	410	
34	h	384	
35	i	725	
36	k	153	
37	2	1560	

2 Entry composition [i](#)

There are 42 unique types of molecules in this entry. The entry contains 85715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosomal protein S2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	204	1611	1032	284	285	10	0	0

- Molecule 2 is a protein called Small ribosomal subunit protein uS3m.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	467	3877	2520	701	643	13	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	22	TRP	ARG	conflict	UNP A0A068BCX1
B	172	LEU	SER	conflict	UNP A0A068BCX1
B	202	LYS	ARG	conflict	UNP A0A068BCX1
B	296	LEU	SER	conflict	UNP A0A068BCX1
B	512	CYS	ARG	conflict	UNP A0A068BCX1
B	524	VAL	ALA	conflict	UNP A0A068BCX1
B	533	LEU	SER	conflict	UNP A0A068BCX1

- Molecule 3 is a protein called Ribosomal protein S4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	333	2806	1793	540	462	11	0	0

- Molecule 4 is a protein called S5 DRBM domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	417	3425	2146	603	664	12	0	0

- Molecule 5 is a protein called bS6m.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	101	Total	C	N	O	S	0	0
			823	535	145	138	5		

- Molecule 6 is a protein called uS7m,Small ribosomal subunit protein uS7m,Small ribosomal subunit protein uS7m.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	157	Total	C	N	O	S	0	0
			1254	793	244	213	4		

- Molecule 7 is a protein called 40S ribosomal protein S15a.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	128	Total	C	N	O	S	0	0
			1037	652	193	189	3		

- Molecule 8 is a protein called 30S ribosomal protein S9.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	211	Total	C	N	O	S	0	0
			1662	1042	306	309	5		

- Molecule 9 is a protein called Small ribosomal subunit protein uS10 domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	113	Total	C	N	O	S	0	0
			941	602	174	159	6		

- Molecule 10 is a protein called Ribosomal protein S11.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	127	Total	C	N	O	S	0	0
			993	612	196	181	4		

- Molecule 11 is a protein called Small ribosomal subunit protein uS12m.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	124	Total	C	N	O	S	0	0
			992	614	209	165	4		

- Molecule 12 is a protein called uS13m.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	L	109	864	523	183	153	5	0	0

- Molecule 13 is a protein called Small ribosomal subunit protein uS14m.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	M	101	843	524	174	140	5	0	0

- Molecule 14 is a protein called Small ribosomal subunit protein uS15c.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
14	N	116	941	593	170	171	7	0	0

- Molecule 15 is a protein called 30S ribosomal protein S16.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
15	O	110	869	550	169	144	6	0	0

- Molecule 16 is a protein called Small ribosomal subunit protein uS17c.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
16	P	91	717	455	135	124	3	0	0

- Molecule 17 is a protein called Small ribosomal subunit protein bS18c.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
17	Q	90	724	460	127	134	3	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Q	250	ILE	VAL	conflict	UNP A0A078FCZ9

- Molecule 18 is a protein called Small ribosomal subunit protein uS19m.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
18	R	166	1299	819	236	237	7	0	0

- Molecule 19 is a protein called Ribosomal protein S21.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
19	S	75	611	386	120	103	2	0	0

- Molecule 20 is a protein called 30S ribosomal protein S31, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
20	T	50	408	260	79	68	1	0	0

- Molecule 21 is a protein called Small ribosomal subunit protein mS23.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
21	U	127	1016	647	185	179	5	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
U	81	GLN	PRO	conflict	UNP A0A3P6DYP2

- Molecule 22 is a protein called mS26.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
22	V	164	1399	874	266	255	4	0	0

- Molecule 23 is a protein called Small ribosomal subunit protein mS29.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
23	W	380	3062	1955	535	559	13	0	0

- Molecule 24 is a protein called BnaC06g06470D protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
24	X	204	1613	1011	262	329	11	0	0

- Molecule 25 is a protein called MOSC domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
25	Y	98	793	495	159	137	2	0	0

- Molecule 26 is a protein called Mitochondrial 28S ribosomal protein S34.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
26	Z	81	660	428	125	105	2	0	0

- Molecule 27 is a protein called EngC GTPase domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
27	7	363	2878	1824	514	526	14	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
7	385	MET	GLN	conflict	UNP A0A0D3CYN0

- Molecule 28 is a protein called Small ribosomal subunit protein mS35 mitochondrial conserved domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
28	a	315	2486	1546	448	482	10	0	0

- Molecule 29 is a protein called IMS import disulfide relay-system CHCH-CHCH-like Cx9C domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
29	b	73	581	361	108	104	8	0	0

- Molecule 30 is a protein called Small ribosomal subunit protein mS38.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	c	26	Total	C	N	O	S	0	0
			232	145	54	32	1		

- Molecule 31 is a protein called IGR domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	d	78	Total	C	N	O	S	0	0
			616	403	110	102	1		

- Molecule 32 is a protein called mS45.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	e	246	Total	C	N	O	S	0	0
			2074	1310	374	380	10		

- Molecule 33 is a protein called mS47.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	f	375	Total	C	N	O	S	0	0
			2898	1825	484	569	20		

- Molecule 34 is a protein called (rape) hypothetical protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	h	310	Total	C	N	O	S	0	0
			2463	1557	438	455	13		

- Molecule 35 is a protein called PROP1-like PPR domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	i	306	Total	C	N	O	S	0	0
			2487	1567	419	487	14		

- Molecule 36 is a protein called RRM domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
36	k	22	Total	C	N	O	S	0	0
			177	113	34	27	3		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
k	9	ARG	LYS	conflict	UNP A0A3P6FEP3
k	15	LYS	ARG	conflict	UNP A0A3P6FEP3

- Molecule 37 is a RNA chain called 18S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
37	2	1560	33424	14925	6097	10842	1560	0	0


- Molecule 38 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		AltConf
38	K	1	Total 1	K 1	0
38	2	9	Total 9	K 9	0

- Molecule 39 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
39	T	1	Total 1	Mg 1	0
39	W	1	Total 1	Mg 1	0
39	7	1	Total 1	Mg 1	0
39	2	82	Total 82	Mg 82	0

- Molecule 40 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
40	W	1	31	10	5	13	3	0

- Molecule 41 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
			Total	Zn	
41	7	1	1	1	0

- Molecule 42 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
42	7	1	32	10	5	14	3	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Ribosomal protein S2

Chain A:  88% 8%



- Molecule 2: Small ribosomal subunit protein uS3m

Chain B:  68% 15% 16%





- Molecule 3: Ribosomal protein S4

Chain C:  81% 11% 8%






• Molecule 18: Small ribosomal subunit protein uS19m




• Molecule 19: Ribosomal protein S21



• Molecule 20: 30S ribosomal protein S31, mitochondrial





• Molecule 21: Small ribosomal subunit protein mS23



• Molecule 22: mS26






- Molecule 29: IMS import disulfide relay-system CHCH-CHCH-like Cx9C domain-containing protein




- Molecule 30: Small ribosomal subunit protein mS38



- Molecule 31: IGR domain-containing protein




- Molecule 32: mS45




- Molecule 33: mS47







• Molecule 36: RRM domain-containing protein



• Molecule 37: 18S rRNA





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	208855	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50.96	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	165000	Depositor
Image detector	TFS FALCON 4i (4k x 4k)	Depositor
Maximum map value	1.059	Depositor
Minimum map value	-0.572	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.023	Depositor
Recommended contour level	0.0377	Depositor
Map size (\AA)	524.87823, 524.87823, 524.87823	wwPDB
Map dimensions	588, 588, 588	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.89265007, 0.89265007, 0.89265007	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 2MG, UR3, 4OC, ATP, K, MG, G7M, MA6, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.12	0/1647	0.34	0/2227
2	B	0.23	0/3969	0.50	1/5320 (0.0%)
3	C	0.15	0/2859	0.38	1/3828 (0.0%)
4	D	0.15	0/3474	0.38	0/4647
5	E	0.11	0/838	0.32	0/1121
6	F	0.16	0/1225	0.42	0/1641
7	G	0.15	0/1054	0.36	0/1418
8	H	0.20	0/1687	0.49	0/2262
9	I	0.19	0/964	0.43	0/1293
10	J	0.12	0/1005	0.34	0/1346
11	K	0.14	0/1009	0.36	0/1345
12	L	0.14	0/874	0.36	0/1165
13	M	0.17	0/856	0.38	0/1134
14	N	0.13	0/958	0.34	0/1280
15	O	0.16	0/890	0.37	0/1196
16	P	0.17	0/730	0.44	1/984 (0.1%)
17	Q	0.12	0/733	0.32	0/978
18	R	0.17	0/1323	0.37	0/1772
19	S	0.15	0/614	0.33	0/809
20	T	0.16	0/416	0.37	0/547
21	U	0.39	2/1037 (0.2%)	0.55	1/1395 (0.1%)
22	V	0.21	0/1423	0.33	0/1895
23	W	0.17	0/3139	0.40	0/4250
24	X	0.22	0/1644	0.46	0/2216
25	Y	0.42	0/807	0.50	0/1077
26	Z	0.12	0/680	0.34	0/914
27	7	0.18	0/2930	0.45	0/3952
28	a	0.28	0/2523	0.60	5/3386 (0.1%)
29	b	0.19	0/589	0.54	0/785
30	c	0.11	0/233	0.27	0/296
31	d	0.16	0/628	0.33	0/840
32	e	0.15	0/2118	0.38	0/2828

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
33	f	0.17	0/2955	0.40	1/3999 (0.0%)
34	h	0.18	0/2511	0.44	0/3391
35	i	0.14	0/2532	0.36	0/3400
36	k	0.07	0/181	0.21	0/239
37	2	0.21	2/37258 (0.0%)	0.34	2/58090 (0.0%)
All	All	0.20	4/90313 (0.0%)	0.38	12/129266 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
37	2	1650	C	O3'-P	-21.93	1.28	1.61
37	2	921	G	O3'-P	14.79	1.83	1.61
21	U	42	ASN	CA-C	-5.67	1.45	1.52
21	U	40	ARG	CA-C	-5.59	1.45	1.53

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
37	2	1650	C	P-O3'-C3'	12.01	138.22	120.20
28	a	211	LYS	N-CA-C	-8.81	101.61	114.39
2	B	212	PRO	CA-N-CD	-8.71	99.81	112.00
16	P	3	PRO	CA-N-CD	-8.68	99.84	112.00
37	2	1650	C	O3'-P-O5'	8.60	116.90	104.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1611	0	1633	13	0
2	B	3877	0	4008	154	0
3	C	2806	0	2971	29	0
4	D	3425	0	3451	25	0
5	E	823	0	879	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	F	1254	0	1268	16	0
7	G	1037	0	1058	10	0
8	H	1662	0	1700	17	0
9	I	941	0	970	13	0
10	J	993	0	1034	27	0
11	K	992	0	1051	14	0
12	L	864	0	891	5	0
13	M	843	0	884	18	0
14	N	941	0	978	3	0
15	O	869	0	898	6	0
16	P	717	0	755	8	0
17	Q	724	0	744	22	0
18	R	1299	0	1307	8	0
19	S	611	0	687	18	0
20	T	408	0	444	2	0
21	U	1016	0	1062	19	0
22	V	1399	0	1427	15	0
23	W	3062	0	3029	35	0
24	X	1613	0	1559	66	0
25	Y	793	0	834	15	0
26	Z	660	0	675	7	0
27	7	2878	0	2946	41	0
28	a	2486	0	2376	109	0
29	b	581	0	592	21	0
30	c	232	0	278	2	0
31	d	616	0	674	5	0
32	e	2074	0	2074	16	0
33	f	2898	0	2852	80	0
34	h	2463	0	2488	66	0
35	i	2487	0	2450	21	0
36	k	177	0	179	2	0
37	2	33424	0	16844	214	0
38	2	9	0	0	0	0
38	K	1	0	0	0	0
39	2	82	0	0	0	0
39	7	1	0	0	0	0
39	T	1	0	0	0	0
39	W	1	0	0	0	0
40	W	31	0	12	0	0
41	7	1	0	0	0	0
42	7	32	0	12	0	0
All	All	85715	0	69974	882	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 882 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:231:ARG:HD3	24:X:426:GLU:CD	1.22	1.53
19:S:93:ILE:CG2	29:b:86:LEU:HD13	1.02	1.49
19:S:93:ILE:HG21	29:b:86:LEU:CD1	1.02	1.47
2:B:231:ARG:CD	24:X:426:GLU:OE1	1.66	1.43
34:h:302:HIS:CE1	34:h:356:SER:HB3	1.58	1.38

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	202/212 (95%)	198 (98%)	4 (2%)	0	100	100
2	B	457/554 (82%)	430 (94%)	24 (5%)	3 (1%)	19	22
3	C	329/362 (91%)	324 (98%)	5 (2%)	0	100	100
4	D	413/501 (82%)	396 (96%)	17 (4%)	0	100	100
5	E	99/138 (72%)	99 (100%)	0	0	100	100
6	F	146/157 (93%)	140 (96%)	6 (4%)	0	100	100
7	G	126/129 (98%)	124 (98%)	2 (2%)	0	100	100
8	H	207/383 (54%)	198 (96%)	8 (4%)	1 (0%)	25	30
9	I	111/228 (49%)	108 (97%)	3 (3%)	0	100	100
10	J	125/304 (41%)	119 (95%)	6 (5%)	0	100	100
11	K	122/125 (98%)	120 (98%)	2 (2%)	0	100	100
12	L	107/154 (70%)	104 (97%)	3 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
13	M	99/155 (64%)	99 (100%)	0	0	100	100
14	N	114/414 (28%)	114 (100%)	0	0	100	100
15	O	108/136 (79%)	106 (98%)	2 (2%)	0	100	100
16	P	89/110 (81%)	87 (98%)	2 (2%)	0	100	100
17	Q	88/237 (37%)	87 (99%)	1 (1%)	0	100	100
18	R	162/212 (76%)	157 (97%)	5 (3%)	0	100	100
19	S	73/100 (73%)	73 (100%)	0	0	100	100
20	T	48/94 (51%)	46 (96%)	2 (4%)	0	100	100
21	U	125/192 (65%)	121 (97%)	4 (3%)	0	100	100
22	V	162/193 (84%)	158 (98%)	4 (2%)	0	100	100
23	W	376/483 (78%)	371 (99%)	5 (1%)	0	100	100
24	X	202/496 (41%)	199 (98%)	3 (2%)	0	100	100
25	Y	96/102 (94%)	95 (99%)	0	1 (1%)	13	14
26	Z	79/153 (52%)	78 (99%)	1 (1%)	0	100	100
27	7	357/496 (72%)	340 (95%)	16 (4%)	1 (0%)	37	45
28	a	311/424 (73%)	305 (98%)	5 (2%)	1 (0%)	37	45
29	b	71/80 (89%)	63 (89%)	8 (11%)	0	100	100
30	c	24/128 (19%)	24 (100%)	0	0	100	100
31	d	76/110 (69%)	76 (100%)	0	0	100	100
32	e	244/383 (64%)	234 (96%)	10 (4%)	0	100	100
33	f	373/410 (91%)	364 (98%)	9 (2%)	0	100	100
34	h	308/384 (80%)	297 (96%)	11 (4%)	0	100	100
35	i	302/725 (42%)	295 (98%)	7 (2%)	0	100	100
36	k	20/153 (13%)	18 (90%)	2 (10%)	0	100	100
All	All	6351/9617 (66%)	6167 (97%)	177 (3%)	7 (0%)	50	59

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
28	a	210	ASP
2	B	481	ARG
8	H	225	ILE
25	Y	95	ARG
2	B	104	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/182 (97%)	177 (100%)	0	100	100
2	B	421/492 (86%)	420 (100%)	1 (0%)	92	96
3	C	312/343 (91%)	312 (100%)	0	100	100
4	D	370/444 (83%)	370 (100%)	0	100	100
5	E	90/126 (71%)	90 (100%)	0	100	100
6	F	124/124 (100%)	124 (100%)	0	100	100
7	G	112/113 (99%)	112 (100%)	0	100	100
8	H	174/320 (54%)	174 (100%)	0	100	100
9	I	104/197 (53%)	104 (100%)	0	100	100
10	J	109/256 (43%)	109 (100%)	0	100	100
11	K	106/107 (99%)	106 (100%)	0	100	100
12	L	92/128 (72%)	92 (100%)	0	100	100
13	M	90/137 (66%)	90 (100%)	0	100	100
14	N	107/362 (30%)	107 (100%)	0	100	100
15	O	93/117 (80%)	93 (100%)	0	100	100
16	P	78/95 (82%)	78 (100%)	0	100	100
17	Q	75/203 (37%)	75 (100%)	0	100	100
18	R	141/180 (78%)	141 (100%)	0	100	100
19	S	68/90 (76%)	68 (100%)	0	100	100
20	T	44/79 (56%)	44 (100%)	0	100	100
21	U	109/170 (64%)	108 (99%)	1 (1%)	75	86
22	V	147/173 (85%)	147 (100%)	0	100	100
23	W	328/410 (80%)	328 (100%)	0	100	100
24	X	174/395 (44%)	174 (100%)	0	100	100
25	Y	84/86 (98%)	84 (100%)	0	100	100
26	Z	66/119 (56%)	66 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
27	7	321/437 (74%)	321 (100%)	0	100	100
28	a	252/375 (67%)	252 (100%)	0	100	100
29	b	66/71 (93%)	66 (100%)	0	100	100
30	c	25/116 (22%)	25 (100%)	0	100	100
31	d	65/93 (70%)	65 (100%)	0	100	100
32	e	226/339 (67%)	226 (100%)	0	100	100
33	f	319/351 (91%)	319 (100%)	0	100	100
34	h	271/337 (80%)	271 (100%)	0	100	100
35	i	272/629 (43%)	272 (100%)	0	100	100
36	k	17/109 (16%)	17 (100%)	0	100	100
All	All	5629/8305 (68%)	5627 (100%)	2 (0%)	100	100

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	462	ASN
21	U	42	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
13	M	65	GLN
24	X	521	ASN
16	P	27	ASN
21	U	42	ASN
25	Y	93	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
37	2	1555/1560 (99%)	261 (16%)	10 (0%)

5 of 261 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
37	2	4	A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
37	2	5	U
37	2	12	A
37	2	17	G
37	2	30	G

5 of 10 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
37	2	1705	C
37	2	1706	C
37	2	1726	U
37	2	573	G
37	2	998	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
37	UR3	2	1807	37	19,22,23	2.81	8 (42%)	26,32,35	1.32	1 (3%)
37	4OC	2	1664	37	20,23,24	3.40	8 (40%)	26,32,35	0.90	1 (3%)
37	MA6	2	1827	37	18,26,27	1.17	3 (16%)	19,38,41	3.70	2 (10%)
37	G7M	2	525	37	20,26,27	2.19	6 (30%)	17,39,42	1.19	2 (11%)
37	2MG	2	1825	37	18,26,27	2.64	6 (33%)	16,38,41	1.75	4 (25%)
37	MA6	2	1828	37	18,26,27	1.11	2 (11%)	19,38,41	3.72	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
37	UR3	2	1807	37	-	2/7/25/26	0/2/2/2
37	4OC	2	1664	37	-	0/9/29/30	0/2/2/2
37	MA6	2	1827	37	-	5/7/29/30	0/3/3/3
37	G7M	2	525	37	-	3/3/25/26	0/3/3/3
37	2MG	2	1825	37	-	2/5/27/28	0/3/3/3
37	MA6	2	1828	37	-	4/7/29/30	0/3/3/3

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
37	2	1664	4OC	C4-N3	7.51	1.45	1.32
37	2	1825	2MG	C2-N2	7.36	1.49	1.33
37	2	1807	UR3	C2-N1	7.27	1.49	1.38
37	2	1664	4OC	C2-N3	6.60	1.49	1.36
37	2	1664	4OC	C6-C5	6.58	1.50	1.35

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
37	2	1828	MA6	N1-C6-N6	-15.12	101.15	117.06
37	2	1827	MA6	N1-C6-N6	-14.73	101.55	117.06
37	2	1827	MA6	N3-C2-N1	-5.66	119.83	128.68
37	2	1828	MA6	N3-C2-N1	-5.30	120.40	128.68
37	2	1807	UR3	C4-N3-C2	-4.96	119.90	124.56

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
37	2	1825	2MG	O4'-C4'-C5'-O5'
37	2	1827	MA6	O4'-C4'-C5'-O5'
37	2	1827	MA6	C3'-C4'-C5'-O5'
37	2	1827	MA6	C5-C6-N6-C10
37	2	1828	MA6	C5-C6-N6-C9

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
37	2	1664	4OC	1	0
37	2	1827	MA6	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
37	2	1825	2MG	1	0
37	2	1828	MA6	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 98 ligands modelled in this entry, 96 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
40	ATP	W	501	39	26,33,33	0.64	0	31,52,52	0.75	2 (6%)
42	GTP	7	602	39	26,34,34	1.13	2 (7%)	32,54,54	1.51	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
40	ATP	W	501	39	-	1/18/38/38	0/3/3/3
42	GTP	7	602	39	-	4/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
42	7	602	GTP	C5-C6	-3.99	1.39	1.47
42	7	602	GTP	C2-N3	2.07	1.38	1.33

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
42	7	602	GTP	C5-C6-N1	3.12	119.46	113.95
42	7	602	GTP	PB-O3B-PG	-3.09	122.24	132.83
42	7	602	GTP	C8-N7-C5	3.03	108.75	102.99
42	7	602	GTP	C3'-C2'-C1'	2.83	105.23	100.98
42	7	602	GTP	C2-N1-C6	-2.78	119.98	125.10

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
42	7	602	GTP	O4'-C4'-C5'-O5'
40	W	501	ATP	PA-O3A-PB-O1B
42	7	602	GTP	PB-O3A-PA-O2A
42	7	602	GTP	PG-O3B-PB-O2B
42	7	602	GTP	C5'-O5'-PA-O1A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
37	2	6

The worst 5 of 6 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	2	1142:A	O3'	1379:C	P	16.35
1	2	137:A	O3'	142:U	P	15.19
1	2	84:A	O3'	129:A	P	11.35
1	2	1842:U	O3'	1845:A	P	7.73
1	2	921:G	O3'	922:G	P	1.83


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-50014. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 294




Y Index: 294




Z Index: 294


6.2.2 Raw map



X Index: 294



Y Index: 294




Z Index: 294


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 300




Y Index: 290




Z Index: 280


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.0377. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_50014_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 5856 nm³; this corresponds to an approximate mass of 5290 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.441 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.441\AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.27	-	-
Author-provided FSC curve	2.27	2.79	2.34
Unmasked-calculated*	3.74	7.82	3.95

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.74 differs from the reported value 2.27 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-50014 and PDB model 9EVT. Per-residue inclusion information can be found in section 3 on page 14.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0377 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0377).

9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0377) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9010	 0.5020
2	 0.9740	 0.5800
7	 0.9100	 0.5250
A	 0.9270	 0.5410
B	 0.8560	 0.4390
C	 0.9460	 0.5680
D	 0.8790	 0.4990
E	 0.9420	 0.5380
F	 0.8380	 0.3910
G	 0.9690	 0.6320
H	 0.8250	 0.4610
I	 0.9630	 0.6190
J	 0.8640	 0.4510
K	 0.9590	 0.6220
L	 0.9010	 0.4900
M	 0.9680	 0.6220
N	 0.9220	 0.5590
O	 0.9820	 0.6530
P	 0.9430	 0.5960
Q	 0.8900	 0.5110
R	 0.6400	 0.2540
S	 0.8690	 0.4610
T	 0.9770	 0.6230
U	 0.6340	 0.2520
V	 0.8850	 0.4630
W	 0.8980	 0.4600
X	 0.8460	 0.4240
Y	 0.9150	 0.5370
Z	 0.9210	 0.4870
a	 0.8600	 0.4050
b	 0.5850	 0.1760
c	 0.8300	 0.4810
d	 0.9270	 0.5540
e	 0.9270	 0.5540
f	 0.8440	 0.3950



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
h	 0.6240	 0.1420
i	 0.5840	 0.1720
k	 0.8490	 0.3750