

Full wwPDB X-ray Structure Validation Report (i)

Oct 20, 2025 – 01:28 pm BST

PDB ID : 9EZW / pdb 00009ezw

Title : Structure of the single-domain antibody VHH_h3 in complex with human

Vsig4

Authors: Markovic, I.; Savvides, S.N.

Deposited on : 2024-04-14

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 2.0

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.46

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\#\text{Entries, resolution range}(\mathring{\mathbf{A}}))$
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	Λ	127	2%	
1	A	121	87% 4%	6% 7%
1	С	127	82%	11% 7%
1	E	127	9% 87%	12% •
	_		2%	12,0
1	G	127	83%	11% 6%
2	В	134	89%	• 7%



 $Continued\ from\ previous\ page...$

Mol	Chain	Length	Quality of chain		
	Б	104	4%		
2	D	134	86%	7%	7%
_	_		7%		
2	F'	134	87%	7%	5%
			3%		
2	Н	134	83%	10%	7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	F	201	-	-	X	-



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8368 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called V-set and immunoglobulin domain-containing protein 4.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace			
1	٨	118	Total	С	N	О	S	0	0	0
1	A	110	942	591	167	181	3	0	U	U
1	С	118	Total	С	N	N O S	1	0		
1		110	955	599	171	182	3	U	1	U
1	Е	126	Total	С	N	О	S	0	2	0
1	<u> 1</u> 2	120	1013	632	184	194	3	0		U
1	G	119	Total	С	N	О	S	0	3	0
1	G	119	980	613	177	187	3	0	3	U

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	119	ALA	-	expression tag	UNP Q9Y279
A	120	ALA	-	expression tag	UNP Q9Y279
A	121	ALA	-	expression tag	UNP Q9Y279
A	122	HIS	-	expression tag	UNP Q9Y279
A	123	HIS	-	expression tag	UNP Q9Y279
A	124	HIS	-	expression tag	UNP Q9Y279
A	125	HIS	-	expression tag	UNP Q9Y279
A	126	HIS	-	expression tag	UNP Q9Y279
A	127	HIS	-	expression tag	UNP Q9Y279
С	119	ALA	-	expression tag	UNP Q9Y279
С	120	ALA	-	expression tag	UNP Q9Y279
С	121	ALA	-	expression tag	UNP Q9Y279
С	122	HIS	-	expression tag	UNP Q9Y279
С	123	HIS	-	expression tag	UNP Q9Y279
С	124	HIS	-	expression tag	UNP Q9Y279
С	125	HIS	-	expression tag	UNP Q9Y279
С	126	HIS	-	expression tag	UNP Q9Y279
С	127	HIS	-	expression tag	UNP Q9Y279
Е	119	ALA	-	expression tag	UNP Q9Y279
Е	120	ALA	-	expression tag	UNP Q9Y279
Е	121	ALA	-	expression tag	UNP Q9Y279



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
Е	122	HIS	-	expression tag	UNP Q9Y279
Е	123	HIS	-	expression tag	UNP Q9Y279
E	124	HIS	-	expression tag	UNP Q9Y279
Е	125	HIS	-	expression tag	UNP Q9Y279
E	126	HIS	-	expression tag	UNP Q9Y279
Е	127	HIS	-	expression tag	UNP Q9Y279
G	119	ALA	-	expression tag	UNP Q9Y279
G	120	ALA	-	expression tag	UNP Q9Y279
G	121	ALA	-	expression tag	UNP Q9Y279
G	122	HIS	-	expression tag	UNP Q9Y279
G	123	HIS	-	expression tag	UNP Q9Y279
G	124	HIS	-	expression tag	UNP Q9Y279
G	125	HIS	-	expression tag	UNP Q9Y279
G	126	HIS	-	expression tag	UNP Q9Y279
G	127	HIS		expression tag	UNP Q9Y279

• Molecule 2 is a protein called single-domain antibody VHH_h3.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	125	Total	С	N	О	S	0	1	0
	Б	120	975	610	171	190	4	U	1	U
2	D	125	Total	С	N	О	S	0	0	0
	D	120	969	607	170	188	4	U	U	0
2	F	127	Total	С	N	О	S	0	1	0
2	Г	127	984	616	173	191	4	0	1	U
2	Н	195	Total	С	N	О	S	0	9	0
2	2 H	125	984	612	173	195	4	U	3	0

• Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0
3	Н	1	Total C O 4 2 2	0	0

 \bullet Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3)$ (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	1	Total C O 7 4 3	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	70	Total O 70 70	0	0
5	В	58	Total O 58 58	0	0
5	С	57	Total O 57 57	0	0
5	D	50	Total O 50 50	0	0
5	Е	54	Total O 54 54	0	0
5	F	75	Total O 75 75	0	0
5	G	70	Total O 70 70	0	0
5	Н	85	Total O 85 85	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: V-set and immunoglobulin domain-containing protein 4



• Molecule 2: single-domain antibody VHH_h3







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	60.87Å 129.94Å 144.28Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.97 - 1.90	Depositor
Resolution (A)	64.97 - 1.90	EDS
% Data completeness	98.3 (64.97-1.90)	Depositor
(in resolution range)	98.3 (64.97-1.90)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.11 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
D D.	0.190 , 0.228	Depositor
R, R_{free}	0.190 , 0.228	DCC
R_{free} test set	8949 reflections (9.84%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38, 40.2	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8368	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5
1	A	0.30	0/964	0.49	0/1315
1	С	0.28	0/978	0.48	0/1333
1	Е	0.25	0/1038	0.45	0/1417
1	G	0.27	0/1003	0.48	0/1367
2	В	0.32	0/998	0.47	0/1350
2	D	0.31	0/992	0.48	0/1341
2	F	0.33	0/1006	0.51	0/1359
2	Н	0.34	0/1006	0.53	0/1360
All	All	0.30	0/7985	0.49	0/10842

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	942	0	923	4	0
1	С	955	0	938	7	0
1	Е	1013	0	971	10	0
1	G	980	0	963	10	0
2	В	975	0	908	6	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	969	0	910	7	0
2	F	984	0	923	9	0
2	Н	984	0	916	9	0
3	A	8	0	12	1	0
3	С	4	0	6	0	0
3	Ε	8	0	12	0	0
3	F	4	0	6	4	0
3	G	12	0	18	1	0
3	Н	4	0	6	1	0
4	G	7	0	10	2	0
5	A	70	0	0	1	0
5	В	58	0	0	0	0
5	С	57	0	0	0	0
5	D	50	0	0	1	0
5	Ε	54	0	0	0	0
5	F	75	0	0	0	0
5	G	70	0	0	0	0
5	Н	85	0	0	0	0
All	All	8368	0	7522	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({ m \AA})$	$-$ overlap (\AA)
2:F:65:GLY:H	3:F:201:EDO:H22	1.45	0.81
2:B:109:TYR:OH	1:G:5[B]:GLU:OE2	2.00	0.79
2:B:0:GLN:N	2:B:0:GLN:OE1	2.18	0.77
1:A:102:ASP:OD2	1:A:104:ASN:ND2	2.22	0.73
2:D:2:GLN:HG2	2:F:116:GLN:O	1.93	0.69
1:E:100:THR:OG1	1:E:102[B]:ASP:OD1	2.11	0.68
2:F:82:MET:HE2	2:F:85:LEU:HD21	1.75	0.68
2:D:19:LEU:HG	2:D:82:MET:HE2	1.78	0.64
1:E:34:LEU:HD11	1:E:50:LEU:HD11	1.80	0.62
1:C:13:PRO:HG2	1:C:16:GLY:HA3	1.81	0.61
1:C:29:GLY:O	1:C:101:PRO:HD3	2.03	0.59
2:D:82:MET:HB3	2:D:85:LEU:HD21	1.83	0.59
1:E:18:VAL:HB	1:E:84:LEU:HD11	1.85	0.58
1:E:14:TRP:CD1	1:E:86:MET:HE2	2.38	0.57
4:G:204:PEG:O4	2:H:102:ARG:NH2	2.40	0.55



Continued from previous page...

Continuea from pre		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:86:MET:O	1:C:89:ARG:HG3	2.07	0.55
1:C:34:LEU:HD11	1:C:50:LEU:HD21	1.90	0.54
2:H:21:CYS:HB3	2:H:78:VAL:HG13	1.90	0.53
2:B:0:GLN:HE22	2:H:40:PRO:N	2.08	0.52
2:H:99:ARG:HE	2:H:112[B]:GLU:CD	2.17	0.52
1:E:115:ARG:HG2	1:E:115:ARG:HH11	1.75	0.51
3:A:202:EDO:H22	5:A:369:HOH:O	2.10	0.51
1:G:18:VAL:HB	1:G:84:LEU:HD11	1.94	0.50
1:G:36:LYS:HG2	1:G:50:LEU:HD23	1.94	0.49
2:F:11:VAL:HG11	2:F:85:LEU:HD13	1.96	0.48
1:G:100:THR:HB	1:G:101:PRO:HD2	1.96	0.48
1:E:79:LEU:HD11	1:E:81:LEU:HD13	1.97	0.47
2:D:88:GLU:HB3	5:D:239:HOH:O	2.16	0.46
1:E:67:LEU:C	1:E:67:LEU:HD23	2.41	0.46
2:B:82:MET:HB3	2:B:85:LEU:HD21	1.98	0.46
1:A:25:ASP:OD2	1:G:1[A]:ARG:N	2.48	0.45
2:H:2:GLN:HG2	2:H:4:VAL:HG23	1.98	0.45
1:C:14:TRP:CD2	1:C:15:LYS:HG2	2.52	0.45
1:C:18:VAL:HB	1:C:84:LEU:HD11	1.98	0.45
1:C:6:VAL:HB	1:C:110:LYS:HG2	1.98	0.45
1:G:36:LYS:HE2	1:G:95:GLU:OE2	2.17	0.44
2:F:7:GLY:O	2:F:17:LEU:HD22	2.18	0.44
1:E:2:PRO:HG3	1:E:27:LEU:HD22	2.00	0.44
2:H:25:GLY:O	2:H:26:ARG:NH1	2.42	0.43
2:F:65:GLY:N	3:F:201:EDO:H22	2.22	0.43
1:G:34:LEU:HD11	1:G:50:LEU:HD21	2.01	0.43
1:G:34:LEU:HD23	4:G:204:PEG:H12	2.01	0.43
1:A:67:LEU:HG	1:A:79:LEU:HD11	2.01	0.43
2:F:65:GLY:H	3:F:201:EDO:C2	2.26	0.43
2:B:67:PHE:CE2	2:B:82:MET:HE3	2.55	0.42
2:F:21:CYS:HB3	2:F:78:VAL:CG1	2.50	0.42
1:E:115:ARG:HG2	1:E:115:ARG:NH1	2.33	0.42
2:H:33:MET:HE3	2:H:33:MET:HB3	1.88	0.42
1:A:18:VAL:HB	1:A:84:LEU:HD11	2.01	0.42
2:B:11:VAL:HG11	2:B:85:LEU:HD12	2.02	0.41
1:G:94:CYS:O	1:G:109:ASP:HA	2.21	0.41
2:D:108:GLN:HG2	2:D:114:TRP:CH2	2.55	0.41
1:E:37:TRP:CE2	1:E:79:LEU:HB2	2.56	0.41
2:H:28:PHE:O	2:H:71:ARG:NH2	2.54	0.41
2:H:113:TYR:CZ	3:H:201:EDO:H22	2.56	0.41
2:D:7:GLY:O	2:D:17:LEU:HD11	2.20	0.40



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:F:59:TYR:CE2	3:F:201:EDO:H12	2.57	0.40
2:D:17:LEU:HD12	2:D:18:ARG:H	1.85	0.40
1:G:33:VAL:O	3:G:201:EDO:H22	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	$116/127\ (91\%)$	111 (96%)	5 (4%)	0	100	100
1	C	117/127~(92%)	113 (97%)	4 (3%)	0	100	100
1	E	$126/127\ (99\%)$	123 (98%)	3 (2%)	0	100	100
1	G	$119/127\ (94\%)$	113 (95%)	6 (5%)	0	100	100
2	В	124/134~(92%)	121 (98%)	3 (2%)	0	100	100
2	D	123/134~(92%)	119 (97%)	4 (3%)	0	100	100
2	F	124/134~(92%)	121 (98%)	3 (2%)	0	100	100
2	Н	$126/134\ (94\%)$	123 (98%)	3 (2%)	0	100	100
All	All	975/1044~(93%)	944 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	108/115 (94%)	108 (100%)	0	100	100
1	С	110/115 (96%)	110 (100%)	0	100	100
1	E	113/115 (98%)	113 (100%)	0	100	100
1	G	112/115 (97%)	111 (99%)	1 (1%)	75	77
2	В	97/103 (94%)	97 (100%)	0	100	100
2	D	97/103 (94%)	97 (100%)	0	100	100
2	F	97/103 (94%)	97 (100%)	0	100	100
2	Н	99/103 (96%)	99 (100%)	0	100	100
All	All	833/872 (96%)	832 (100%)	1 (0%)	92	93

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	79	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	HIS
1	A	91	HIS
1	С	71	HIS
1	С	91	HIS
2	D	2	GLN
2	D	81	GLN
2	D	83	ASN
2	F	119	GLN
1	G	91	HIS
1	G	117	GLN
2	Н	81	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Dag	Link	В	ond leng	$_{ m gths}$	В	ond ang	gles
IVIOI	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	EDO	A	201	-	3,3,3	0.45	0	2,2,2	0.42	0
3	EDO	С	201	-	3,3,3	0.47	0	2,2,2	0.41	0
3	EDO	Е	201	-	3,3,3	0.47	0	2,2,2	0.41	0
3	EDO	G	201	-	3,3,3	0.47	0	2,2,2	0.49	0
3	EDO	F	201	-	3,3,3	0.35	0	2,2,2	0.51	0
3	EDO	Н	201	-	3,3,3	0.45	0	2,2,2	0.35	0
3	EDO	G	202	-	3,3,3	0.54	0	2,2,2	0.16	0
4	PEG	G	204	-	6,6,6	0.10	0	5,5,5	0.18	0
3	EDO	G	203	-	3,3,3	0.44	0	2,2,2	0.42	0
3	EDO	Е	202	-	3,3,3	0.58	0	2,2,2	0.18	0
3	EDO	A	202	-	3,3,3	0.43	0	2,2,2	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	201	-	-	0/1/1/1	-
3	EDO	С	201	-	-	0/1/1/1	-
3	EDO	E	201	-	-	0/1/1/1	-
3	EDO	G	201	_	-	0/1/1/1	-
3	EDO	F	201	-	-	1/1/1/1	-
3	EDO	Н	201	_	-	0/1/1/1	-
3	EDO	G	202	-	-	1/1/1/1	-
4	PEG	G	204	-	-	1/4/4/4	-
3	EDO	G	203	-	-	0/1/1/1	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	Е	202	-	-	0/1/1/1	-
3	EDO	A	202	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	201	EDO	O1-C1-C2-O2
3	G	202	EDO	O1-C1-C2-O2
4	G	204	PEG	O1-C1-C2-O2


There are no ring outliers.


5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	201	EDO	1	0
3	F	201	EDO	4	0
3	Н	201	EDO	1	0
4	G	204	PEG	2	0
3	A	202	EDO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	$118/127\ (92\%)$	0.01	2 (1%) 69	71	22, 32, 51, 63	0
1	С	$118/127\ (92\%)$	0.26	5 (4%) 41	43	18, 36, 58, 62	1 (0%)
1	E	$126/127\ (99\%)$	0.55	12 (9%) 15	16	21, 38, 62, 74	2 (1%)
1	G	119/127 (93%)	0.03	2 (1%) 69	71	15, 33, 50, 74	3 (2%)
2	В	$125/134\ (93\%)$	0.16	2 (1%) 70	72	18, 33, 53, 71	1 (0%)
2	D	$125/134\ (93\%)$	0.34	6 (4%) 36	37	22, 35, 54, 80	0
2	F	127/134~(94%)	0.11	9 (7%) 23	24	19, 31, 51, 79	2 (1%)
2	Н	125/134~(93%)	-0.11	4 (3%) 50	52	14, 29, 43, 68	4 (3%)
All	All	983/1044 (94%)	0.17	42 (4%) 40	42	14, 33, 56, 80	13 (1%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	126	HIS	5.1
2	F	127	ALA	4.8
1	G	119	ALA	4.4
1	Е	120	ALA	4.4
1	Е	119	ALA	4.2
2	F	0	GLN	4.0
2	D	124	SER	3.9
2	F	25	GLY	3.8
2	D	0	GLN	3.5
1	Е	121	ALA	3.4
1	Е	14	TRP	3.3
1	Е	125	HIS	3.3
2	Н	25	GLY	3.1
2	F	1	VAL	3.0
2	Н	26	ARG	2.9
2	В	124	SER	2.6



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	126	ALA	2.6
1	Е	124	HIS	2.6
2	D	8	GLY	2.6
2	D	40	PRO	2.4
2	Н	28	PHE	2.4
1	С	14	TRP	2.3
1	Е	61	ALA	2.3
2	В	25	GLY	2.3
2	F	27	THR	2.2
1	G	61	ALA	2.2
1	С	54	SER	2.2
2	F	109	TYR	2.2
1	Е	83	THR	2.1
1	Е	104	ASN	2.1
1	Е	122	HIS	2.1
2	Н	27	THR	2.1
2	D	1	VAL	2.1
1	Е	123	HIS	2.1
1	A	61	ALA	2.1
2	D	112	GLU	2.1
1	A	118	LYS	2.1
1	С	65	GLY	2.0
1	С	118	LYS	2.0
1	С	61	ALA	2.0
2	F	2	GLN	2.0
2	F	40	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	EDO	A	202	4/4	0.75	0.15	48,49,50,59	0
4	PEG	G	204	7/7	0.80	0.15	41,45,57,58	0
3	EDO	Е	202	4/4	0.83	0.17	39,40,42,47	0
3	EDO	Н	201	4/4	0.87	0.14	41,45,47,48	0
3	EDO	G	202	4/4	0.87	0.13	44,47,52,54	0
3	EDO	G	201	4/4	0.88	0.11	40,42,45,47	0
3	EDO	G	203	4/4	0.89	0.12	48,49,53,54	0
3	EDO	F	201	4/4	0.90	0.11	31,41,42,43	0
3	EDO	Е	201	4/4	0.91	0.10	49,53,56,65	0
3	EDO	A	201	4/4	0.92	0.12	38,41,42,45	0
3	EDO	С	201	4/4	0.93	0.14	46,48,51,53	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around PEG G 204:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO E 202:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around EDO H 201:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO G 202:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around EDO G 201:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO G 203:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around EDO F 201:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO E 201:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around EDO A 201:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

