



wwPDB EM Validation Summary Report ⓘ

Feb 20, 2025 – 12:05 pm GMT

PDB ID : 9F0J
EMDB ID : EMD-50111
Title : Cryo-EM structure of the I923V MDA5-dsRNA filament without nucleotide
Authors : Singh, R.; Herrero del Valle, A.; Modis, Y.
Deposited on : 2024-04-16
Resolution : 3.35 Å (reported)
Based on initial model : 7BKQ

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **FAILED**
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Clashscore | 210492 | 15764 |
| Ramachandran outliers | 207382 | 16835 |
| Sidechain outliers | 206894 | 16415 |
| RNA backbone | 6643 | 2191 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 1028 | |
| 2 | X | 15 | |
| 3 | Z | 15 | |

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11368 atoms, of which 5586 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon-induced helicase C domain-containing protein 1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|------|------|-----|-----|---------|-------|---|
| | | | Total | C | H | N | O | | | S |
| 1 | A | 640 | 10403 | 3273 | 5261 | 888 | 948 | 33 | 0 | 0 |

There are 42 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|-----------------------|------------|
| A | -20 | MET | - | initiating methionine | UNP Q8R5F7 |
| A | -19 | GLY | - | expression tag | UNP Q8R5F7 |
| A | -18 | SER | - | expression tag | UNP Q8R5F7 |
| A | -17 | SER | - | expression tag | UNP Q8R5F7 |
| A | -16 | HIS | - | expression tag | UNP Q8R5F7 |
| A | -15 | HIS | - | expression tag | UNP Q8R5F7 |
| A | -14 | HIS | - | expression tag | UNP Q8R5F7 |
| A | -13 | HIS | - | expression tag | UNP Q8R5F7 |
| A | -12 | HIS | - | expression tag | UNP Q8R5F7 |
| A | -11 | HIS | - | expression tag | UNP Q8R5F7 |
| A | -10 | SER | - | expression tag | UNP Q8R5F7 |
| A | -9 | SER | - | expression tag | UNP Q8R5F7 |
| A | -8 | GLY | - | expression tag | UNP Q8R5F7 |
| A | -7 | ARG | - | expression tag | UNP Q8R5F7 |
| A | -6 | GLU | - | expression tag | UNP Q8R5F7 |
| A | -5 | ASN | - | expression tag | UNP Q8R5F7 |
| A | -4 | LEU | - | expression tag | UNP Q8R5F7 |
| A | -3 | TYR | - | expression tag | UNP Q8R5F7 |
| A | -2 | PHE | - | expression tag | UNP Q8R5F7 |
| A | -1 | GLN | - | expression tag | UNP Q8R5F7 |
| A | 0 | GLY | - | expression tag | UNP Q8R5F7 |
| A | 1 | HIS | - | expression tag | UNP Q8R5F7 |
| A | 2 | MET | - | expression tag | UNP Q8R5F7 |
| A | ? | - | ASP | deletion | UNP Q8R5F7 |
| A | ? | - | LYS | deletion | UNP Q8R5F7 |
| A | ? | - | SER | deletion | UNP Q8R5F7 |
| A | ? | - | ASP | deletion | UNP Q8R5F7 |
| A | ? | - | ASP | deletion | UNP Q8R5F7 |

Continued on next page...

Continued from previous page...

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| A | ? | - | GLU | deletion | UNP Q8R5F7 |
| A | ? | - | ALA | deletion | UNP Q8R5F7 |
| A | ? | - | SER | deletion | UNP Q8R5F7 |
| A | ? | - | SER | deletion | UNP Q8R5F7 |
| A | ? | - | CYS | deletion | UNP Q8R5F7 |
| A | ? | - | ASN | deletion | UNP Q8R5F7 |
| A | ? | - | ASP | deletion | UNP Q8R5F7 |
| A | ? | - | GLN | deletion | UNP Q8R5F7 |
| A | ? | - | LEU | deletion | UNP Q8R5F7 |
| A | ? | - | LYS | deletion | UNP Q8R5F7 |
| A | ? | - | GLY | deletion | UNP Q8R5F7 |
| A | ? | - | ASP | deletion | UNP Q8R5F7 |
| A | ? | - | VAL | deletion | UNP Q8R5F7 |
| A | 923 | VAL | ILE | engineered mutation | UNP Q8R5F7 |

- Molecule 2 is a RNA chain called RNA (5'-UCCAUGC GCAUGACG-3').

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|-----|-----|----|-----|---------|-------|---|
| | | | Total | C | H | N | O | | | P |
| 2 | X | 15 | 480 | 142 | 162 | 56 | 105 | 15 | 0 | 0 |

- Molecule 3 is a RNA chain called RNA (5'-CGUCAUGC GCAUGGA-3').

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|-----|-----|----|-----|---------|-------|---|
| | | | Total | C | H | N | O | | | P |
| 3 | Z | 15 | 484 | 143 | 163 | 58 | 105 | 15 | 0 | 0 |

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

| Mol | Chain | Residues | Atoms | | AltConf |
|-----|-------|----------|-------|----|---------|
| | | | Total | Zn | |
| 4 | A | 1 | 1 | 1 | 0 |



- Molecule 3: RNA (5'-CGUCAUGCGCAUGGA-3')

Chain Z:  100%

There are no outlier residues recorded for this chain.

4 Experimental information

| Property | Value | Source |
|------------------------------------|---|-----------|
| EM reconstruction method | HELICAL | Depositor |
| Imposed symmetry | HELICAL, twist=91.63°, rise=44.84 Å, axial sym=C1 | Depositor |
| Number of segments used | 547562 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{Å}^2$) | 44 | Depositor |
| Minimum defocus (nm) | 500 | Depositor |
| Maximum defocus (nm) | 2500 | Depositor |
| Magnification | 105000 | Depositor |
| Image detector | GATAN K3 BIOQUANTUM (6k x 4k) | Depositor |

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|---------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 0.28 | 0/5222 | 0.46 | 0/7011 |
| 2 | X | 0.39 | 0/354 | 0.76 | 0/549 |
| 3 | Z | 0.41 | 0/358 | 0.80 | 0/556 |
| All | All | 0.30 | 0/5934 | 0.52 | 0/8116 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 5142 | 5261 | 5261 | 89 | 0 |
| 2 | X | 318 | 162 | 163 | 1 | 0 |
| 3 | Z | 321 | 163 | 163 | 0 | 0 |
| 4 | A | 1 | 0 | 0 | 0 | 0 |
| All | All | 5782 | 5586 | 5587 | 89 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 89 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 1:A:779:ARG:NH2 | 1:A:797:ASP:OD1 | 2.08 | 0.87 |
| 1:A:960:ILE:HD11 | 1:A:971:MET:SD | 2.29 | 0.72 |
| 1:A:703:LYS:O | 1:A:707:THR:HG23 | 1.91 | 0.71 |
| 1:A:511:CYS:SG | 1:A:519:ILE:HD11 | 2.31 | 0.70 |
| 1:A:472:LEU:O | 1:A:476:ASN:N | 2.26 | 0.68 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|-----|-------|----------------|-----------|---------|----------|-----------------------|
| 1 | A | 622/1028 (60%) | 590 (95%) | 32 (5%) | 0 | 100 100 |

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|-----|-------|---------------|------------|----------|-----------------------|
| 1 | A | 572/918 (62%) | 572 (100%) | 0 | 100 100 |

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 469 | ASN |
| 1 | A | 476 | ASN |
| 1 | A | 578 | HIS |

5.3.3 RNA [i](#)

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|-------------|-------------------|-----------------|
| 2 | X | 14/15 (93%) | 0 | 0 |
| 3 | Z | 14/15 (93%) | 0 | 0 |
| All | All | 28/30 (93%) | 0 | 0 |

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.