

wwPDB EM Validation Summary Report (i)

Nov 19, 2022 – 11:54 pm GMT

PDB ID : 6F38

EMDB ID : EMD-4177

Title : Cryo-EM structure of two dynein tail domains bound to dynactin and HOOK3 Authors : Lau, C.K.; Urnavicius, L.; Elshenawy, M.M.; Morales-Rios, E.; Motz, C.;

Yildiz, A.; Carter, A.P.

Deposited on : 2017-11-28

Resolution : 6.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.4, CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 6.70 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} ext{Whole archive} \ (\# ext{Entries}) \end{array}$	${ m EM~structures} \ (\#{ m Entries})$
Ramachandran outliers	154571	4023

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	376	98%	
1	В	376	98%	
1	С	376	98%	••
1	D	376	98%	
1	Е	376	98%	••
1	F	376	98%	•
1	G	376	97%	
1	I	376	98%	
2	Н	375	99%	
3	J	390	94%	• 5%
4	K	286	97%	•





Continued from previous page...

Mol	Chain	Length	Quality of chain	
5	L	272	99%	
6	M	587	100%	
7	N	616	100%	
8	0	65	100%	
8	Р	65	100% 54%	
9	Q	87	100%	
9	R	87	9%	
10	U	190	89%	• 9%
11	V	182	90%	• 9%
12	X	223	26%	
12	X	223	25%	
13	Y	264	100%	
14	Z	52	100%	
15	a	68	76%	24%
16	b	90	<u>.</u>	
			82% 6%	• 17%
17	С	50	70% 7%	• 26%
18	d	29	83% 19%	17%
19	е	1455	63% 9%	36%
19	f	1455	64%	36%
19	m	1455	63%	36%
19	n	1455	63%	36%
20	g	612	65%	35%
20	h	612	6 5% •	35%
20	О	612	9%	35%
			30%	
20	p	612	65%	34%

Continued on next page...



Continued from previous page...

Mol	Chain		Quality of	chain
			54%	
21	i	492	54%	46%
			28%	
21	j	492	55%	44%
			48%	
21	q	492	54%	46%
			54%	
21	r	492	54%	46%
20	,	0.0	14%	
22	k	96	97%	•
00	1	O.C.	28%	
22	1	96	97%	•
00		O.C.	48%	
22	S	96	39%	• •
22	+	96		
	t	90	97%	•
23	7	53		
23	Z	99	100%	



2 Entry composition (i)

There are 25 unique types of molecules in this entry. The entry contains 68864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ARP1 actin related protein 1 homolog A.

Mol	Chain	Residues		Ator	ns		AltConf	Trace	
1	A	370	Total	С	N	О	0	0	
1	A	370	1821	1082	369	370	0	0	
1	В	370	Total	С	N	О	0	0	
1	Б	370	1822	1082	370	370	0	0	
1	С	370	Total	С	N	О	0	0	
1		370	1822	1082	370	370	0	0	
1	D	370	Total	С	N	О	0	0	
1	ט	D 370	1822	1082	370	370	U	0	
1	E	E	370	Total	С	N	О	0	0
1	<u> 1</u> 2	370	1822	1082	370	370	U	0	
1	F	370	Total	С	N	О	0	0	
1	I'	370	1822	1082	370	370	0	0	
1	С	370	Total	С	N	О	0	0	
1	1 G	370	1822	1082	370	370	U	0	
1	ı T	I 370	Total	С	N	О	0	0	
1	1	310	1822	1082	370	370			

• Molecule 2 is a protein called Actin, cytoplasmic 1.

Mol	Chain	Residues		Ator	$\mathbf{n}\mathbf{s}$	AltConf	Trace	
2	Н	370	Total 1822	C 1082	N 370	O 370	0	0

• Molecule 3 is a protein called Actin related protein 10 homolog.

Mo	ol	Chain	Residues		Ator	$\mathbf{n}\mathbf{s}$	AltConf	Trace	
3		J	369	Total 1819	C 1081	N 369	O 369	0	0

• Molecule 4 is a protein called Capping protein (Actin filament) muscle Z-line, alpha 1.



Mol	Chain	Residues	Atoms				AltConf	Trace
4	K	278	Total 1378	C 822	N 278	O 278	0	0

• Molecule 5 is a protein called F-actin capping protein beta subunit.

Mol	Chain	Residues	Atoms				AltConf	Trace
5	L	269	Total	С	N	О	0	0
		230	1327	789	269	269		

• Molecule 6 is a protein called Dynactin Subunit 2.

Mol	Chain	Residues	Atoms				AltConf	Trace
6	M	587	Total 2935	C 1761	N 587	O 587	0	0

• Molecule 7 is a protein called Dynactin Subunit 2.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	N	616	Total	С	N	O	0	0
1	11	010	3080	1848	616	616	U	0

• Molecule 8 is a protein called Dynactin Subunit 3.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	О	65	Total 323	C 193			0	0
8	Р	65	Total 323	C 193			0	0

• Molecule 9 is a protein called Dynactin Subunit 2.

Mol	Chain	Residues	Atoms				AltConf	Trace
9	Q	87	Total 435				0	0
9	R	87	Total 435	C 261			0	0

• Molecule 10 is a protein called Dynactin 6.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
10	TT	172	Total	С	N	О	0	1
10	U	172	843	500	171	172	0	1



• Molecule 11 is a protein called Dynactin subunit 5.

Mol	Chain	Residues	Atoms				AltConf	Trace
11	V	165	Total 812	C 482	N 165	O 165	0	0

• Molecule 12 is a protein called HOOK3.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
19	V	223	Total	С	N	О	0	0
	220	1115	669	223	223	0		
19	37	222	Total	С	N	О	0	0
12 x	222	1110	666	222	222	0	U	

 \bullet Molecule 13 is a protein called Dynactin Subunit 4.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
13	Y	264	Total 1320	C 792	N 264	O 264	0	0

• Molecule 14 is a protein called Dynactin Subunit 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
14	Z	52	Total 260	C 156	N 52	O 52	0	0

• Molecule 15 is a protein called Dynactin subunit 2.

Mol	Chain	Residues	Atoms				AltConf	Trace
15	a	52	Total 259	C 155	N 52	O 52	0	0

• Molecule 16 is a protein called Dynactin subunit 2.

Mol	Chain	Residues	Atoms				AltConf	Trace
16	b	75	Total 369	C 219	N 75	O 75	0	0

• Molecule 17 is a protein called Dynactin subunit 2.

Mol	Chain	Residues		Aton	ıs	AltConf	Trace	
17	c	37	Total 184	C 110	N 37	O 37	0	0



• Molecule 18 is a protein called Dynactin subunit 2.

Mol	Chain	Residues	Atoms				AltConf	Trace
10	J	24	Total	С	N	О	0	0
10	a	24	119	71	24	24	0	U

• Molecule 19 is a protein called Cytoplasmic dynein 1 heavy chain 1.

Mol	Chain	Residues		Ator	ns	AltConf	Trace	
19	е	926	Total	С	N	О	0	0
19	е	920	4598	2746	926	926	U	0
19	f	929	Total	С	N	О	0	0
19	1	929	4613	2755	929	929	0	0
19	m	926	Total	С	N	О	0	0
19	m	920	4598	2746	926	926	0	0
19	10	928	Total	С	N	О	0	0
19	n	920	4608	2752	928	928		

• Molecule 20 is a protein called Cytoplasmic dynein 1 intermediate chain 2.

Mol	Chain	Residues		Ator	AltConf	Trace		
20	ď	397	Total	С	N	О	0	0
20	g	391	1961	1167	397	397	U	0
20	h	400	Total	$^{\mathrm{C}}$	N	O	0	0
20	20 11	400	1975	1175	400	400	0	U
20	0	397	Total	\mathbf{C}	N	O	0	0
20	О	391	1961	1167	397	397	0	0
20	n	401	Total	С	N	О	0	1
20	p	401	1977	1176	401	400	U	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
g	484	SER	THR	conflict	UNP Q13409
g	499	GLY	ASP	conflict	UNP Q13409
h	484	SER	THR	conflict	UNP Q13409
h	499	GLY	ASP	conflict	UNP Q13409
О	484	SER	THR	conflict	UNP Q13409
О	499	GLY	ASP	conflict	UNP Q13409
р	484	SER	THR	conflict	UNP Q13409
p	499	GLY	ASP	conflict	UNP Q13409

• Molecule 21 is a protein called Cytoplasmic dynein 1 light intermediate chain 2.



Mol	Chain	Residues		Ato	ms		AltConf	Trace	
21	i	268	Total	С	N	О	0	0	
21	1		1327	791	268	268	0		
21	i	275	Total	С	N	O	0	0	
21	J	210	1362	812	275	275	U		
21	a	268	Total	С	N	O	0	0	
21	q	200	1327	791	268	268			
21	r	268	Total	С	N	О	0	0	
21	r	∠08	1327	791	268	268			

• Molecule 22 is a protein called Dynein light chain roadblock-type 1.


Mol	Chain	Residues		Aton	ıs		AltConf	Trace	
22	k	93	Total	С	N	О	0	0	
22	K	90	462	276	93	93	U		
22	1	93	Total	С	N	O	0	0	
22	1	90	462	276	93	93		U	
22	g	93	Total	С	N	Ο	0	0	
22 s	90	462	276	93	93	0	U		
22	+	93	Total	С	N	О	0	0	
22	t	93	462	276	93	93		U	

• Molecule 23 is a protein called Dynactin Subunit 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
23	Z	53	Total 265	C 159	N 53	O 53	0	0

• Molecule 24 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).






Mol	Chain	Residues		Ato	oms			AltConf	
24	A	1	Total	С	N	О	Р	0	
24	A	1	27	10	5	10	2	U	
24	В	1	Total	С	N	О	Р	0	
24	Б	1	27	10	5	10	2	U	
24	С	1	Total	С	N	О	Р	0	
24		1	27	10	5	10	2	U	
24	D	1	Total	С	N	О	Р	0	
24	ע	1	27	10	5	10	2	U	
24	Е	1	Total	С	N	О	Р	0	
24	Ľ	1	27	10	5	10	2	U	
24	F	1	Total	С	N	Ο	Р	0	
24	I.	1	27	10	5	10	2	U	
24	G	1	Total	\mathbf{C}	N	Ο	Р	0	
24	G	1	27	10	5	10	2	O	
24	I	1	Total	С	N	Ο	Р	0	
24	1	1	27	10	5	10	2	0	
24	J	1	Total	С	N	Ο	Р	0	
24	J	1	27	10	5	10	2		

 \bullet Molecule 25 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3).$






Mol	Chain	Residues	Atoms					AltConf
25	п	1	Total	С	N	О	Р	0
25	п	1	31	10	5	13	3	U




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ARP1 actin related protein 1 homolog A

















• Molecule 16: Dynactin subunit 2











• Molecule 19: Cytoplasmic dynein 1 heavy chain 1









LYS ASP VAL LEU LEU VAL ALA GLN GLN


• Molecule 19: Cytoplasmic dynein 1 heavy chain 1








• Molecule 19: Cytoplasmic dynein 1 heavy chain 1








• Molecule 20: Cytoplasmic dynein 1 intermediate chain 2

Chain h: 65% . 35%










• Molecule 21: Cytoplasmic dynein 1 light intermediate chain 2












 \bullet Molecule 22: Dynein light chain roadblock-type 1

Chain s: 96% ...





• Molecule 22: Dynein light chain roadblock-type 1

39% Chain t: 97%



• Molecule 23: Dynactin Subunit 1

Chain z: 100%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	23407	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	45	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.237	Depositor
Minimum map value	-0.074	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.06	Depositor
Map size (Å)	852.0, 852.0, 852.0	wwPDB
Map dimensions	600, 600, 600	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.42, 1.42, 1.42	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Clasia.	Во	nd lengths	В	ond angles
Mol	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.43	0/1820	0.76	0/2529
1	В	0.47	0/1821	0.79	0/2531
1	С	0.46	0/1821	0.81	$1/2531 \ (0.0\%)$
1	D	0.49	0/1821	0.76	0/2531
1	Е	0.48	0/1821	0.81	0/2531
1	F	0.49	0/1821	0.81	$2/2531 \ (0.1\%)$
1	G	0.53	1/1821 (0.1%)	0.84	3/2531 (0.1%)
1	I	0.45	0/1821	0.76	$1/2531 \ (0.0\%)$
2	Н	0.46	0/1821	0.75	0/2531
3	J	0.44	0/1816	0.82	$2/2522 \ (0.1\%)$
4	K	0.38	0/1377	0.80	0/1919
5	L	0.43	0/1326	0.73	0/1844
10	U	0.38	0/841	0.65	0/1168
11	V	0.29	0/811	0.61	0/1126
15	a	0.43	0/257	0.84	0/356
16	b	0.51	1/367~(0.3%)	0.87	0/507
17	С	0.40	0/182	0.90	1/251~(0.4%)
18	d	0.35	0/118	0.73	0/163
19	е	0.29	0/4593	0.52	3/6403~(0.0%)
19	f	0.33	0/4608	0.68	3/6424 (0.0%)
19	m	0.33	0/4593	0.62	3/6403~(0.0%)
19	n	0.30	0/4603	0.56	3/6417~(0.0%)
20	g	0.29	0/1959	0.70	0/2725
20	h	0.34	0/1972	0.82	3/2741 (0.1%)
20	О	0.35	0/1959	0.82	$2/2725 \ (0.1\%)$
20	р	0.29	0/1975	0.66	0/2747
21	i	0.26	0/1322	0.60	0/1835
21	j	0.31	0/1357	0.75	0/1884
21	q	0.26	0/1322	0.52	0/1835
21	r	0.27	0/1322	0.56	0/1835
22	k	0.28	0/461	0.59	0/642
22	1	0.31	0/461	0.63	0/642



Mol	Chain	Bo	nd lengths	Bond angles		
MIOI	Moi Chain		# Z > 5	RMSZ	# Z >5	
22	s	0.36	0/461	0.79	0/642	
22	t	0.32	0/461	0.76	0/642	
All	All	0.38	$2/56912 \ (0.0\%)$	0.70	27/79175 (0.0%)	

All (2) bond length outliers are listed below:

	Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
	1	G	65	HIS	CA-CB	-8.23	1.35	1.53
Ī	16	b	20	TYR	CA-CB	-5.29	1.42	1.53

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
20	О	412	SER	C-N-CA	12.66	153.34	121.70
3	J	247	ASP	C-N-CA	8.94	144.06	121.70
19	m	1030	PRO	N-CA-CB	7.99	112.88	103.30
1	F	112	GLU	C-N-CA	7.70	140.96	121.70
19	n	1030	PRO	N-CA-CB	7.68	112.51	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	n Analysed Favoured Allowed		Outliers	Percentiles	
1	A	368/376~(98%)	311 (84%)	56 (15%)	1 (0%)	41 76
1	В	368/376~(98%)	318 (86%)	49 (13%)	1 (0%)	41 76

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	С	368/376~(98%)	315 (86%)	52 (14%)	1 (0%)	41	76
1	D	368/376~(98%)	316 (86%)	52 (14%)	0	100	100
1	E	368/376~(98%)	315 (86%)	51 (14%)	2 (0%)	29	69
1	F	$368/376\ (98\%)$	310 (84%)	57 (16%)	1 (0%)	41	76
1	G	$368/376\ (98\%)$	308 (84%)	58 (16%)	2 (0%)	29	69
1	I	$368/376\ (98\%)$	308 (84%)	59 (16%)	1 (0%)	41	76
2	Н	$368/375\ (98\%)$	316 (86%)	52 (14%)	0	100	100
3	J	363/390~(93%)	296 (82%)	67 (18%)	0	100	100
4	K	276/286 (96%)	240 (87%)	35 (13%)	1 (0%)	34	72
5	L	267/272 (98%)	232 (87%)	35 (13%)	0	100	100
10	U	169/190 (89%)	126 (75%)	41 (24%)	2 (1%)	13	50
11	V	163/182 (90%)	115 (71%)	46 (28%)	2 (1%)	13	50
15	a	48/68 (71%)	33 (69%)	15 (31%)	0	100	100
16	b	71/90 (79%)	51 (72%)	20 (28%)	0	100	100
17	c	33/50 (66%)	25 (76%)	7 (21%)	1 (3%)	4	28
18	d	22/29~(76%)	17 (77%)	5 (23%)	0	100	100
19	e	916/1455~(63%)	770 (84%)	140 (15%)	6 (1%)	22	63
19	f	919/1455~(63%)	786 (86%)	130 (14%)	3 (0%)	41	76
19	m	$916/1455\ (63\%)$	787 (86%)	125 (14%)	4 (0%)	34	72
19	n	918/1455~(63%)	798 (87%)	114 (12%)	6 (1%)	22	63
20	g	393/612 (64%)	296 (75%)	96 (24%)	1 (0%)	41	76
20	h	394/612~(64%)	305 (77%)	88 (22%)	1 (0%)	41	76
20	О	393/612 (64%)	301 (77%)	91 (23%)	1 (0%)	41	76
20	р	397/612~(65%)	274 (69%)	122 (31%)	1 (0%)	41	76
21	i	258/492~(52%)	225 (87%)	33 (13%)	0	100	100
21	j	265/492~(54%)	206 (78%)	57 (22%)	2 (1%)	19	60
21	q	258/492~(52%)	202 (78%)	56 (22%)	0	100	100
21	r	258/492~(52%)	204 (79%)	54 (21%)	0	100	100
22	k	91/96~(95%)	64 (70%)	27 (30%)	0	100	100
22	1	91/96~(95%)	66 (72%)	25 (28%)	0	100	100
22	s	91/96~(95%)	69 (76%)	21 (23%)	1 (1%)	14	52

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
22	t	91/96~(95%)	75 (82%)	16 (18%)	0	100 100
All	All	11373/15560 (73%)	9380 (82%)	1952 (17%)	41 (0%)	38 72

5 of 41 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
10	U	79	PRO
19	е	1026	MET
19	е	1027	PRO
19	е	1030	PRO
19	f	1030	PRO

5.3.2 Protein sidechains (i)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Trino	Chain	n Res	Link	Во	ond leng	ths	В	ond ang	les
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
24	ADP	С	800	-	24,29,29	0.84	0	29,45,45	1.71	5 (17%)
24	ADP	G	800	-	24,29,29	0.97	1 (4%)	29,45,45	1.40	4 (13%)
24	ADP	I	800	-	24,29,29	0.95	1 (4%)	29,45,45	1.57	5 (17%)
24	ADP	В	800	-	24,29,29	0.92	0	29,45,45	1.57	6 (20%)
24	ADP	D	800	-	24,29,29	0.89	0	29,45,45	1.67	6 (20%)
24	ADP	F	800	-	24,29,29	0.88	0	29,45,45	1.63	4 (13%)
25	ATP	Н	401	-	26,33,33	0.96	1 (3%)	31,52,52	1.69	6 (19%)
24	ADP	J	800	-	24,29,29	0.94	1 (4%)	29,45,45	1.66	5 (17%)
24	ADP	A	800	-	24,29,29	0.97	1 (4%)	29,45,45	1.41	4 (13%)
24	ADP	Е	800	-	24,29,29	0.95	1 (4%)	29,45,45	1.54	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
24	ADP	С	800	-	-	5/12/32/32	0/3/3/3
24	ADP	G	800	-	-	5/12/32/32	0/3/3/3
24	ADP	I	800	-	-	7/12/32/32	0/3/3/3
24	ADP	В	800	-	-	4/12/32/32	0/3/3/3
24	ADP	D	800	-	-	5/12/32/32	0/3/3/3
24	ADP	F	800	-	-	3/12/32/32	0/3/3/3
25	ATP	Н	401	-	-	7/18/38/38	0/3/3/3
24	ADP	J	800	-	-	3/12/32/32	0/3/3/3
24	ADP	A	800	-	-	4/12/32/32	0/3/3/3
24	ADP	Е	800	-	-	2/12/32/32	0/3/3/3

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
24	A	800	ADP	C5-C4	2.20	1.46	1.40
24	J	800	ADP	C5-C4	2.16	1.46	1.40
24	I	800	ADP	C5-C4	2.15	1.46	1.40
24	Е	800	ADP	C5-C4	2.06	1.46	1.40
24	G	800	ADP	C5-C4	2.05	1.46	1.40

The worst 5 of 49 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
25	Н	401	ATP	PB-O3B-PG	-5.38	114.35	132.83
24	С	800	ADP	PA-O3A-PB	-4.62	116.96	132.83
24	J	800	ADP	PA-O3A-PB	-4.55	117.22	132.83
24	F	800	ADP	PA-O3A-PB	-4.39	117.77	132.83
25	Н	401	ATP	PA-O3A-PB	-4.02	119.05	132.83

There are no chirality outliers.

5 of 45 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
24	В	800	ADP	C5'-O5'-PA-O2A
24	С	800	ADP	C5'-O5'-PA-O2A
24	С	800	ADP	C5'-O5'-PA-O3A
24	D	800	ADP	C3'-C4'-C5'-O5'
24	Е	800	ADP	C3'-C4'-C5'-O5'


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
































5.7 Other polymers (i)

There are no such residues in this entry.

Torsions



Rings

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
13	Y	12
7	N	5
6	M	5
8	О	3
8	P	3
9	R	2
9	Q	2

The worst 5 of 32 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	N	367:UNK	С	401:UNK	N	98.98
1	N	171:UNK	С	201:UNK	N	96.72
1	M	171:UNK	С	201:UNK	N	91.43
1	M	367:UNK	С	401:UNK	N	84.01
1	M	625:UNK	С	701:UNK	N	69.30




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-4177. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

6.2.1 Primary map








Y Index: 300


6.2.2 Raw map







Y Index: 300



Z Index: 300

The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map








Y Index: 325




Z Index: 274


6.3.2 Raw map



X Index: 289



Y Index: 325




Z Index: 274

The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.06. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $1246~\mathrm{nm^3}$; this corresponds to an approximate mass of $1125~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.149 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.149 Å $^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
rtesolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	6.70	-	-	
Author-provided FSC curve	-	-	-	
Unmasked-calculated*	9.71	26.18	11.07	

^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 9.71 differs from the reported value 6.7 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-4177 and PDB model 6F38. Per-residue inclusion information can be found in section 3 on page 12.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.06 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.06).



9.4 Atom inclusion (i)



At the recommended contour level, 81% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.06) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7938	0.2300
A	0.9708	0.2770
В	0.9854	0.2960
С	0.9659	0.2910
D	0.9865	0.2990
E	0.9794	0.2990
F	0.9908	0.3010
G	0.9919	0.2930
Н	0.9930	0.2970
I	0.9816	0.2740
J	0.9827	0.2810
K	0.9739	0.2600
L	0.9894	0.2780
M	0.9581	0.2450
N	0.9526	0.2620
О	0.9907	0.3010
Р	0.9969	0.2980
Q	0.4529	0.1500
R	0.9908	0.2830
U	0.8766	0.1760
V	0.9914	0.2640
X	0.7040	0.2110
Y	0.9538	0.3210
Z	0.9538	0.2960
a	0.9498	0.2980
b	0.9268	0.2800
c	0.8967	0.3030
d	0.8992	0.3020
e	0.6712	0.1900
f	0.8407	0.2310
g	0.7659	0.1410
h	0.9448	0.2210
i	0.0136	0.0040
j	0.4949	0.1510
k	0.8506	0.1460



Continued on next page...



$Continued\ from\ previous\ page...$

Chain	Atom inclusion	Q-score
l	0.7035	0.1570
m	0.6973	0.2240
n	0.6128	0.2010
О	0.8210	0.1830
p	0.5114	0.1360
q	0.1372	0.1770
r	0.0136	0.1600
s	0.4978	0.1090
t	0.6277	0.1230
X	0.7009	0.1930
Z	0.8906	0.2550

