



wwPDB X-ray Structure Validation Summary Report ⓘ

May 31, 2025 – 01:18 pm BST

PDB ID : 9FHG / pdb_00009fhg
Title : Crystallographic structure of AcrB V612N in LTO state
Authors : Lazarova, M.; Pos, K.M.
Deposited on : 2024-05-27
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2511 (3.00-3.00)
Clashscore	180529	2866 (3.00-3.00)
Ramachandran outliers	177936	2778 (3.00-3.00)
Sidechain outliers	177891	2781 (3.00-3.00)
RSRZ outliers	164620	2523 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1057	 3% 70% 28% ..
1	B	1057	 4% 70% 27% .
1	C	1057	 5% 70% 27% ..
2	D	169	 % 58% 34% . 8%
2	E	169	 2% 59% 31% . 10%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 25972 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multidrug efflux pump subunit AcrB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1044	Total	C	N	O	S	0	0	0
			7944	5105	1316	1479	44			
1	B	1033	Total	C	N	O	S	0	0	0
			7850	5051	1296	1459	44			
1	C	1033	Total	C	N	O	S	0	0	0
			7850	5051	1296	1459	44			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	612	ASN	VAL	engineered mutation	UNP P31224
A	1050	LEU	-	expression tag	UNP P31224
A	1051	GLU	-	expression tag	UNP P31224
A	1052	HIS	-	expression tag	UNP P31224
A	1053	HIS	-	expression tag	UNP P31224
A	1054	HIS	-	expression tag	UNP P31224
A	1055	HIS	-	expression tag	UNP P31224
A	1056	HIS	-	expression tag	UNP P31224
A	1057	HIS	-	expression tag	UNP P31224
B	612	ASN	VAL	engineered mutation	UNP P31224
B	1050	LEU	-	expression tag	UNP P31224
B	1051	GLU	-	expression tag	UNP P31224
B	1052	HIS	-	expression tag	UNP P31224
B	1053	HIS	-	expression tag	UNP P31224
B	1054	HIS	-	expression tag	UNP P31224
B	1055	HIS	-	expression tag	UNP P31224
B	1056	HIS	-	expression tag	UNP P31224
B	1057	HIS	-	expression tag	UNP P31224
C	612	ASN	VAL	engineered mutation	UNP P31224
C	1050	LEU	-	expression tag	UNP P31224
C	1051	GLU	-	expression tag	UNP P31224
C	1052	HIS	-	expression tag	UNP P31224
C	1053	HIS	-	expression tag	UNP P31224

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1054	HIS	-	expression tag	UNP P31224
C	1055	HIS	-	expression tag	UNP P31224
C	1056	HIS	-	expression tag	UNP P31224
C	1057	HIS	-	expression tag	UNP P31224


- Molecule 2 is a protein called DARPIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	156	Total	C	N	O	S	0	0	0
			1177	741	206	229	1			
2	E	152	Total	C	N	O	S	0	0	0
			1151	726	202	222	1			

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Multidrug efflux pump subunit AcrB



Chain B: 4% 70% 27%

100 data points are displayed in a grid. The top row labels are: V1022, P1023, F1025, F1026, V1027, P1028, V1029, R1030, A939, K940, T943, E947, L952, P961, D966, V968, R969, A970, H971, L972, P973, R974, L976, P977, F982, R983, L984, G985, V986, P987, R988, L989, V990, P991, A995, W996, S997, Q1000, V1003, V1007, A1014, T1015, A1018, I1019. The bottom row labels are: SER, ARG, LYS, ASN, GLU, ASP, ASP, ILE, HIS, HIS, SER, HIS, THR, VAL, ASP, HIS, HIS, LEU, HIS, HIS, HIS, HIS, HIS, HIS, F982, R983, L984, G985, V986, P987, R988, L989, V990, P991, A995, W996, S997, Q1000, V1003, V1007, A1014, T1015, A1018, I1019.

Chain C:

5%


70%

27%


..

Item	Category
M1	Yellow
P2	Yellow
N3	Yellow
F4	Yellow
F5	Yellow
I6	Yellow
D7	Yellow
R8	Yellow
P9	Yellow
I10	Yellow
I15	Yellow
I19	Yellow
M20	Yellow
L21	Yellow
A22	Yellow
G23	Yellow
G24	Yellow
L25	Yellow
V32	Yellow
A33	Yellow
Q34	Yellow
I38	Yellow
A39	Yellow
S46	Yellow
A47	Yellow
S48	Yellow
Y49	Yellow
D53	Yellow
A54	Yellow
A55	Yellow
T56	Yellow
V57	Yellow
T62	Yellow
M68	Yellow
M69	Yellow
M70	Yellow
G71	Yellow
M76	Yellow
S80	Yellow
N81	Yellow
S82	Yellow
G86	Yellow
T87	Yellow
V88	Yellow
Q89	Yellow
I90	Yellow
E95	Yellow


Item	Category
D101	Yellow
A102	Yellow
A103	Yellow
Q104	Yellow
V105	Yellow
Q106	Yellow
V107	Yellow
Q112	Yellow
L113	Yellow
A114	Yellow
M115	Yellow
L117	Yellow
L118	Yellow
P119	Yellow
Q120	Yellow
E121	Yellow
V122	Yellow
Q123	Yellow
Q124	Yellow
Q125	Yellow
S128	Yellow
S133	Yellow
S134	Yellow
V142	Yellow
T145	Yellow
D146	Yellow
E152	Yellow
S155	Yellow
A159	Yellow
M162	Yellow
A165	Yellow
V175	Yellow
Q176	Yellow
L177	Yellow
S180	Yellow
Q181	Yellow
Y182	Yellow
A183	Yellow
M184	Yellow
R185	Yellow
I186	Yellow
M187	Yellow
M188	Yellow
P189	Yellow
I190	Yellow



• Molecule 2: DARPIN



• Molecule 2: DARPIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	145.81Å 161.40Å 245.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.50 – 3.00 49.50 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.50-3.00) 99.7 (49.50-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.20.1	Depositor
R, R_{free}	0.224 , 0.283 0.224 , 0.283	Depositor DCC
R_{free} test set	5793 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	75.9	Xtriage
Anisotropy	0.660	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 21.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	25972	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	1/8096 (0.0%)	0.65	1/10992 (0.0%)
1	B	0.44	1/8000 (0.0%)	0.64	1/10864 (0.0%)
1	C	0.41	1/8000 (0.0%)	0.63	0/10864
2	D	0.41	0/1196	0.64	0/1626
2	E	0.39	0/1170	0.61	0/1591
All	All	0.42	3/26462 (0.0%)	0.64	2/35937 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	426	PRO	CA-C	13.51	1.59	1.51
1	A	897	ILE	CA-CB	8.70	1.59	1.53
1	C	146	ASP	CA-C	5.67	1.60	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	426	PRO	O-C-N	10.69	126.22	121.31
1	A	461	GLY	N-CA-C	5.10	120.51	111.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7944	0	8081	220	0
1	B	7850	0	7998	208	0
1	C	7850	0	7998	212	0
2	D	1177	0	1159	47	0
2	E	1151	0	1136	42	0
All	All	25972	0	26372	698	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 698 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:153:SER:HB3	2:E:161:LEU:HD23	1.51	0.90
1:C:76:MET:HE3	1:C:95:GLU:HA	1.57	0.84
1:B:939:ALA:O	1:B:943:ILE:HG13	1.82	0.80
1:C:544:LEU:HA	1:C:547:ILE:HD12	1.64	0.78
1:C:15:ILE:O	1:C:19:ILE:HG13	1.82	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1042/1057 (99%)	972 (93%)	63 (6%)	7 (1%)	19	54
1	B	1031/1057 (98%)	972 (94%)	57 (6%)	2 (0%)	44	77
1	C	1031/1057 (98%)	961 (93%)	67 (6%)	3 (0%)	37	70
2	D	154/169 (91%)	139 (90%)	15 (10%)	0	100	100
2	E	150/169 (89%)	137 (91%)	13 (9%)	0	100	100
All	All	3408/3509 (97%)	3181 (93%)	215 (6%)	12 (0%)	30	66

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	71	GLY
1	A	126	GLY
1	A	678	THR
1	A	1037	ASN
1	A	388	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	850/863 (98%)	828 (97%)	22 (3%)	41	72
1	B	839/863 (97%)	819 (98%)	20 (2%)	44	74
1	C	839/863 (97%)	816 (97%)	23 (3%)	40	71
2	D	120/132 (91%)	116 (97%)	4 (3%)	33	67
2	E	117/132 (89%)	114 (97%)	3 (3%)	41	72
All	All	2765/2853 (97%)	2693 (97%)	72 (3%)	41	72

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	577	GLN
2	E	165	LEU
1	C	626	ILE
2	D	45	VAL
1	B	82	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	871	ASN
1	C	872	GLN
2	D	142	GLN
1	A	846	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	797	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	1044/1057 (98%)	0.10	36 (3%)	48	28	47, 80, 115, 136	0
1	B	1033/1057 (97%)	0.13	44 (4%)	40	23	53, 79, 104, 121	0
1	C	1033/1057 (97%)	0.14	50 (4%)	36	21	49, 73, 94, 109	0
2	D	156/169 (92%)	-0.02	2 (1%)	74	54	67, 80, 98, 108	0
2	E	152/169 (89%)	0.09	3 (1%)	64	43	71, 86, 107, 116	0
All	All	3418/3509 (97%)	0.12	135 (3%)	44	26	47, 77, 106, 136	0

The worst 5 of 135 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	618	ALA	6.2
1	B	284	GLN	5.9
1	B	407	ASP	5.3
1	A	537	SER	5.2
1	B	134	SER	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.