

wwPDB X-ray Structure Validation Summary Report (i)

Jun 29, 2025 - 08:34 am BST

PDB ID	:	$9\mathrm{FHY} \ / \ \mathrm{pdb} \ 00009\mathrm{fhy}$
Title	:	Bacteroides ovatus polysaccharide lyase family 38 (BoPL38) wild type in com-
		plex with an unsaturated mannuronic acid tetramer at pH 3.5
Authors	:	Tandrup, T.; Wilkens, C.
Deposited on	:	2024-05-28
Resolution	:	1.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.4, CSD as 541 be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
_		10.1	2%		
	A	404	86%	8%	6%
1	D	10.1	4%		_
	В	404	86%	8%	6%
1	C	10.1	5%		_
	С	404	88%	6%	6%
1	D	10.1	5%		_
	D	404	86%	8%	6%
2		9			
2	Ε	3	100%		



Mol	Chain	Length	Quality of chain
2	F	3	100%
2	G	3	100%
2	Н	3	100%



9FHY

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 13124 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	380	Total	С	Ν	0	\mathbf{S}	0	0 2	0
	А	300	3066	1961	519	573	13	0		0
1	В	380	Total	С	Ν	0	S	0	2	0
	ГБ		3064	1961	519	571	13			
1	С	380	Total	С	Ν	0	S	0	0	0
			3052	1952	516	571	13	0		0
1	1 D 380	280	Total	С	Ν	0	S	0	0	0
		380	3052	1952	516	571	13	U		0

• Molecule 1 is a protein called Alginate lyase family protein.

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	1	MET	-	initiating methionine	UNP A0A5M5BWR5
А	2	GLY	-	expression tag	UNP A0A5M5BWR5
А	3	SER	-	expression tag	UNP A0A5M5BWR5
А	4	SER	-	expression tag	UNP A0A5M5BWR5
А	5	HIS	-	expression tag	UNP A0A5M5BWR5
А	6	HIS	-	expression tag	UNP A0A5M5BWR5
A	7	HIS	-	expression tag	UNP A0A5M5BWR5
A	8	HIS	-	expression tag	UNP A0A5M5BWR5
A	9	HIS	-	expression tag	UNP A0A5M5BWR5
А	10	HIS	-	expression tag	UNP A0A5M5BWR5
A	11	SER	-	expression tag	UNP A0A5M5BWR5
A	12	SER	-	expression tag	UNP A0A5M5BWR5
A	13	GLY	-	expression tag	UNP A0A5M5BWR5
A	14	LEU	-	expression tag	UNP A0A5M5BWR5
А	15	VAL	-	expression tag	UNP A0A5M5BWR5
А	16	PRO	-	expression tag	UNP A0A5M5BWR5
А	17	ARG	-	expression tag	UNP A0A5M5BWR5
А	18	GLY	-	expression tag	UNP A0A5M5BWR5
А	19	SER	-	expression tag	UNP A0A5M5BWR5
А	20	HIS	-	expression tag	UNP A0A5M5BWR5
А	21	MET	-	expression tag	UNP A0A5M5BWR5



	Continued from previous page Comment Reference Chain Residue Modelled Actual Comment Reference									
Chain			Actual	Comment						
A	22	ALA	-	expression tag	UNP A0A5M5BWR5					
A	23	SER	-	expression tag	UNP A0A5M5BWR5					
B	1	MET	-	initiating methionine	UNP A0A5M5BWR5					
В	2	GLY	-	expression tag	UNP A0A5M5BWR5					
В	3	SER	-	expression tag	UNP A0A5M5BWR5					
В	4	SER	-	expression tag	UNP A0A5M5BWR5					
В	5	HIS	-	expression tag	UNP A0A5M5BWR5					
В	6	HIS	-	expression tag	UNP A0A5M5BWR5					
В	7	HIS	-	expression tag	UNP A0A5M5BWR5					
В	8	HIS	-	expression tag	UNP A0A5M5BWR5					
В	9	HIS	-	expression tag	UNP A0A5M5BWR5					
В	10	HIS	-	expression tag	UNP A0A5M5BWR5					
В	11	SER	-	expression tag	UNP A0A5M5BWR5					
В	12	SER	-	expression tag	UNP A0A5M5BWR5					
В	13	GLY	-	expression tag	UNP A0A5M5BWR5					
В	14	LEU	-	expression tag	UNP A0A5M5BWR5					
В	15	VAL	-	expression tag	UNP A0A5M5BWR5					
В	16	PRO	-	expression tag	UNP A0A5M5BWR5					
В	17	ARG	-	expression tag	UNP A0A5M5BWR5					
В	18	GLY	-	expression tag	UNP A0A5M5BWR5					
В	19	SER	-	expression tag	UNP A0A5M5BWR5					
В	20	HIS	-	expression tag	UNP A0A5M5BWR5					
В	21	MET	-	expression tag	UNP A0A5M5BWR5					
В	22	ALA	-	expression tag	UNP A0A5M5BWR5					
В	23	SER	-	expression tag	UNP A0A5M5BWR5					
С	1	MET	_	initiating methionine	UNP A0A5M5BWR5					
С	2	GLY	-	expression tag	UNP A0A5M5BWR5					
С	3	SER	_	expression tag	UNP A0A5M5BWR5					
С	4	SER	-	expression tag	UNP A0A5M5BWR5					
С	5	HIS	_	expression tag	UNP A0A5M5BWR5					
С	6	HIS	_	expression tag	UNP A0A5M5BWR5					
С	7	HIS	-	expression tag	UNP A0A5M5BWR5					
С	8	HIS	-	expression tag	UNP A0A5M5BWR5					
C	9	HIS	-	expression tag	UNP A0A5M5BWR5					
С	10	HIS	-	expression tag	UNP A0A5M5BWR5					
С	11	SER	-	expression tag	UNP A0A5M5BWR5					
С	12	SER	-	expression tag	UNP A0A5M5BWR5					
С	13	GLY	-	expression tag	UNP A0A5M5BWR5					
C	14	LEU	_	expression tag	UNP A0A5M5BWR5					
C	15	VAL	_	expression tag	UNP A0A5M5BWR5					
C	16	PRO	_	expression tag	UNP A0A5M5BWR5					
C	10	ARG	_	expression tag	UNP A0A5M5BWR5					
~	· ·				entinued on nert nage					



Chain	Residue	Modelled	Actual	Comment	Reference
С	18	GLY	_	expression tag	UNP A0A5M5BWR5
С	19	SER	_	expression tag	UNP A0A5M5BWR5
С	20	HIS	-	expression tag	UNP A0A5M5BWR5
С	21	MET	-	expression tag	UNP A0A5M5BWR5
С	22	ALA	-	expression tag	UNP A0A5M5BWR5
С	23	SER	-	expression tag	UNP A0A5M5BWR5
D	1	MET	-	initiating methionine	UNP A0A5M5BWR5
D	2	GLY	-	expression tag	UNP A0A5M5BWR5
D	3	SER	-	expression tag	UNP A0A5M5BWR5
D	4	SER	-	expression tag	UNP A0A5M5BWR5
D	5	HIS	-	expression tag	UNP A0A5M5BWR5
D	6	HIS	-	expression tag	UNP A0A5M5BWR5
D	7	HIS	-	expression tag	UNP A0A5M5BWR5
D	8	HIS	-	expression tag	UNP A0A5M5BWR5
D	9	HIS	-	expression tag	UNP A0A5M5BWR5
D	10	HIS	-	expression tag	UNP A0A5M5BWR5
D	11	SER	-	expression tag	UNP A0A5M5BWR5
D	12	SER	-	expression tag	UNP A0A5M5BWR5
D	13	GLY	-	expression tag	UNP A0A5M5BWR5
D	14	LEU	-	expression tag	UNP A0A5M5BWR5
D	15	VAL	-	expression tag	UNP A0A5M5BWR5
D	16	PRO	-	expression tag	UNP A0A5M5BWR5
D	17	ARG	-	expression tag	UNP A0A5M5BWR5
D	18	GLY	-	expression tag	UNP A0A5M5BWR5
D	19	SER	-	expression tag	UNP A0A5M5BWR5
D	20	HIS	-	expression tag	UNP A0A5M5BWR5
D	21	MET	-	expression tag	UNP A0A5M5BWR5
D	22	ALA	-	expression tag	UNP A0A5M5BWR5
D	23	SER	-	expression tag	UNP A0A5M5BWR5

• Molecule 2 is an oligosaccharide called 4-deoxy-alpha-L-erythro-hex-4-enopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
2	Е	3	Total C 36 18		0	0	0
2	F	3	Total C 36 18	O 18	0	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	G	3	Total C O 36 18 18	0	0	0
2	Н	3	Total C O 36 18 18	0	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	220	Total O 220 220	0	0
4	В	213	Total O 213 213	0	0
4	С	150	Total O 150 150	0	0
4	D	153	Total O 153 153	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Alginate lyase family protein



 \bullet Molecule 2: 4-deoxy-alpha-L-erythro-hex-4-enopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid

Chain E:

100%

BEM1 BEM2 MAW3

 \bullet Molecule 2: 4-deoxy-alpha-L-erythro-hex-4-enopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid

Chain F:

100%

BEM1 BEM2 MAW3

 \bullet Molecule 2: 4-deoxy-alpha-L-erythro-hex-4-enopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid

Chain G:

100%

BEM1 BEM2 MAV3

 \bullet Molecule 2: 4-deoxy-alpha-L-erythro-hex-4-enopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid-(1-4)-beta-D-mannopyranuronic acid

Chain H:

100%

BEM1 BEM2 MAW3



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	197.87Å 88.70Å 147.35Å	Depositor
a, b, c, α , β , γ	90.00° 120.39° 90.00°	Depositor
Resolution (Å)	58.48 - 1.76	Depositor
Resolution (A)	58.48 - 1.76	EDS
% Data completeness	81.4 (58.48-1.76)	Depositor
(in resolution range)	81.4 (58.48-1.76)	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.68 (at 1.76 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.21_5207	Depositor
D D.	0.217 , 0.244	Depositor
R, R_{free}	0.217 , 0.244	DCC
R_{free} test set	10866 reflections (4.97%)	wwPDB-VP
Wilson B-factor $(Å^2)$	32.5	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.36 , 27.6	EDS
L-test for twinning ²	$ \langle L \rangle = 0.46, \langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13124	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 11.85% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BEM, MAW, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.37	0/3150	0.52	0/4268	
1	В	0.37	0/3148	0.55	0/4266	
1	С	0.37	0/3130	0.53	0/4242	
1	D	0.36	0/3130	0.53	0/4242	
All	All	0.37	0/12558	0.53	0/17018	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3066	0	3041	22	0
1	В	3064	0	3044	25	0
1	С	3052	0	3022	18	0
1	D	3052	0	3022	19	0
2	Е	36	0	19	0	0
2	F	36	0	19	0	0
2	G	36	0	19	0	0
2	Н	36	0	19	0	0
3	В	5	0	0	0	0



Mol	Chain	Non-H	Non-H H(model)		Clashes	Symm-Clashes	
3	D	5	0	0	0	0	
4	А	220	0	0	7	0	
4	В	213	0	0	8	0	
4	С	150	0	0	2	0	
4	D	153	0	0	2	0	
All	All	13124	0	12205	85	0	

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:387:LYS:NZ	4:A:501:HOH:O	1.87	1.07
1:D:42:GLN:NE2	4:D:601:HOH:O	2.05	0.89
1:B:360:ARG:NH1	4:B:603:HOH:O	2.13	0.74
1:C:360:ARG:NH1	4:C:501:HOH:O	2.21	0.72
1:B:162:LYS:NZ	4:B:602:HOH:O	2.12	0.71

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percen	tiles
1	А	380/404~(94%)	374~(98%)	6(2%)	0	100	100
1	В	380/404~(94%)	373~(98%)	7~(2%)	0	100	100
1	С	378/404~(94%)	371~(98%)	7(2%)	0	100	100
1	D	378/404~(94%)	371~(98%)	7 (2%)	0	100	100
All	All	1516/1616~(94%)	1489~(98%)	27~(2%)	0	100	100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	326/344~(95%)	323~(99%)	3~(1%)	75	65	
1	В	326/344~(95%)	326 (100%)	0	100	100	
1	С	324/344~(94%)	322~(99%)	2 (1%)	84	78	
1	D	324/344~(94%)	322~(99%)	2 (1%)	84	78	
All	All	1300/1376~(94%)	1293 (100%)	7 (0%)	86	82	

5 of 7 residues with a non-rotameric side chain are listed below:

Mol	Chain	Chain Res	
1	С	148	LYS
1	С	198	VAL
1	D	175	VAL
1	D	45	GLU
1	А	184	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	173	GLN
1	D	375	GLN
1	В	173	GLN
1	В	315	GLN
1	В	358	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	ths	Bond angles		
MOI	туре	Ullalli	nes	S LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	BEM	Е	1	2	13,13,13	1.03	1 (7%)	18,19,19	1.49	2 (11%)
2	BEM	Е	2	2	12,12,13	1.06	0	14,17,19	1.47	3 (21%)
2	MAW	Е	3	2	10,11,12	1.33	1 (10%)	$13,\!15,\!17$	2.42	4 (30%)
2	BEM	F	1	2	13,13,13	0.98	0	18,19,19	1.18	1 (5%)
2	BEM	F	2	2	12,12,13	1.20	1 (8%)	14,17,19	1.33	3 (21%)
2	MAW	F	3	2	10,11,12	1.26	1 (10%)	$13,\!15,\!17$	2.01	2 (15%)
2	BEM	G	1	2	13,13,13	1.05	1 (7%)	18,19,19	1.31	2 (11%)
2	BEM	G	2	2	12,12,13	1.11	1 (8%)	14,17,19	1.08	0
2	MAW	G	3	2	10,11,12	1.23	0	$13,\!15,\!17$	1.86	3 (23%)
2	BEM	Н	1	2	13,13,13	1.04	1 (7%)	18,19,19	1.39	2 (11%)
2	BEM	Н	2	2	12,12,13	1.18	1 (8%)	14,17,19	0.88	0
2	MAW	Н	3	2	10,11,12	1.33	2 (20%)	$13,\!15,\!17$	2.04	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEM	Е	1	2	-	0/4/24/24	0/1/1/1
2	BEM	Е	2	2	-	0/4/21/24	0/1/1/1
2	MAW	Е	3	2	-	0/4/17/20	0/1/1/1
2	BEM	F	1	2	-	0/4/24/24	0/1/1/1
2	BEM	F	2	2	-	0/4/21/24	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAW	F	3	2	-	0/4/17/20	0/1/1/1
2	BEM	G	1	2	-	0/4/24/24	0/1/1/1
2	BEM	G	2	2	-	0/4/21/24	0/1/1/1
2	MAW	G	3	2	-	0/4/17/20	0/1/1/1
2	BEM	Н	1	2	-	0/4/24/24	0/1/1/1
2	BEM	Н	2	2	-	0/4/21/24	0/1/1/1
2	MAW	Н	3	2	-	0/4/17/20	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
2	F	3	MAW	C5-C6	2.74	1.55	1.48
2	F	2	BEM	O5-C1	-2.70	1.39	1.43
2	Е	3	MAW	C5-C6	2.65	1.55	1.48
2	Е	1	BEM	O6B-C6	-2.29	1.23	1.30
2	Н	2	BEM	O6B-C6	-2.24	1.23	1.30

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	Ε	3	MAW	O5-C5-C4	-6.35	119.45	124.81
2	F	3	MAW	O5-C5-C4	-5.34	120.30	124.81
2	G	3	MAW	O5-C5-C4	-4.55	120.97	124.81
2	Н	3	MAW	O5-C5-C4	-4.47	121.04	124.81
2	Н	3	MAW	C2-C3-C4	-3.87	107.02	112.32

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mo	Type	Chain	Res	Link	B	gths	В	ond ang	gles	
INIO!	Type	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	SO4	D	501	-	4,4,4	0.66	0	$6,\!6,\!6$	0.07	0
3	SO4	В	501	-	4,4,4	0.66	0	$6,\!6,\!6$	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	380/404~(94%)	0.35	8 (2%) 63 69	20, 32, 43, 53	2 (0%)
1	В	380/404~(94%)	0.40	15 (3%) 44 50	17, 33, 44, 54	2 (0%)
1	С	380/404~(94%)	0.56	21 (5%) 32 35	25, 36, 48, 58	0
1	D	380/404~(94%)	0.58	19 (5%) 35 38	24, 37, 49, 55	0
All	All	1520/1616~(94%)	0.47	63 (4%) 42 48	17, 34, 47, 58	4 (0%)

The worst 5 of 63 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	74	LYS	3.8
1	С	50	ARG	3.6
1	В	191[A]	ARG	3.3
1	D	23	SER	3.2
1	D	360	ARG	3.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	BEM	Н	1	13/13	0.84	0.13	40,46,54,54	0
2	BEM	Е	1	13/13	0.90	0.09	32,39,46,51	0




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BEM	F	1	13/13	0.92	0.09	$35,\!41,\!51,\!51$	0
2	BEM	G	1	13/13	0.93	0.09	$38,\!46,\!53,\!56$	0
2	BEM	Н	2	12/13	0.93	0.09	32,37,39,39	0
2	MAW	Н	3	11/12	0.95	0.08	30,34,36,36	0
2	BEM	G	2	12/13	0.96	0.06	32,37,39,40	0
2	MAW	G	3	11/12	0.96	0.07	31,34,36,36	0
2	BEM	Е	2	12/13	0.96	0.07	26,30,33,35	0
2	MAW	F	3	11/12	0.96	0.06	26,29,31,31	0
2	MAW	Е	3	11/12	0.96	0.06	26,27,30,30	0
2	BEM	F	2	12/13	0.97	0.05	30,32,34,35	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.











6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	D	501	5/5	0.34	0.17	103,106,122,123	0
3	SO4	В	501	5/5	0.83	0.17	36,41,45,46	5

6.5 Other polymers (i)

There are no such residues in this entry.

