

# wwPDB EM Validation Summary Report (i)

Feb 5, 2025 – 04:28 pm GMT

PDB ID : 9G0O

EMDB ID : EMD-50941

Title : Xenopus borealis undecorated microtubule - 14 protofilament, 3-start helix

Authors: Troman, L.A.; Moores, C.A.

Deposited on : 2024-07-08

Resolution : 3.30 Å(reported)

Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113

Mogul : 1.8.4, CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : 1.9.13

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$		
Ramachandran outliers	207382	16835		
Sidechain outliers	206894	16415		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	445	96%	
1	В	445	96%	• •
1	С	445		• •
1	D	445		
1	Е	445		
1	F	445		
2	a	449	96%	•
2	b	449	96%	•
2	c	449	96%	•

Continued on next page...



Continued from previous page...

	J	1	1 9	
Mol	Chain	Length	Quality of chain	
2	d	449	96%	
2	е	449	96%	
2	f	449	96%	



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 80238 atoms, of which 39306 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Tubulin beta chain.

Mol	Chain	Residues			Atom	.S			AltConf	Trace
1	A	430	Total	С	Н	N	О	S	0	0
1	Λ	450	6631	2119	3258	578	650	26	0	0
1	В	430	Total	С	Н	N	О	S	0	0
1	Б	450	6631	2119	3258	578	650	26	0	0
1	С	430	Total	С	Н	N	О	S	0	0
1		450	6631	2119	3258	578	650	26	0	0
1	D	430	Total	С	Н	N	О	S	0	0
1	ט	450	6631	2119	3258	578	650	26	0	0
1	Е	430	Total	С	Н	N	О	S	0	0
1	<u> 1</u> 2	450	6631	2119	3258	578	650	26	0	0
1	F	430	Total	С	Н	N	О	S	0	0
1	1'	430	6631	2119	3258	578	650	26		

• Molecule 2 is a protein called Tubulin alpha chain.

Mol	Chain	Residues			Atom	S			AltConf	Trace
2		432	Total	С	Н	N	О	S	0	0
	a	432	6681	2139	3293	577	648	24	0	0
2	b	432	Total	С	Н	N	О	S	0	0
2	D	432	6681	2139	3293	577	648	24	U	U
2		432	Total	С	Н	N	О	S	0	0
	С	402	6681	2139	3293	577	648	24	0	U
2	d	432	Total	С	Н	N	О	S	0	0
2	u	402	6681	2139	3293	577	648	24		0
2	e	432	Total	С	Н	N	О	S	0	0
	C	432	6681	2139	3293	577	648	24	0	U
2	f	432	Total	С	Н	N	О	S	0	0
	1	402	6681	2139	3293	577	648	24		

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	5	LEU	ILE	conflict	UNP Q5EB23
a	32	ARG	GLN	conflict	UNP Q5EB23

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
a	36	VAL	ILE	conflict	UNP Q5EB23
a	37	SER	PRO	conflict	UNP Q5EB23
a	39	ASP	GLU	conflict	UNP Q5EB23
a	40	HIS	LYS	conflict	UNP Q5EB23
a	44	ILE	THR	conflict	UNP Q5EB23
a	77	VAL	ILE	conflict	UNP Q5EB23
a	81	PRO	HIS	conflict	UNP Q5EB23
a	116	THR	SER	conflict	UNP Q5EB23
a	118	MET	LEU	conflict	UNP Q5EB23
a	339	THR	SER	conflict	UNP Q5EB23
b	5	LEU	ILE	conflict	UNP Q5EB23
b	32	ARG	GLN	conflict	UNP Q5EB23
b	36	VAL	ILE	conflict	UNP Q5EB23
b	37	SER	PRO	conflict	UNP Q5EB23
b	39	ASP	GLU	conflict	UNP Q5EB23
b	40	HIS	LYS	conflict	UNP Q5EB23
b	44	ILE	THR	conflict	UNP Q5EB23
b	77	VAL	ILE	conflict	UNP Q5EB23
b	81	PRO	HIS	conflict	UNP Q5EB23
b	116	THR	SER	conflict	UNP Q5EB23
b	118	MET	LEU	conflict	UNP Q5EB23
b	339	THR	SER	conflict	UNP Q5EB23
c	5	LEU	ILE	conflict	UNP Q5EB23
С	32	ARG	GLN	conflict	UNP Q5EB23
c	36	VAL	ILE	conflict	UNP Q5EB23
c	37	SER	PRO	conflict	UNP Q5EB23
c	39	ASP	GLU	conflict	UNP Q5EB23
c	40	HIS	LYS	conflict	UNP Q5EB23
С	44	ILE	THR	conflict	UNP Q5EB23
С	77	VAL	ILE	conflict	UNP Q5EB23
С	81	PRO	HIS	conflict	UNP Q5EB23
С	116	THR	SER	conflict	UNP Q5EB23
С	118	MET	LEU	conflict	UNP Q5EB23
c	339	THR	SER	conflict	UNP Q5EB23
d	5	LEU	ILE	conflict	UNP Q5EB23
d	32	ARG	GLN	conflict	UNP Q5EB23
d	36	VAL	ILE	conflict	UNP Q5EB23
d	37	SER	PRO	conflict	UNP Q5EB23
d	39	ASP	GLU	conflict	UNP Q5EB23
d	40	HIS	LYS	conflict	UNP Q5EB23
d	44	ILE	THR	conflict	UNP Q5EB23
d	77	VAL	ILE	conflict	UNP Q5EB23

Continued on next page...




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
d	81	PRO	HIS	conflict	UNP Q5EB23
d	116	THR	SER	conflict	UNP Q5EB23
d	118	MET	LEU	conflict	UNP Q5EB23
d	339	THR	SER	conflict	UNP Q5EB23
е	5	LEU	ILE	conflict	UNP Q5EB23
е	32	ARG	GLN	conflict	UNP Q5EB23
е	36	VAL	ILE	conflict	UNP Q5EB23
е	37	SER	PRO	conflict	UNP Q5EB23
е	39	ASP	GLU	conflict	UNP Q5EB23
е	40	HIS	LYS	conflict	UNP Q5EB23
е	44	ILE	THR	conflict	UNP Q5EB23
е	77	VAL	ILE	conflict	UNP Q5EB23
е	81	PRO	HIS	conflict	UNP Q5EB23
е	116	THR	SER	conflict	UNP Q5EB23
е	118	MET	LEU	conflict	UNP Q5EB23
е	339	THR	SER	conflict	UNP Q5EB23
f	5	LEU	ILE	conflict	UNP Q5EB23
f	32	ARG	GLN	conflict	UNP Q5EB23
f	36	VAL	ILE	conflict	UNP Q5EB23
f	37	SER	PRO	conflict	UNP Q5EB23
f	39	ASP	GLU	conflict	UNP Q5EB23
f	40	HIS	LYS	conflict	UNP Q5EB23
f	44	ILE	THR	conflict	UNP Q5EB23
f	77	VAL	ILE	conflict	UNP Q5EB23
f	81	PRO	HIS	conflict	UNP Q5EB23
f	116	THR	SER	conflict	UNP Q5EB23
f	118	MET	LEU	conflict	UNP Q5EB23
f	339	THR	SER	conflict	UNP Q5EB23

• Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues		Ato	oms			AltConf
3	A	1	Total	С	N	О	Р	0
3	Λ	1	28	10	5	11	2	U
3	В	1	Total	С	N	О	Р	0
3	Ъ	1	28	10	5	11	2	U
3	C	1	Total	С	N	О	Р	0
J		1	28	10	5	11	2	U
3	D	1	Total	С	N	О	Р	0
	D	1	28	10	5	11	2	U
3	E	1	Total	$\mathbf{C}$	N	Ο	Р	0
	Ш	1	28	10	5	11	2	O
3	F	1	Total	$\mathbf{C}$	N	Ο	Р	0
	I.	1	28	10	5	11	2	

 $\bullet$  Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3)$  (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Ato	oms			AltConf	
4		1	Total	С	N	О	Р	0	
4	a	1	32	10	5	14	3	0	
4	b	1	Total	С	N	О	Р	0	
4	D	1	32	10	5	14	3	0	
4	c	1	Total	С	N	О	Р	0	
4	C	1	32	10	5	14	3	U	
4	d	1	Total	С	N	О	Р	0	
4	u	1	32	10	5	14	3	U	
4	e	1	Total	С	N	Ο	Р	0	
4	C	1	32	10	5	14	3	U	
1	f	1	Total	С	N	Ο	Р	0	
4	1	1	32	10	5	14	3	U	

 $\bullet$  Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
5	a	1	Total Mg 1 1	0
5	b	1	Total Mg 1 1	0
5	С	1	Total Mg 1 1	0
5	d	1	Total Mg 1 1	0
5	е	1	Total Mg 1 1	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
5	f	1	Total Mg 1 1	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tubulin beta chain

Chain A: 96%



• Molecule 1: Tubulin beta chain

Chain B: 96% ...




• Molecule 1: Tubulin beta chain

Chain C: 96% · ·




• Molecule 1: Tubulin beta chain

Chain D: 96% ...




• Molecule 1: Tubulin beta chain

Chain E:



• Molecule 1: Tubulin beta chain





# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	255980	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	47.4	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	2100	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.069	Depositor
Minimum map value	-0.031	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.004	Depositor
Recommended contour level	0.015	Depositor
Map size (Å)	443.872, 443.872, 443.872	wwPDB
Map dimensions	416, 416, 416	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.067, 1.067, 1.067	Depositor



# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GDP, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.25	0/3448	0.48	0/4673
1	В	0.25	0/3448	0.48	0/4673
1	С	0.25	0/3448	0.48	0/4673
1	D	0.25	0/3448	0.48	0/4673
1	Е	0.25	0/3448	0.48	0/4673
1	F	0.25	0/3448	0.48	0/4673
2	a	0.24	0/3463	0.46	0/4693
2	b	0.24	0/3463	0.46	0/4693
2	c	0.24	0/3463	0.46	0/4693
2	d	0.24	0/3463	0.46	0/4693
2	е	0.24	0/3463	0.46	0/4693
2	f	0.24	0/3463	0.46	0/4693
All	All	0.25	0/41466	0.47	0/56196

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	428/445~(96%)	413 (96%)	15 (4%)	0	100	100
1	В	428/445~(96%)	413 (96%)	15 (4%)	0	100	100
1	С	428/445 (96%)	413 (96%)	15 (4%)	0	100	100
1	D	428/445 (96%)	413 (96%)	15 (4%)	0	100	100
1	E	428/445 (96%)	413 (96%)	15 (4%)	0	100	100
1	F	428/445 (96%)	413 (96%)	15 (4%)	0	100	100
2	a	428/449 (95%)	415 (97%)	13 (3%)	0	100	100
2	b	428/449 (95%)	415 (97%)	13 (3%)	0	100	100
2	c	428/449 (95%)	415 (97%)	13 (3%)	0	100	100
2	d	428/449 (95%)	415 (97%)	13 (3%)	0	100	100
2	е	428/449 (95%)	415 (97%)	13 (3%)	0	100	100
2	f	428/449 (95%)	415 (97%)	13 (3%)	0	100	100
All	All	5136/5364 (96%)	4968 (97%)	168 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$368/380 \ (97\%)$	363 (99%)	5 (1%)	62 78
1	В	368/380 (97%)	363 (99%)	5 (1%)	62 78

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	368/380~(97%)	363 (99%)	5 (1%)	62 78
1	D	368/380 (97%)	363 (99%)	5 (1%)	62 78
1	E	368/380~(97%)	363 (99%)	5 (1%)	62 78
1	F	368/380 (97%)	363 (99%)	5 (1%)	62 78
2	a	368/380~(97%)	367 (100%)	1 (0%)	91 94
2	b	368/380~(97%)	367 (100%)	1 (0%)	91 94
2	c	368/380~(97%)	367 (100%)	1 (0%)	91 94
2	d	368/380 (97%)	367 (100%)	1 (0%)	91 94
2	e	368/380~(97%)	367 (100%)	1 (0%)	91 94
2	f	368/380 (97%)	367 (100%)	1 (0%)	91 94
All	All	4416/4560 (97%)	4380 (99%)	36 (1%)	77 87

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	364	SER
2	f	111	LYS
1	F	386	THR
2	С	111	LYS
1	С	137	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	347	ASN
2	е	60	HIS
1	С	347	ASN
1	D	204	ASN
1	D	347	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 6 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Т	Clasia.	Das	T !1.	Во	ond leng	ths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GTP	e	502	5	26,34,34	1.17	2 (7%)	32,54,54	1.61	7 (21%)
4	GTP	d	502	5	26,34,34	1.17	2 (7%)	32,54,54	1.61	7 (21%)
3	GDP	F	501	-	24,30,30	0.97	1 (4%)	30,47,47	1.32	4 (13%)
4	GTP	c	502	5	26,34,34	1.17	2 (7%)	32,54,54	1.60	7 (21%)
3	GDP	С	501	-	24,30,30	0.97	1 (4%)	30,47,47	1.31	4 (13%)
4	GTP	f	502	5	26,34,34	1.18	2 (7%)	32,54,54	1.61	7 (21%)
3	GDP	Е	501	-	24,30,30	0.97	1 (4%)	30,47,47	1.32	4 (13%)
3	GDP	A	501	-	24,30,30	0.97	1 (4%)	30,47,47	1.32	4 (13%)
3	GDP	D	501	-	24,30,30	0.98	1 (4%)	30,47,47	1.31	4 (13%)
4	GTP	a	501	5	26,34,34	1.17	2 (7%)	32,54,54	1.61	7 (21%)
3	GDP	В	501	-	24,30,30	0.97	1 (4%)	30,47,47	1.32	4 (13%)
4	GTP	b	501	5	26,34,34	1.18	2 (7%)	32,54,54	1.61	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	е	502	5	-	9/18/38/38	0/3/3/3
4	GTP	d	502	5	-	9/18/38/38	0/3/3/3
3	GDP	F	501	-	-	4/12/32/32	0/3/3/3
4	GTP	С	502	5	-	9/18/38/38	0/3/3/3
3	GDP	С	501	-	-	4/12/32/32	0/3/3/3
4	GTP	f	502	5	-	9/18/38/38	0/3/3/3
3	GDP	E	501	-	-	4/12/32/32	0/3/3/3
3	GDP	A	501	-	-	4/12/32/32	0/3/3/3
3	GDP	D	501	-	-	4/12/32/32	0/3/3/3
4	GTP	a	501	5	-	9/18/38/38	0/3/3/3
3	GDP	В	501	-	-	4/12/32/32	0/3/3/3
4	GTP	b	501	5	-	9/18/38/38	0/3/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
4	b	501	GTP	C5-C6	-4.31	1.38	1.47
4	f	502	GTP	C5-C6	-4.31	1.38	1.47
4	a	501	GTP	C5-C6	-4.31	1.38	1.47
4	е	502	GTP	C5-C6	-4.29	1.38	1.47
4	d	502	GTP	C5-C6	-4.26	1.38	1.47

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	501	GDP	PA-O3A-PB	-3.79	119.82	132.83
3	В	501	GDP	PA-O3A-PB	-3.78	119.86	132.83
3	D	501	GDP	PA-O3A-PB	-3.78	119.86	132.83
3	С	501	GDP	PA-O3A-PB	-3.76	119.91	132.83
3	Е	501	GDP	PA-O3A-PB	-3.76	119.91	132.83

There are no chirality outliers.

5 of 78 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	GDP	C5'-O5'-PA-O1A
3	A	501	GDP	C5'-O5'-PA-O2A
3	В	501	GDP	C5'-O5'-PA-O1A
3	В	501	GDP	C5'-O5'-PA-O2A
3	С	501	GDP	C5'-O5'-PA-O1A





There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











































### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




### 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-50941. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



#### 6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




# 6.2 Central slices (i)

#### 6.2.1 Primary map








X Index: 208


Y Index: 208

Z Index: 208


#### 6.2.2 Raw map







Y Index: 208




Z Index: 208


The images above show central slices of the map in three orthogonal directions.




### 6.3 Largest variance slices (i)

#### 6.3.1 Primary map









X Index: 110


Y Index: 105

Z Index: 156

#### 6.3.2 Raw map







X Index: 109

Y Index: 106


Z Index: 194

The images above show the largest variance slices of the map in three orthogonal directions.




### 6.4 Orthogonal standard-deviation projections (False-color) (i)

#### 6.4.1 Primary map



#### 6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




# 6.5 Orthogonal surface views (i)

#### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.015. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




### 6.6 Mask visualisation (i)


This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

### $6.6.1 \quad \mathrm{emd\_50941\_msk\_1.map} \ \ \mathbf{\mathring{i}}$








# 7 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 7.2 Volume estimate (i)




The volume at the recommended contour level is  $1117 \text{ nm}^3$ ; this corresponds to an approximate mass of 1009 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum (i)




<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.303  $\rm \mathring{A}^{-1}$ 



## 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.303  $\rm \mathring{A}^{-1}$ 



### 8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
rtesolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.30	-	-
Author-provided FSC curve	3.32	3.92	3.35
Unmasked-calculated*	4.20	7.60	4.28


<sup>\*</sup>Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.20 differs from the reported value 3.3 by more than 10 %



## 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-50941 and PDB model 9G0O. Per-residue inclusion information can be found in section 3 on page 10.


### 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.015).



### 9.4 Atom inclusion (i)



At the recommended contour level, 77% of all backbone atoms, 77% of all non-hydrogen atoms, are inside the map.



### 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.015) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7720	0.4460
A	0.7650	0.4140
В	0.7920	0.4710
С	0.7870	0.4390
D	0.7730	0.4220
E	0.8030	0.4790
F	0.7950	0.4510
a	0.7670	0.4270
b	0.7970	0.4720
С	0.7790	0.4440
d	0.7480	0.4220
e	0.7890	0.4730
f	0.7730	0.4420



