



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:17 PM UTC

PDB ID : 1G1X / pdb_00001g1x
Title : STRUCTURE OF RIBOSOMAL PROTEINS S15, S6, S18, AND 16S RIBOSOMAL RNA
Authors : Agalarov, S.C.; Prasad, G.S.; Funke, P.M.; Stout, C.D.; Williamson, J.R.
Deposited on : 2000-10-13
Resolution : 2.60 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RNA backbone	3983	1014 (2.84-2.36)





The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	D	41	41% 44% 7% 5% .
1	I	41	29% 41% 27% .
2	E	44	27% 43% 25% 5%
2	J	44	23% 52% 14% 11%
3	A	98	32% 50% 17% .
3	F	98	15% 58% 21% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	B	88	 40% 44% 15% .
4	G	88	 34% 50% 11% 5%
5	C	88	 11% 28% 11% . 48%
5	H	88	 13% 34% 7% . 45%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S RIBOSOMAL RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	D	40	Total 880	C 391	N 178	O 272	P 39	0	0	0
1	I	41	Total 902	C 401	N 183	O 278	P 40	0	0	0

- Molecule 2 is a RNA chain called 16S RIBOSOMAL RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	E	44	Total 932	C 417	N 167	O 305	P 43	0	0	0
2	J	44	Total 932	C 417	N 167	O 305	P 43	0	0	0

- Molecule 3 is a protein called 30S RIBOSOMAL PROTEIN S6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	98	Total 825	C 521	N 151	O 150	S 3	0	0	0
3	F	98	Total 825	C 521	N 151	O 150	S 3	0	0	0

- Molecule 4 is a protein called 30S RIBOSOMAL PROTEIN S15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	88	Total 737	C 460	N 147	O 128	S 2	0	0	0
4	G	88	Total 737	C 460	N 147	O 128	S 2	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	79	GLU	ALA	conflict	UNP Q5SJ76
B	80	ILE	LEU	conflict	UNP Q5SJ76
B	81	VAL	ILE	conflict	UNP Q5SJ76
B	86	LEU	ILE	conflict	UNP Q5SJ76
G	79	GLU	ALA	conflict	UNP Q5SJ76
G	80	ILE	LEU	conflict	UNP Q5SJ76
G	81	VAL	ILE	conflict	UNP Q5SJ76
G	86	LEU	ILE	conflict	UNP Q5SJ76

- Molecule 5 is a protein called 30S RIBOSOMAL PROTEIN S18.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	C	46	379	245	73	61	0	0	0
5	H	48	394	254	76	64	0	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: 16S RIBOSOMAL RNA

Chain D: 



- Molecule 1: 16S RIBOSOMAL RNA

Chain I: 




- Molecule 2: 16S RIBOSOMAL RNA

Chain E: 



- Molecule 2: 16S RIBOSOMAL RNA

Chain J: 





- Molecule 3: 30S RIBOSOMAL PROTEIN S6

Chain A: 




- Molecule 3: 30S RIBOSOMAL PROTEIN S6

Chain F: 



- Molecule 4: 30S RIBOSOMAL PROTEIN S15

Chain B: 




- Molecule 4: 30S RIBOSOMAL PROTEIN S15

Chain G: 





- Molecule 5: 30S RIBOSOMAL PROTEIN S18

Chain C: 



- Molecule 5: 30S RIBOSOMAL PROTEIN S18

Chain H: 



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	169.50Å 169.50Å 113.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.60	Depositor
% Data completeness (in resolution range)	99.9 (50.00-2.60)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	0.28	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.266 , 0.298	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7543	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	D	0.51	0/990	0.80	1/1549 (0.1%)
1	I	0.55	0/1015	0.87	2/1588 (0.1%)
2	E	0.53	0/1040	0.93	4/1619 (0.2%)
2	J	0.53	0/1040	0.95	8/1619 (0.5%)
3	A	0.67	0/838	1.16	6/1129 (0.5%)
3	F	0.75	0/838	1.21	9/1129 (0.8%)
4	B	0.73	0/748	1.05	2/996 (0.2%)
4	G	0.70	0/748	1.05	5/996 (0.5%)
5	C	0.62	0/381	1.28	7/506 (1.4%)
5	H	0.71	0/396	1.39	9/525 (1.7%)
All	All	0.62	0/8034	1.03	53/11656 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2
1	I	0	3
2	J	0	2
All	All	0	7

There are no bond length outliers.

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	6	VAL	N-CA-C	11.45	121.92	110.82
5	H	65	ILE	N-CA-C	-10.55	100.61	110.53
3	F	25	ILE	N-CA-C	-8.87	102.19	110.53
5	H	64	ARG	CA-C-N	8.40	131.31	120.56
5	H	64	ARG	C-N-CA	8.40	131.31	120.56

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	652	U	Sidechain
1	D	674	G	Sidechain
1	I	659	U	Sidechain
1	I	672	U	Sidechain
1	I	674	G	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	880	0	438	30	0
1	I	902	0	449	40	0
2	E	932	0	479	36	0
2	J	932	0	479	40	0
3	A	825	0	841	96	0
3	F	825	0	841	117	0
4	B	737	0	772	65	0
4	G	737	0	772	64	0
5	C	379	0	421	56	0
5	H	394	0	437	60	0
All	All	7543	0	5929	549	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 41.

The worst 5 of 549 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:652:U:O2'	1:I:653:A:H5''	1.37	1.21
1:D:659:U:H5''	3:F:36:ARG:HH22	1.11	1.10
4:G:4:LYS:HD3	4:G:5:GLU:H	1.31	0.96
5:H:53:ARG:HA	5:H:56:THR:HG23	1.47	0.95
1:D:659:U:H5''	3:F:36:ARG:NH2	1.82	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	96/98 (98%)	74 (77%)	18 (19%)	4 (4%)	2	3
3	F	96/98 (98%)	68 (71%)	17 (18%)	11 (12%)	0	0
4	B	86/88 (98%)	75 (87%)	9 (10%)	2 (2%)	5	9
4	G	86/88 (98%)	71 (83%)	11 (13%)	4 (5%)	2	2
5	C	40/88 (46%)	26 (65%)	9 (22%)	5 (12%)	0	0
5	H	42/88 (48%)	31 (74%)	10 (24%)	1 (2%)	4	9
All	All	446/548 (81%)	345 (77%)	74 (17%)	27 (6%)	1	1

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	C	63	GLN
3	F	12	PRO
3	F	43	LEU
3	F	47	ARG
3	F	93	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	89/89 (100%)	79 (89%)	10 (11%)	6	12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	F	89/89 (100%)	73 (82%)	16 (18%)	2	3
4	B	80/80 (100%)	67 (84%)	13 (16%)	2	4
4	G	80/80 (100%)	64 (80%)	16 (20%)	1	2
5	C	41/77 (53%)	36 (88%)	5 (12%)	5	10
5	H	43/77 (56%)	41 (95%)	2 (5%)	23	48
All	All	422/492 (86%)	360 (85%)	62 (15%)	3	6

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	2	ARG
4	G	57	MET
3	F	40	VAL
4	G	56	LEU
4	G	87	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	94	GLN
4	G	45	HIS
5	H	36	ASN
4	G	70	GLN
4	B	12	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	D	39/41 (95%)	5 (12%)	2 (5%)
1	I	40/41 (97%)	9 (22%)	2 (5%)
2	E	43/44 (97%)	13 (30%)	6 (13%)
2	J	43/44 (97%)	12 (27%)	4 (9%)
All	All	165/170 (97%)	39 (23%)	14 (8%)

5 of 39 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	D	583	A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	652	U
1	D	653	A
1	D	654	G
1	D	665	A

5 of 14 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	E	753	A
1	I	652	U
2	J	753	A
2	J	733	A
2	J	748	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.