



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 04:54 AM UTC

PDB ID : 9G3Z / pdb_00009g3z
EMDB ID : EMD-51019
Title : Structure of the Open gamma-Tubulin Ring Complex from Pig Brain
Authors : Munoz-Hernandez, H.; Wiczorek, M.
Deposited on : 2024-07-12
Resolution : 4.30 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Q-score	-	25397	4585 (3.80 - 4.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	O	79	 62% 85% 15%
1	P	79	 81% 15%
1	Q	79	 80% 19%
2	Y	155	 33% 67%
3	a	451	 94%





Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	b	451	94%
3	c	451	92% 5%
3	d	451	91% 6%
3	e	451	93%
3	f	451	94%
3	g	451	91% 5%
3	h	451	95%
3	i	451	95%
3	j	451	93%
3	k	451	94%
3	l	451	15% 94%
3	m	451	95% 97%
3	n	451	92% 97%
4	w	1663	98%
4	x	1663	98%
5	B	910	67% 32%
5	D	910	68% 32%
5	F	910	68% 32%
5	H	910	80% 19%
5	N	910	38% 76% 22%
6	A	905	72% 26%
6	C	905	70% 26%
6	E	905	73% 26%
6	G	905	73% 26%
6	M	905	14% 85% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	L	1715	
8	I	667	
8	K	667	
9	J	1042	

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 113079 atoms, of which 34747 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitotic spindle organizing protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	H	N	O		
1	O	67	497	199	164	67	67	0	0
1	P	67	497	199	164	67	67	0	0
1	Q	64	464	190	146	64	64	0	0

- Molecule 2 is a protein called Mitotic-spindle organizing protein 2A isoform X4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	H	N	O		
2	Y	51	379	150	127	51	51	0	0

- Molecule 3 is a protein called Tubulin gamma chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	H	N	O		
3	a	444	3161	1306	967	444	444	0	0
3	m	444	3161	1306	967	444	444	0	0
3	b	442	3146	1299	963	442	442	0	0
3	c	440	3129	1293	956	440	440	0	0
3	d	426	3030	1252	926	426	426	0	0
3	e	432	3074	1270	940	432	432	0	0
3	f	433	3081	1273	942	433	433	0	0
3	g	433	3081	1273	942	433	433	0	0
3	h	439	3122	1291	953	439	439	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
3	i	438	Total	C	H	N	O	0	0
			3115	1288	951	438	438		
3	n	442	Total	C	H	N	O	0	0
			3146	1299	963	442	442		
3	k	438	Total	C	H	N	O	0	0
			3115	1288	951	438	438		
3	j	439	Total	C	H	N	O	0	0
			3122	1292	952	439	439		
3	l	435	Total	C	H	N	O	0	0
			3094	1279	945	435	435		

- Molecule 4 is a protein called CDK5 regulatory subunit-associated protein 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	w	33	Total	C	H	N	O	0	0
			230	99	65	33	33		
4	x	34	Total	C	H	N	O	0	0
			237	102	67	34	34		

- Molecule 5 is a protein called Gamma-tubulin complex component 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	N	707	Total	C	H	N	O	0	0
			5082	2092	1576	707	707		
5	B	621	Total	C	H	N	O	0	0
			4466	1839	1385	621	621		
5	H	739	Total	C	H	N	O	0	0
			5322	2189	1655	739	739		
5	D	621	Total	C	H	N	O	0	0
			4466	1839	1385	621	621		
5	F	621	Total	C	H	N	O	0	0
			4466	1839	1385	621	621		

- Molecule 6 is a protein called Gamma-tubulin complex component.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	M	782	Total	C	H	N	O	0	0
			5606	2316	1726	782	782		
6	A	669	Total	C	H	N	O	0	0
			4792	1983	1471	669	669		
6	C	669	Total	C	H	N	O	0	0
			4792	1983	1471	669	669		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
6	E	669	Total	C	H	N	O	0	0
			4792	1983	1471	669	669		
6	G	669	Total	C	H	N	O	0	0
			4792	1983	1471	669	669		

- Molecule 7 is a protein called Tubulin gamma complex associated protein 6.


Mol	Chain	Residues	Atoms					AltConf	Trace
7	L	773	Total	C	H	N	O	0	0
			5548	2269	1733	773	773		

- Molecule 8 is a protein called Gamma-tubulin complex component.


Mol	Chain	Residues	Atoms					AltConf	Trace
8	I	587	Total	C	H	N	O	0	0
			4180	1732	1274	587	587		
8	K	587	Total	C	H	N	O	0	0
			4180	1732	1274	587	587		

- Molecule 9 is a protein called Gamma-tubulin complex component.


Mol	Chain	Residues	Atoms					AltConf	Trace
9	J	664	Total	C	H	N	O	0	0
			4714	1967	1419	664	664		




• Molecule 3: Tubulin gamma chain




• Molecule 3: Tubulin gamma chain




• Molecule 3: Tubulin gamma chain





• Molecule 3: Tubulin gamma chain





• Molecule 3: Tubulin gamma chain




• Molecule 3: Tubulin gamma chain



• Molecule 3: Tubulin gamma chain



• Molecule 3: Tubulin gamma chain





• Molecule 3: Tubulin gamma chain








• Molecule 3: Tubulin gamma chain




• Molecule 3: Tubulin gamma chain




• Molecule 3: Tubulin gamma chain








• Molecule 5: Gamma-tubulin complex component 3




• Molecule 5: Gamma-tubulin complex component 3





• Molecule 5: Gamma-tubulin complex component 3







• Molecule 5: Gamma-tubulin complex component 3



• Molecule 6: Gamma-tubulin complex component



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	80608	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	55	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	2900	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.846	Depositor
Minimum map value	-0.251	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.034	Depositor
Recommended contour level	0.125	Depositor
Map size (\AA)	542.7187, 542.7187, 542.7187	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.41333, 1.41333, 1.41333	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	O	0.09	0/332	0.29	0/462
1	P	0.25	0/332	0.55	0/462
1	Q	0.22	0/317	0.45	0/441
2	Y	0.16	0/251	0.35	0/348
3	a	0.15	0/2193	0.46	0/3053
3	b	0.15	0/2181	0.42	0/3034
3	c	0.17	0/2171	0.43	0/3020
3	d	0.16	0/2101	0.44	0/2921
3	e	0.16	0/2132	0.43	0/2966
3	f	0.17	0/2137	0.42	0/2973
3	g	0.18	0/2137	0.41	0/2973
3	h	0.19	0/2167	0.40	0/3015
3	i	0.19	0/2162	0.44	0/3008
3	j	0.19	0/2168	0.47	0/3017
3	k	0.15	0/2162	0.40	0/3008
3	l	0.13	0/2147	0.37	0/2987
3	m	0.11	0/2193	0.35	0/3053
3	n	0.13	0/2181	0.41	0/3034
4	w	0.12	0/164	0.31	0/228
4	x	0.12	0/169	0.28	0/235
5	B	0.19	0/3077	0.43	0/4287
5	D	0.21	0/3077	0.44	0/4287
5	F	0.23	0/3077	0.45	0/4287
5	H	0.24	0/3662	0.46	0/5102
5	N	0.13	0/3502	0.41	3/4879 (0.1%)
6	A	0.18	0/3316	0.46	0/4620
6	C	0.21	0/3316	0.45	0/4620
6	E	0.23	0/3316	0.46	0/4620
6	G	0.24	0/3316	0.44	0/4620
6	M	0.12	0/3875	0.37	1/5399 (0.0%)
7	L	0.19	0/3805	0.52	0/5281
8	I	0.24	0/2901	0.44	0/4036
8	K	0.19	0/2901	0.40	0/4036
9	J	0.23	0/3288	0.45	0/4577

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
All	All	0.19	0/78226	0.43	4/108889 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	a	0	1
3	b	0	1
3	j	0	1
3	m	0	1
3	n	0	3
5	N	0	1
All	All	0	8

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	N	576	PHE	N-CA-C	7.81	127.43	110.80
5	N	574	GLY	N-CA-C	7.06	129.91	113.18
6	M	231	ARG	N-CA-C	-6.96	105.41	114.04
5	N	575	ASP	N-CA-C	6.35	118.08	111.03

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
5	N	575	ASP	Peptide
3	a	284	SER	Peptide
3	b	284	SER	Peptide
3	m	284	SER	Peptide
3	n	261	ILE	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	333	164	164	0	0
1	P	333	164	164	1	0
1	Q	318	146	146	1	0
2	Y	252	127	129	0	0
3	a	2194	967	967	9	0
3	b	2183	963	963	10	0
3	c	2173	956	956	13	0
3	d	2104	926	926	7	0
3	e	2134	940	940	6	0
3	f	2139	942	942	5	0
3	g	2139	942	942	12	0
3	h	2169	953	953	6	0
3	i	2164	951	951	5	0
3	j	2170	952	952	11	0
3	k	2164	951	951	9	0
3	l	2149	945	945	6	0
3	m	2194	967	967	2	0
3	n	2183	963	963	2	0
4	w	165	65	65	0	0
4	x	170	67	67	0	0
5	B	3081	1385	1385	9	0
5	D	3081	1385	1385	3	0
5	F	3081	1385	1385	3	0
5	H	3667	1655	1655	7	0
5	N	3506	1576	1576	5	0
6	A	3321	1471	1471	10	0
6	C	3321	1471	1471	17	0
6	E	3321	1471	1471	7	0
6	G	3321	1471	1471	7	0
6	M	3880	1726	1726	7	0
7	L	3815	1733	1733	30	0
8	I	2906	1274	1274	7	0
8	K	2906	1274	1274	17	0
9	J	3295	1419	1419	12	0
All	All	78332	34747	34749	237	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 237 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:L:513:CYS:O	7:L:1363:ASP:N	2.06	0.88
6:C:396:TYR:O	6:C:469:CYS:N	2.08	0.87
6:A:236:GLN:N	6:A:245:THR:O	2.16	0.79
8:K:209:PRO:O	8:K:255:LYS:N	2.15	0.79
3:i:318:ILE:N	3:i:380:ASN:O	2.15	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	65/79 (82%)	65 (100%)	0	0	100	100
1	P	65/79 (82%)	63 (97%)	1 (2%)	1 (2%)	8	38
1	Q	62/79 (78%)	59 (95%)	3 (5%)	0	100	100
2	Y	49/155 (32%)	49 (100%)	0	0	100	100
3	a	442/451 (98%)	431 (98%)	10 (2%)	1 (0%)	43	77
3	b	438/451 (97%)	429 (98%)	8 (2%)	1 (0%)	43	77
3	c	436/451 (97%)	429 (98%)	7 (2%)	0	100	100
3	d	420/451 (93%)	414 (99%)	6 (1%)	0	100	100
3	e	428/451 (95%)	422 (99%)	6 (1%)	0	100	100
3	f	429/451 (95%)	421 (98%)	8 (2%)	0	100	100
3	g	429/451 (95%)	421 (98%)	8 (2%)	0	100	100
3	h	435/451 (96%)	427 (98%)	8 (2%)	0	100	100
3	i	434/451 (96%)	426 (98%)	8 (2%)	0	100	100
3	j	435/451 (96%)	422 (97%)	12 (3%)	1 (0%)	43	77
3	k	434/451 (96%)	426 (98%)	8 (2%)	0	100	100
3	l	431/451 (96%)	423 (98%)	8 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	m	442/451 (98%)	431 (98%)	10 (2%)	1 (0%)	43	77
3	n	438/451 (97%)	425 (97%)	13 (3%)	0	100	100
4	w	31/1663 (2%)	31 (100%)	0	0	100	100
4	x	32/1663 (2%)	32 (100%)	0	0	100	100
5	B	613/910 (67%)	608 (99%)	5 (1%)	0	100	100
5	D	613/910 (67%)	608 (99%)	5 (1%)	0	100	100
5	F	613/910 (67%)	607 (99%)	6 (1%)	0	100	100
5	H	729/910 (80%)	718 (98%)	11 (2%)	0	100	100
5	N	699/910 (77%)	682 (98%)	16 (2%)	1 (0%)	48	83
6	A	659/905 (73%)	644 (98%)	14 (2%)	1 (0%)	43	77
6	C	659/905 (73%)	646 (98%)	12 (2%)	1 (0%)	43	77
6	E	659/905 (73%)	644 (98%)	14 (2%)	1 (0%)	43	77
6	G	659/905 (73%)	645 (98%)	13 (2%)	1 (0%)	43	77
6	M	772/905 (85%)	743 (96%)	27 (4%)	2 (0%)	36	71
7	L	753/1715 (44%)	677 (90%)	72 (10%)	4 (0%)	24	62
8	I	577/667 (86%)	567 (98%)	9 (2%)	1 (0%)	43	77
8	K	577/667 (86%)	568 (98%)	8 (1%)	1 (0%)	43	77
9	J	650/1042 (62%)	636 (98%)	14 (2%)	0	100	100
All	All	15607/23198 (67%)	15239 (98%)	350 (2%)	18 (0%)	49	83

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
6	A	321	LEU
6	C	321	LEU
6	E	321	LEU
6	G	321	LEU
5	N	574	GLY

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
6	G	1
6	A	1
6	C	1
6	E	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	G	850:ASP	C	851:CYS	N	3.27
1	A	850:ASP	C	851:CYS	N	3.01
1	C	850:ASP	C	851:CYS	N	2.91
1	E	850:ASP	C	851:CYS	N	2.88


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-51019. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192



Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 187




Y Index: 235




Z Index: 143


6.3.2 Raw map



X Index: 187



Y Index: 235




Z Index: 143


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map




6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.125. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 2585 nm³; this corresponds to an approximate mass of 2335 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.233 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.233\AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.30	-	-
Author-provided FSC curve	4.34	7.13	4.49
Unmasked-calculated*	7.99	16.18	8.23

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.99 differs from the reported value 4.3 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-51019 and PDB model 9G3Z. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.125 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.125).







































































9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.125) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9000	 0.2050
A	 0.9730	 0.1930
B	 0.9930	 0.2460
C	 0.9750	 0.2560
D	 0.9950	 0.2660
E	 0.9970	 0.2850
F	 0.9980	 0.2920
G	 0.9970	 0.3080
H	 0.9970	 0.3280
I	 0.9980	 0.3210
J	 0.9930	 0.2710
K	 0.9990	 0.2400
L	 0.9780	 0.1730
M	 0.8300	 0.0830
N	 0.5200	 0.0780
O	 0.2550	 0.1000
P	 0.9880	 0.3270
Q	 0.9810	 0.2500
Y	 0.9880	 0.1320
a	 0.9900	 0.1040
b	 0.9840	 0.1460
c	 0.9720	 0.1720
d	 0.9900	 0.1550
e	 0.9870	 0.1940
f	 0.9910	 0.2120
g	 0.9950	 0.2450
h	 0.9980	 0.3020
i	 0.9990	 0.2860
j	 0.9910	 0.1900
k	 0.9850	 0.1170
l	 0.8240	 0.0550
m	 0.0430	 0.0450
n	 0.0710	 0.0320
w	 0.8610	 -0.0210
x	 0.9770	 0.0520

