



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2025 – 06:28 am GMT

PDB ID : 9GBL  
Title : Human Angiotensin-1 converting enzyme C-domain in complex with a diprolyl inhibitor- SG18  
Authors : Gregory, K.S.; Acharya, K.R.; Cozier, G.E.  
Deposited on : 2024-07-31  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	597	 2% 76% 17%
2	B	4	 100%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4939 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Angiotensin-converting enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	575	4696	3011	802	858	25	0	2	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	64	GLY	GLU	engineered mutation	UNP P12821
A	90	GLN	ASN	engineered mutation	UNP P12821
A	155	GLN	ASN	engineered mutation	UNP P12821
A	337	GLN	ASN	engineered mutation	UNP P12821
A	586	GLN	ASN	engineered mutation	UNP P12821

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	4	49	28	2	19	0	0	0


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Cl 2 2	0	0

- Molecule 5 is (2S)-1-[(2S)-2-[[[(1S)-1-[(2S)-1-(4-carboxyphenyl)carbonylpyrrolidin-2-yl]-2-oxidanyl-2-oxidanylidene-ethyl]amino]propanoyl]pyrrolidine-2-carboxylic acid (three-letter code: A1IJY) (formula: C<sub>22</sub>H<sub>27</sub>N<sub>3</sub>O<sub>8</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 33 22 3 8	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	158	Total O 158 158	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Angiotensin-converting enzyme



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.82Å 85.68Å 135.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.47 – 2.40 72.36 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (72.47-2.40) 99.9 (72.36-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.04 (at 2.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.178 , 0.249 0.179 , 0.252	Depositor DCC
$R_{free}$ test set	1320 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.7	Xtrriage
Anisotropy	0.134	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 28.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4939	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, A1IJY, NAG, CL, ZN, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	2/4831 (0.0%)	1.33	43/6564 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	479	GLU	CD-OE1	5.13	1.31	1.25
1	A	176	GLU	CD-OE1	5.07	1.31	1.25

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	191	ARG	CA-CB-CG	12.97	141.94	113.40
1	A	53	ARG	CB-CA-C	-9.83	90.74	110.40
1	A	253	ARG	NE-CZ-NH2	-9.54	115.53	120.30
1	A	337	GLN	CB-CA-C	-8.61	93.18	110.40
1	A	169	MET	CG-SD-CE	8.58	113.92	100.20
1	A	142	MET	CG-SD-CE	-8.00	87.39	100.20
1	A	480	ASN	CB-CA-C	-7.88	94.64	110.40
1	A	191	ARG	CG-CD-NE	-7.48	96.09	111.80
1	A	561	ARG	NE-CZ-NH1	7.13	123.86	120.30
1	A	150	THR	OG1-CB-CG2	-6.85	94.25	110.00
1	A	614	LEU	CB-CG-CD2	6.81	122.58	111.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	139	LEU	CB-CG-CD1	-6.67	99.66	111.00
1	A	337	GLN	N-CA-CB	6.58	122.44	110.60
1	A	257	ARG	CD-NE-CZ	6.56	132.78	123.60
1	A	395	LYS	CD-CE-NZ	-6.32	97.17	111.70
1	A	448	MET	CG-SD-CE	6.26	110.22	100.20
1	A	561	ARG	NE-CZ-NH2	-6.19	117.20	120.30
1	A	493	GLN	N-CA-CB	6.08	121.55	110.60
1	A	53	ARG	NE-CZ-NH2	-5.94	117.33	120.30
1	A	350	VAL	N-CA-CB	5.88	124.45	111.50
1	A	318	GLU	N-CA-CB	5.87	121.17	110.60
1	A	585	PRO	N-CA-CB	-5.86	96.15	102.60
1	A	86	MET	CG-SD-CE	5.86	109.58	100.20
1	A	191	ARG	NE-CZ-NH2	5.77	123.19	120.30
1	A	66	ASN	CB-CA-C	5.77	121.93	110.40
1	A	466	GLN	N-CA-CB	5.67	120.81	110.60
1	A	457	PHE	N-CA-CB	-5.66	100.41	110.60
1	A	507	ASP	CB-CA-C	5.62	121.63	110.40
1	A	567	LYS	CB-CG-CD	5.54	125.99	111.60
1	A	468	ARG	N-CA-CB	5.51	120.52	110.60
1	A	561	ARG	CD-NE-CZ	5.51	131.31	123.60
1	A	81	LEU	CB-CG-CD2	5.40	120.17	111.00
1	A	431	ASN	CB-CA-C	5.30	120.99	110.40
1	A	609	LEU	N-CA-CB	-5.24	99.91	110.40
1	A	245	LEU	CB-CG-CD2	5.21	119.86	111.00
1	A	163	PRO	N-CA-CB	-5.21	96.87	102.60
1	A	48	VAL	N-CA-CB	5.20	122.93	111.50
1	A	308	GLN	CB-CA-C	-5.19	100.02	110.40
1	A	463	LEU	CB-CG-CD2	-5.17	102.22	111.00
1	A	547	LEU	CB-CG-CD1	-5.11	102.32	111.00
1	A	576	GLU	N-CA-CB	5.11	119.79	110.60
1	A	196	PHE	CB-CA-C	-5.09	100.21	110.40
1	A	87	GLN	CB-CA-C	5.06	120.53	110.40

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	191	ARG	Sidechain
1	A	402	ARG	Sidechain
1	A	468	ARG	Sidechain
1	A	489	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4696	0	4544	42	0
2	B	49	0	43	0	0
3	A	1	0	0	0	0
4	A	2	0	0	0	0
5	A	33	0	0	1	0
6	A	158	0	0	0	0
All	All	4939	0	4587	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:ASP:OD2	1:A:106:GLN:HG2	1.81	0.81
1:A:399:VAL:HG12	1:A:402:ARG:NH1	1.97	0.79
1:A:241:GLN:HB2	1:A:242:PRO:HD3	1.76	0.66
1:A:46:LYS:O	1:A:50:GLU:HG3	1.96	0.65
1:A:514:ILE:HB	1:A:515:PRO:HD3	1.84	0.59
1:A:76:GLU:N	1:A:76:GLU:OE1	2.38	0.55
1:A:470:ARG:HB3	1:A:476:ILE:HG13	1.89	0.54
1:A:533:PHE:O	1:A:537:LEU:HG	2.09	0.52
1:A:485:TRP:CD2	1:A:508:PRO:HG3	2.48	0.48
1:A:543:HIS:CD2	1:A:550:CYS:HB2	2.49	0.47
1:A:395:LYS:O	1:A:402:ARG:NH2	2.47	0.46
1:A:579:GLN:HA	1:A:584:GLN:O	2.15	0.46
1:A:476:ILE:HG23	1:A:480:ASN:HB3	1.97	0.46
1:A:159:LEU:HD12	1:A:159:LEU:N	2.31	0.46
1:A:488:LEU:HD22	1:A:492:TYR:HE2	1.80	0.46
1:A:300:ASP:HB3	1:A:303:GLU:HB2	1.99	0.45
1:A:358:ASP:HB2	1:A:391:PHE:CD2	2.51	0.44
1:A:457:PHE:CE2	1:A:461:SER:HB3	2.52	0.44
1:A:275:LEU:HA	1:A:286:ILE:HD13	1.99	0.44
1:A:325:SER:O	1:A:554:GLN:HA	2.18	0.44
1:A:241:GLN:CB	1:A:242:PRO:HD3	2.46	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:THR:HG21	1:A:375:LEU:HD13	2.00	0.43
1:A:162:GLU:HA	1:A:163:PRO:HA	1.84	0.43
1:A:565:ALA:HB2	1:A:581:ILE:HG13	1.99	0.43
1:A:477:THR:O	1:A:480:ASN:HB2	2.18	0.43
1:A:273:HIS:CE1	1:A:498:PRO:HB3	2.53	0.43
1:A:460:PHE:HB2	1:A:493:GLN:NE2	2.34	0.42
1:A:144:THR:O	1:A:145:THR:C	2.57	0.42
1:A:304:ALA:O	1:A:308:GLN:HG2	2.20	0.42
1:A:562:LEU:HD23	1:A:562:LEU:HA	1.93	0.42
1:A:335:PHE:CD1	1:A:335:PHE:C	2.92	0.42
1:A:468:ARG:NH2	1:A:518:VAL:O	2.51	0.42
1:A:593:LEU:HD23	1:A:593:LEU:HA	1.82	0.42
1:A:134:GLU:OE2	1:A:199:LYS:HE2	2.21	0.41
1:A:433:LEU:HD12	1:A:434:SER:H	1.86	0.41
1:A:473:ASP:OD1	1:A:475:SER:OG	2.33	0.41
1:A:355:SER:HA	5:A:704:A1IJY:O21	2.20	0.41
1:A:493:GLN:HB3	1:A:495:LEU:HG	2.03	0.41
1:A:67:TRP:CZ3	1:A:368:LYS:HB2	2.56	0.40
1:A:514:ILE:HB	1:A:515:PRO:CD	2.49	0.40
1:A:154:PRO:O	1:A:156:GLY:N	2.54	0.40
1:A:511:LYS:O	1:A:515:PRO:HD2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	571/597 (96%)	547 (96%)	22 (4%)	2 (0%)	30 44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	ASN
1	A	347	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	503/520 (97%)	482 (96%)	21 (4%)	25 43

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	SER
1	A	78	SER
1	A	79	LYS
1	A	83	GLN
1	A	86	MET
1	A	100	ARG
1	A	140	LEU
1	A	242	PRO
1	A	290	VAL
1	A	322	PHE
1	A	325	SER
1	A	346	ASP
1	A	350	VAL
1	A	371	THR
1	A	390	TYR
1	A	394	TYR
1	A	440	ASP
1	A	475	SER
1	A	500	PRO
1	A	585	PRO
1	A	594	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2,1	14,14,15	0.54	0	17,19,21	1.58	3 (17%)
2	NAG	B	2	2	14,14,15	0.44	0	17,19,21	1.46	4 (23%)
2	BMA	B	3	2	11,11,12	1.04	1 (9%)	15,15,17	0.98	1 (6%)
2	FUC	B	4	2	10,10,11	1.12	1 (10%)	14,14,16	2.14	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1
2	FUC	B	4	2	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4	FUC	C2-C3	-3.00	1.48	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	BMA	C4-C5	2.16	1.57	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	FUC	O3-C3-C2	-5.34	99.78	109.99
2	B	1	NAG	C1-C2-N2	3.52	116.50	110.49
2	B	1	NAG	C2-N2-C7	-3.33	118.17	122.90
2	B	4	FUC	C1-C2-C3	3.00	113.35	109.67
2	B	4	FUC	O2-C2-C3	-2.94	104.24	110.14
2	B	2	NAG	C2-N2-C7	2.79	126.87	122.90
2	B	2	NAG	O4-C4-C3	-2.76	103.96	110.35
2	B	2	NAG	O3-C3-C2	2.73	115.12	109.47
2	B	4	FUC	C3-C4-C5	2.41	113.52	109.77
2	B	4	FUC	O2-C2-C1	2.40	114.07	109.15
2	B	3	BMA	C1-O5-C5	2.24	115.22	112.19
2	B	1	NAG	O5-C5-C6	2.20	110.66	107.20
2	B	2	NAG	C1-C2-N2	-2.15	106.82	110.49

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
2	B	3	BMA	O5-C5-C6-O6
2	B	3	BMA	C4-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	2	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	A1IJY	A	704	3	35,35,35	3.27	18 (51%)	43,50,50	2.44	17 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	A1IJY	A	704	3	-	7/36/56/56	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	704	A1IJY	C06-N10	-11.60	1.32	1.47
5	A	704	A1IJY	C09-N10	6.29	1.59	1.47
5	A	704	A1IJY	C27-N26	5.12	1.57	1.47
5	A	704	A1IJY	C05-N04	-4.89	1.39	1.46
5	A	704	A1IJY	C11-N10	4.56	1.45	1.34
5	A	704	A1IJY	C13-C12	3.42	1.45	1.39
5	A	704	A1IJY	C30-N26	-3.38	1.40	1.47
5	A	704	A1IJY	C14-C15	3.15	1.44	1.39
5	A	704	A1IJY	C02-N26	3.08	1.41	1.34
5	A	704	A1IJY	C07-C06	3.02	1.58	1.53
5	A	704	A1IJY	C08-C09	-2.99	1.41	1.51
5	A	704	A1IJY	C15-C16	2.87	1.55	1.49
5	A	704	A1IJY	C12-C11	2.86	1.54	1.50
5	A	704	A1IJY	C05-C06	2.76	1.57	1.54
5	A	704	A1IJY	C28-C27	-2.58	1.42	1.51
5	A	704	A1IJY	C29-C30	2.14	1.59	1.53
5	A	704	A1IJY	C19-C15	2.13	1.42	1.39
5	A	704	A1IJY	C08-C07	2.04	1.60	1.51

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	704	A1IJY	O32-C31-O33	-6.68	108.93	124.09
5	A	704	A1IJY	O18-C16-C15	5.45	129.00	114.85
5	A	704	A1IJY	C19-C15-C16	-4.76	111.03	120.39
5	A	704	A1IJY	C13-C14-C15	-4.54	115.50	120.78
5	A	704	A1IJY	C29-C30-C31	-4.27	104.35	111.36
5	A	704	A1IJY	O32-C31-C30	4.19	127.91	113.38
5	A	704	A1IJY	C25-C03-C02	-4.13	102.24	109.73
5	A	704	A1IJY	O18-C16-O17	-3.18	116.29	123.35
5	A	704	A1IJY	C19-C15-C14	3.16	123.09	118.59
5	A	704	A1IJY	C14-C15-C16	2.75	125.79	120.39
5	A	704	A1IJY	O23-C22-O24	-2.74	117.87	124.09
5	A	704	A1IJY	O17-C16-C15	-2.68	114.30	121.45

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	704	A1IJY	C27-N26-C02	-2.60	116.83	125.88
5	A	704	A1IJY	C20-C12-C13	2.46	122.08	118.59
5	A	704	A1IJY	C30-N26-C02	2.31	129.38	121.41
5	A	704	A1IJY	C25-C03-N04	-2.24	104.11	108.85
5	A	704	A1IJY	C19-C20-C12	-2.18	118.24	120.78

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	704	A1IJY	N04-C05-C06-C07
5	A	704	A1IJY	C22-C05-C06-C07
5	A	704	A1IJY	N04-C05-C22-O23
5	A	704	A1IJY	N04-C05-C22-O24
5	A	704	A1IJY	C22-C05-N04-C03
5	A	704	A1IJY	C22-C05-C06-N10
5	A	704	A1IJY	N04-C05-C06-N10

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	704	A1IJY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	575/597 (96%)	-0.43	11 (1%) 66 62	11, 32, 63, 107	2 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	347	GLY	3.9
1	A	617	PRO	3.7
1	A	40	ASP	3.4
1	A	344	PRO	2.7
1	A	297	PRO	2.6
1	A	154	PRO	2.5
1	A	155	GLN	2.5
1	A	82	LEU	2.4
1	A	123	GLU	2.1
1	A	79	LYS	2.1
1	A	346	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BMA	B	3	11/12	0.71	0.15	56,76,86,89	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	1	14/15	0.86	0.11	39,46,73,78	0
2	NAG	B	2	14/15	0.89	0.10	51,59,70,71	0
2	FUC	B	4	10/11	0.90	0.09	34,41,49,58	10

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	A1IJY	A	704	33/33	0.96	0.06	20,28,35,40	0
4	CL	A	703	1/1	0.99	0.02	24,24,24,24	0
4	CL	A	702	1/1	0.99	0.03	27,27,27,27	0
3	ZN	A	701	1/1	1.00	0.01	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.