



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 08:19 PM UTC

PDB ID : 2GVZ / pdb_00002gvz
Title : Crystal Structure of Complex of Gs- with The Catalytic Domains of Mammalian Adenylyl Cyclase: Complex with MANT-ATP and Mn
Authors : Mou, T.-C.; Sprang, S.R.
Deposited on : 2006-05-03
Resolution : 3.27 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1303 (3.30-3.26)
Clashscore	190562	1354 (3.30-3.26)
Ramachandran outliers	187476	1334 (3.30-3.26)
Sidechain outliers	187428	1333 (3.30-3.26)
RSRZ outliers	180081	1303 (3.30-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	 32% 40% 10% 16%
2	B	212	 39% 43% 7% 11%
3	C	394	 35% 43% 6% 16%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5751 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylate cyclase type 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	189	1476	929	259	271	17	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	MET	-	initiating methionine	UNP P30803
A	357	HIS	-	expression tag	UNP P30803
A	358	HIS	-	expression tag	UNP P30803
A	359	HIS	-	expression tag	UNP P30803
A	360	HIS	-	expression tag	UNP P30803
A	361	HIS	-	expression tag	UNP P30803
A	362	HIS	-	expression tag	UNP P30803
A	476	MET	VAL	engineered mutation	UNP P30803

- Molecule 2 is a protein called Adenylate cyclase type 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	188	1457	930	239	278	10	0	0	0

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(s), alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	330	2702	1714	470	505	13	0	0	0

- Molecule 4 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mn	0	0
			2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Mn			
4	C	1	1	1		0	0

- Molecule 5 is METHYLPIPERAZINOFORSKOLIN (CCD ID: FKP) (formula: $C_{30}H_{50}N_2O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	39	30	2	7	0	0

- Molecule 6 is 3'-O-[2-(METHYLAMINO)BENZOYL]ADENOSINE 5'-(TETRAHYDROGEN TRIPHOSPHATE) (CCD ID: ONA) (formula: $C_{18}H_{23}N_6O_{14}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
6	A	1	41	18	6	14	3	0	0

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
7	C	1	1	1	0	0

- Molecule 8 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (CCD ID: GSP) (formula: C₁₀H₁₆N₅O₁₃P₃S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
8	C	1	32	10	5	13	3	1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Adenylate cyclase type 5



• Molecule 2: Adenylate cyclase type 2



• Molecule 3: Guanine nucleotide-binding protein G(s), alpha subunit



T350	R283	M216	P138	Y65
A351	T284	M219	D139	ASN
S352	V287	H220	F140	GLY
D354	L288	M221	P143	PHE
G355	L289	F222	E144	ASN
H357	F290	D223	F145	GLY
Y358	L291	V224	F146	GLU
G359	N292	G225	Y147	GLY
Y360	K293	G226	E148	GLU
T364	Q294	Q227	H149	GLU
G365	D295	R228	L150	ASP
A366	L296	D229	A150	PRO
V367	L297	E230	L153	GLN
D368	V301	W154	E155	ALA
T369	G304	R231	R232	ALA
E370	K305	K233	E156	ALA
N371	S306	W234	D156	ARG
I372	R307	I235	E157	SER
R373	S308	Q236	G158	ASN
R374	I309	C237	Y159	SER
V375	D310	F238	R160	SER
F376	Y311	T242	A161	ASP
N377	F312	I244	C162	G86
D378	P313	I245	Y163	G87
C379	E314	F246	E164	K88
R380	F315	V247	R165	K89
D381	F318	V248	S166	T90
I382	Y319	A249	N167	R91
I383	T319	S250	E168	V92
Q384	T320	S251	Y169	I95
R385	P321	S252	Q170	N88
M386	A324	Y253	L171	L99
L388	T325	M254	I172	K100
ARG	P326	M255	D173	E101
GLN	E327	V256	C174	A102
TYR	P328	I257	A175	A103
GLU	G329	R258	Q176	I103
LEU	E330	E259	Y177	E104
LEU	E331	D260	F178	I105
	D331	N261	L179	I106
	F332	R265	D180	V107
	R333	L266	K181	A108
	V334	A269	I182	A109
	T335	L272	D183	M110
	R336	F275	V184	L113
	A337	K274	I185	L119
	F340	S275	K186	M124
	I341	I276	Q195	N127
	D342	W277	C200	R127
	E344	N278	R201	V128
	F345	N279	V202	V128
	L346	R280	L203	L131
	R347	W281	T204	I131
	I348	L282	S205	L132
	S349		G206	S133
			I207	V134
			V214	V137

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	116.80Å 132.10Å 69.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.97 – 3.27 14.97 – 3.27	Depositor EDS
% Data completeness (in resolution range)	85.8 (14.97-3.27) 75.1 (14.97-3.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 3.25Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.275 , 0.330 0.260 , 0.309	Depositor DCC
R_{free} test set	774 reflections (5.32%)	wwPDB-VP
Wilson B-factor (Å ²)	55.9	Xtrriage
Anisotropy	1.226	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 52.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5751	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, GSP, CL, ONA, FKP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/1504	1.12	12/2027 (0.6%)
2	B	0.70	0/1481	1.08	9/1999 (0.5%)
3	C	0.62	0/2759	1.10	18/3733 (0.5%)
All	All	0.63	0/5744	1.10	39/7759 (0.5%)

There are no bond length outliers.

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	388	ASP	N-CA-C	14.41	131.10	113.23
3	C	327	GLU	CA-C-N	14.34	137.77	119.84
3	C	327	GLU	C-N-CA	14.34	137.77	119.84
1	A	525	ARG	N-CA-C	-13.59	88.34	109.95
2	B	912	LEU	N-CA-C	-9.50	100.00	111.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1476	0	1450	132	0
2	B	1457	0	1463	102	0
3	C	2702	0	2650	187	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	2	0	0	0	0
4	C	1	0	0	0	0
5	A	39	0	50	6	0
6	A	41	0	19	2	0
7	C	1	0	0	1	0
8	C	32	0	12	1	0
All	All	5751	0	5644	397	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 397 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:925:LEU:HG	2:B:982:LYS:HZ2	1.23	1.00
3:C:231:ARG:HH21	3:C:272:LEU:HD22	1.26	0.98
3:C:63:LEU:HD13	3:C:373:ARG:HE	1.31	0.94
5:A:1:FKP:H173	5:A:1:FKP:H201	1.51	0.90
3:C:119:LEU:H	3:C:119:LEU:HD12	1.37	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	187/225 (83%)	153 (82%)	30 (16%)	4 (2%)	5 26
2	B	184/212 (87%)	156 (85%)	26 (14%)	2 (1%)	11 39
3	C	326/394 (83%)	270 (83%)	49 (15%)	7 (2%)	5 26
All	All	697/831 (84%)	579 (83%)	105 (15%)	13 (2%)	6 29

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	351	ALA
1	A	382	ILE
1	A	456	ALA
1	A	479	VAL
1	A	480	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/189 (84%)	141 (89%)	17 (11%)	6	24
2	B	161/184 (88%)	149 (92%)	12 (8%)	12	38
3	C	297/351 (85%)	277 (93%)	20 (7%)	15	42
All	All	616/724 (85%)	567 (92%)	49 (8%)	11	36

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	1077	THR
3	C	200	CYS
3	C	40	THR
3	C	119	LEU
3	C	227	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	93	GLN
3	C	213	GLN
3	C	176	GLN
3	C	220	HIS
2	B	965	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ONA	A	100	4	43,44,44	4.12	18 (41%)	63,67,67	2.62	26 (41%)
5	FKP	A	1	-	40,42,42	2.73	20 (50%)	52,68,68	2.74	21 (40%)
8	GSP	C	395	4	33,34,34	1.67	6 (18%)	47,54,54	1.52	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ONA	A	100	4	-	6/32/48/48	0/4/4/4
5	FKP	A	1	-	-	12/14/97/97	0/4/4/4
8	GSP	C	395	4	-	3/21/38/38	0/3/3/3

The worst 5 of 44 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	100	ONA	CA4-CA5	12.12	1.59	1.38
6	A	100	ONA	CA5-CA6	12.09	1.59	1.39
6	A	100	ONA	CA3-CA2	10.52	1.56	1.38
6	A	100	ONA	CA3-CA4	8.60	1.57	1.38
6	A	100	ONA	OA-CA	7.72	1.44	1.22

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1	FKP	C19-C4-C18	-10.98	91.89	107.87
6	A	100	ONA	N1-C2-N3	-7.02	117.95	128.58
6	A	100	ONA	O2B-PB-O1A	6.79	125.63	107.27
6	A	100	ONA	O2B-PB-O1B	6.58	125.05	107.27
5	A	1	FKP	O1-C8-C7	5.99	108.83	103.16

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1	FKP	C6-C7-O4-C21
5	A	1	FKP	C16-C13-C14-C15
5	A	1	FKP	C22-C21-O4-C7
5	A	1	FKP	O5-C21-O4-C7
5	A	1	FKP	C23-C24-C25-N1


There are no ring outliers.


3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	100	ONA	2	0
5	A	1	FKP	6	0
8	C	395	GSP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	189/225 (84%)	0.03	2 (1%) 78 61	30, 87, 99, 116	0
2	B	188/212 (88%)	-0.36	0 100 100	13, 49, 96, 98	0
3	C	330/394 (83%)	-0.34	0 100 100	19, 55, 93, 100	0
All	All	707/831 (85%)	-0.25	2 (0%) 90 82	13, 59, 97, 116	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	400	PHE	3.9
1	A	475	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	ONA	A	100	41/41	0.91	0.11	53,88,97,100	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MN	A	581	1/1	0.93	0.09	64,64,64,64	0
5	FKP	A	1	39/39	0.94	0.09	13,23,87,89	0
7	CL	C	397	1/1	0.95	0.35	59,59,59,59	0
8	GSP	C	395	32/32	0.95	0.08	31,41,73,76	0
4	MN	A	582	1/1	0.97	0.06	31,31,31,31	0
4	MN	C	396	1/1	0.99	0.04	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around FKP A 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GSP C 395:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.