

# wwPDB X-ray Structure Validation Summary Report (i)

Jun 10, 2025 – 04:19 pm BST

PDB ID	:	$9$ H2X / pdb_00009h2x
Title	:	Crystal structure of stabilized A2A adenosine receptor A2AR-StaR2-bRIL in
		complex with compound 7, a novel nanomolar A2A receptor antagonist from
		modern hit-finding with structure-guided de novo design
Authors	:	Tian, G.; Maja, N.
Deposited on		
Resolution	:	1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity		4-5-2 with Phenix2.0rc1
0		
Mogul	:	1.8.4, CSD as $541$ be (2020)
Xtriage (Phenix)	:	2.0rc1
$\mathrm{EDS}$	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072(1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			9%		
1	А	433	82%	7%	11%



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 3586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Adenosine receptor A2a,Soluble cytochrome b562.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	384	Total 3084	C 2014	N 517	O 530	S 23	0	15	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	-8	ASP	-	expression tag	UNP P29274
А	-7	TYR	_	expression tag	UNP P29274
А	-6	LYS	_	expression tag	UNP P29274
А	-5	ASP	-	expression tag	UNP P29274
А	-4	ASP	-	expression tag	UNP P29274
А	-3	ASP	-	expression tag	UNP P29274
А	-2	ASP	-	expression tag	UNP P29274
А	-1	GLY	-	expression tag	UNP P29274
А	0	ALA	-	expression tag	UNP P29274
А	1	PRO	-	expression tag	UNP P29274
А	54	LEU	ALA	conflict	UNP P29274
A	88	ALA	THR	conflict	UNP P29274
А	107	ALA	ARG	conflict	UNP P29274
А	122	ALA	LYS	conflict	UNP P29274
А	154	ALA	ASN	conflict	UNP P29274
А	202	ALA	LEU	conflict	UNP P29274
А	1007	TRP	MET	conflict	UNP P0ABE7
А	1102	ILE	HIS	conflict	UNP P0ABE7
А	1106	LEU	-	linker	UNP P0ABE7
A	235	ALA	LEU	conflict	UNP P29274
А	239	ALA	VAL	conflict	UNP P29274
А	277	ALA	SER	conflict	UNP P29274
А	318	ALA	-	expression tag	UNP P29274
А	319	HIS	-	expression tag	UNP P29274
А	320	HIS	-	expression tag	UNP P29274
А	321	HIS	-	expression tag	UNP P29274
А	322	HIS	-	expression tag	UNP P29274

There are 33 discrepancies between the modelled and reference sequences:




Chain	Residue	Modelled	Actual	Comment	Reference
А	323	HIS	-	expression tag	UNP P29274
А	324	HIS	-	expression tag	UNP P29274
А	325	HIS	-	expression tag	UNP P29274
А	326	HIS	-	expression tag	UNP P29274
А	327	HIS	-	expression tag	UNP P29274
А	328	HIS	-	expression tag	UNP P29274

Continued from previous page...

• Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Na 1 1	0	0


• Molecule 3 is CHOLESTEROL (CCD ID: CLR) (formula:  $C_{27}H_{46}O$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total         C         O           28         27         1	0	0
3	А	1	Total         C         O           28         27         1	0	0
3	А	1	Total         C         O           28         27         1	0	0

• Molecule 4 is OLEIC ACID (CCD ID: OLA) (formula:  $C_{18}H_{34}O_2$ ).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total         C         O           16         14         2	0	0
4	А	1	Total C 11 11	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
4	А	1	Total         C         O           11         9         2	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
4	А	1	Total         C         O           16         14         2	0	0
4	А	1	Total         C         O           20         18         2	0	0
4	А	1	Total         C         O           11         9         2	0	0
4	А	1	Total C 6 6	0	0
4	А	1	Total         C         O           20         18         2	0	0
4	А	1	Total         C         O           10         8         2	0	0
4	А	1	Total         C         O           20         18         2	0	0
4	А	1	Total         C         O           16         14         2	0	0
4	А	1	Total         C         O           18         16         2	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total C 18 18	0	0
4	А	1	Total         C         O           15         13         2	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 9 & 7 & 2 \end{array}$	0	0
4	А	1	Total         C         O           12         10         2	0	0
4	А	1	$\begin{array}{cc} \text{Total} & \text{C} \\ 6 & 6 \end{array}$	0	0
4	А	1	Total         C         O           17         15         2	0	0
4	А	1	Total         C         O           10         8         2	0	0

Continued from previous page...

• Molecule 5 is 4-(furan-2-yl)-6-(6-imidazol-1-ylpyridin-2-yl)-1,3,5-triazin-2-amine (CCD ID: A1IR1) (formula:  $C_{15}H_{11}N_7O$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	А	1	Total 23	C 15	N 7	0 1	0	0


• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	116	Total O 116 116	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Adenosine receptor A2a,Soluble cytochrome b562



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	39.35Å 179.53Å 140.12Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	25.20 - 1.75	Depositor
Resolution (A)	25.20 - 1.75	EDS
% Data completeness	86.3 (25.20-1.75)	Depositor
(in resolution range)	86.3 (25.20-1.75)	EDS
R <sub>merge</sub>	0.41	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.38 (at 1.75 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.11.8	Depositor
D D.	0.206 , $0.226$	Depositor
$R, R_{free}$	0.201 , $0.222$	DCC
$R_{free}$ test set	2200 reflections $(5.02%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	22.6	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35 , $51.6$	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3586	wwPDB-VP
Average B, all atoms $(Å^2)$	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CLR, NA, OLA, A1IR1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.76	0/3156	0.94	2/4295~(0.0%)	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	283	VAL	N-CA-C	5.98	118.56	111.09
1	А	24	ASN	CA-CB-CG	-5.48	107.12	112.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3084	0	3158	20	0
2	А	1	0	0	0	0
3	А	84	0	138	0	0
4	А	278	0	391	5	0
5	А	23	0	0	0	0
6	А	116	0	0	1	0
All	All	3586	0	3687	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29[A]:TRP:NE1	1:A:33:LEU:HD11	2.12	0.65
1:A:29[A]:TRP:HE1	1:A:33:LEU:HD11	1.70	0.56
1:A:1030:LEU:HB3	1:A:1076[B]:LEU:HG	1.88	0.54
1:A:40:VAL:HG11	1:A:116:VAL:CG1	2.40	0.52
1:A:29[A]:TRP:CD1	1:A:33:LEU:CD1	2.92	0.52

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	А	395/433~(91%)	393~(100%)	2(0%)	0	100 100	

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	324/353~(92%)	323 (100%)	1 (0%)	91 88		

All (1) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	138	THR

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such side chains are listed below:

Mol	Chain	Res	Type
1	А	157	GLN
1	А	1006	ASN
1	А	1088	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 26 ligands modelled in this entry, 1 is monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type	Cype Chain Res	Dec	Link	Bo	Bond lengths			Bond angles		
	Type		nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2	
4	OLA	А	1205	-	$15,\!15,\!19$	0.59	0	$15,\!15,\!19$	0.56	0	
4	OLA	А	1220	-	14,14,19	0.61	0	14,14,19	0.57	0	
4	OLA	А	1216	-	19,19,19	0.54	0	19,19,19	0.56	0	
4	OLA	А	1217	-	$15,\!15,\!19$	0.60	0	$15,\!15,\!19$	0.59	0	
3	CLR	А	1204	-	31,31,31	0.40	0	48,48,48	0.50	0	
4	OLA	А	1215	-	9,9,19	0.73	0	9,9,19	0.62	0	



Mol	Type	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
MOI	туре	Ullalli	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
4	OLA	A	1224	-	16,16,19	0.57	0	16, 16, 19	0.51	0
4	OLA	А	1218	-	17,17,19	0.59	0	$17,\!17,\!19$	0.60	0
4	OLA	A	1210	-	$15,\!15,\!19$	0.62	0	$15,\!15,\!19$	0.57	0
4	OLA	А	1213	-	$5,\!5,\!19$	0.17	0	$4,\!4,\!19$	0.18	0
4	OLA	А	1219	-	17,17,19	0.21	0	16, 16, 19	0.24	0
4	OLA	А	1222	-	10, 10, 19	0.65	0	$9,\!9,\!19$	0.75	0
4	OLA	А	1223	-	$5,\!5,\!19$	0.30	0	$4,\!4,\!19$	0.18	0
4	OLA	А	1225	-	$9,\!9,\!19$	0.69	0	$9,\!9,\!19$	0.67	0
4	OLA	A	1211	-	$19,\!19,\!19$	0.53	0	$19,\!19,\!19$	0.53	0
4	OLA	A	1212	-	10,10,19	0.64	0	$10,\!10,\!19$	0.67	0
5	A1IR1	А	1226	-	22,26,26	0.51	0	29,36,36	0.77	1 (3%)
4	OLA	А	1214	-	19,19,19	0.50	0	19,19,19	0.53	0
4	OLA	A	1208	-	10,10,19	0.67	0	$10,\!10,\!19$	0.66	0
4	OLA	A	1221	-	8,8,19	0.73	0	8,8,19	0.70	0
3	CLR	A	1202	-	31,31,31	0.42	0	48,48,48	0.51	0
4	OLA	А	1209	-	7,7,19	0.86	0	7,7,19	0.83	0
3	CLR	А	1203	-	31,31,31	0.36	0	48,48,48	0.53	0
4	OLA	А	1206	-	10,10,19	0.24	0	9,9,19	0.32	0
4	OLA	А	1207	-	7,7,19	0.79	0	$7,\!7,\!19$	0.79	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLA	А	1205	-	-	3/13/13/17	-
4	OLA	А	1220	-	-	3/12/12/17	-
4	OLA	А	1216	_	-	7/17/17/17	-
4	OLA	А	1217	_	-	3/13/13/17	-
3	CLR	А	1204	-	-	1/10/68/68	0/4/4/4
4	OLA	А	1215	-	-	2/7/7/17	-
4	OLA	А	1224	-	-	5/14/14/17	-
4	OLA	А	1218	-	-	9/15/15/17	-
4	OLA	А	1210	-	-	2/13/13/17	-
4	OLA	А	1213	-	-	1/3/3/17	-
4	OLA	А	1219	-	-	3/15/15/17	-
4	OLA	А	1222	-	-	3/6/6/17	-
4	OLA	А	1223	-	-	0/3/3/17	-
4	OLA	А	1225	-	-	0/7/7/17	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLA	А	1211	-	-	5/17/17/17	-
4	OLA	А	1212	-	-	3/8/8/17	-
5	A1IR1	А	1226	-	-	0/8/12/12	0/4/4/4
4	OLA	А	1214	-	-	3/17/17/17	-
4	OLA	А	1208	-	-	2/8/8/17	-
4	OLA	А	1221	-	-	2/6/6/17	-
3	CLR	А	1202	-	-	1/10/68/68	0/4/4/4
4	OLA	А	1209	-	-	2/5/5/17	-
3	CLR	А	1203	-	-	0/10/68/68	0/4/4/4
4	OLA	А	1206	-	-	0/8/8/17	-
4	OLA	А	1207	-	-	2/5/5/17	-

Continued from previous page...

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	А	1226	A1IR1	C13-C14-C15	-2.03	105.71	112.92

There are no chirality outliers.

5 of 62 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	1224	OLA	C1-C2-C3-C4
4	А	1222	OLA	C9-C10-C11-C12
4	А	1224	OLA	C2-C3-C4-C5
4	А	1218	OLA	C12-C13-C14-C15
4	А	1205	OLA	С11-С10-С9-С8

There are no ring outliers.


5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	А	1218	OLA	1	0
4	А	1214	OLA	1	0
4	А	1208	OLA	1	0
4	А	1221	OLA	1	0
4	А	1206	OLA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,




bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	384/433~(88%)	0.49	41 (10%) 12 14	7, 27, 51, 72	15 (3%)

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	1021	ASP	5.3
1	А	1061	PHE	5.0
1	А	290	TYR	4.1
1	А	1	PRO	3.7
1	А	1040	ALA	3.7

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
4	OLA	А	1207	8/20	0.50	0.26	$66,\!66,\!67,\!67$	0
4	OLA	А	1214	20/20	0.67	0.23	64,64,67,67	0




9H2X
------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	OLA	А	1213	6/20	0.69	0.22	42,42,43,43	0
4	OLA	А	1222	12/20	0.70	0.20	44,44,62,62	0
4	OLA	А	1215	10/20	0.72	0.25	64,64,64,64	0
4	OLA	А	1209	8/20	0.75	0.24	57,59,61,61	0
4	OLA	А	1217	16/20	0.76	0.21	64,64,65,65	0
4	OLA	А	1210	16/20	0.77	0.18	34,38,43,43	0
4	OLA	А	1208	11/20	0.77	0.22	68,69,69,69	0
4	OLA	А	1206	11/20	0.77	0.20	46,47,48,48	0
4	OLA	А	1212	11/20	0.78	0.18	$50,\!50,\!53,\!53$	0
4	OLA	А	1216	20/20	0.78	0.18	49,51,55,55	0
4	OLA	А	1223	6/20	0.78	0.26	52,53,53,53	0
4	OLA	А	1211	20/20	0.79	0.17	43,46,54,54	0
4	OLA	А	1220	15/20	0.80	0.16	44,47,54,54	0
4	OLA	А	1219	18/20	0.81	0.17	44,48,49,49	0
4	OLA	А	1225	10/20	0.81	0.15	51,52,54,54	0
4	OLA	А	1224	17/20	0.82	0.20	48,49,51,51	0
4	OLA	А	1221	9/20	0.82	0.19	47,48,51,52	0
4	OLA	А	1205	16/20	0.83	0.15	40,42,45,45	0
4	OLA	А	1218	18/20	0.85	0.16	46,47,50,50	0
3	CLR	А	1203	28/28	0.91	0.09	23,24,30,32	0
3	CLR	А	1202	28/28	0.93	0.07	20,23,32,33	0
3	CLR	А	1204	28/28	0.95	0.07	17,20,36,37	0
2	NA	А	1201	1/1	0.95	0.13	36,36,36,36	0
5	A1IR1	А	1226	23/23	0.97	0.05	13,14,23,25	0


Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






















































### 6.5 Other polymers (i)

There are no such residues in this entry.

