



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 06:36 AM UTC

PDB ID : 9H4R / pdb_00009h4r
Title : Structure of fertilization-blocking monoclonal antibody IE-3 VHVL bound to the ZP-N1 domain of mouse ZP2 (crystal form I)
Authors : Dioguardi, E.; De Sanctis, D.; Jovine, L.
Deposited on : 2024-10-21
Resolution : 1.53 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1003 (1.54-1.54)
Clashscore	190562	1025 (1.54-1.54)
Ramachandran outliers	187476	1007 (1.54-1.54)
Sidechain outliers	187428	1007 (1.54-1.54)
RSRZ outliers	180081	1002 (1.54-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	112	 78% 8% 14%
1	B	112	 82% 14%
2	H	123	 2% 91% 7%
2	X	123	 96%
3	L	117	 3% 90% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Y	117	 84% 15%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zona pellucida sperm-binding protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	96	776	497	123	150	6	0	0	0
1	B	96	775	498	123	148	6	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	SER	ASN	engineered mutation	UNP P20239
A	139	LEU	-	expression tag	UNP P20239
A	140	GLU	-	expression tag	UNP P20239
A	141	HIS	-	expression tag	UNP P20239
A	142	HIS	-	expression tag	UNP P20239
A	143	HIS	-	expression tag	UNP P20239
A	144	HIS	-	expression tag	UNP P20239
A	145	HIS	-	expression tag	UNP P20239
A	146	HIS	-	expression tag	UNP P20239
B	83	SER	ASN	engineered mutation	UNP P20239
B	139	LEU	-	expression tag	UNP P20239
B	140	GLU	-	expression tag	UNP P20239
B	141	HIS	-	expression tag	UNP P20239
B	142	HIS	-	expression tag	UNP P20239
B	143	HIS	-	expression tag	UNP P20239
B	144	HIS	-	expression tag	UNP P20239
B	145	HIS	-	expression tag	UNP P20239
B	146	HIS	-	expression tag	UNP P20239

- Molecule 2 is a protein called Heavy chain variable (VH) domain of anti-ZP2 monoclonal antibody IE-3.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	120	Total	C	N	O	S	0	0	0
			912	568	151	188	5			
2	X	120	Total	C	N	O	S	0	0	0
			912	568	151	188	5			

- Molecule 3 is a protein called Light chain variable (VL) domain of anti-ZP2 monoclonal antibody IE-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	115	Total	C	N	O	S	0	0	0
			883	553	147	180	3			
3	Y	116	Total	C	N	O	S	0	0	0
			894	560	152	179	3			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	132	Total	O	0	0
			132	132		
4	H	82	Total	O	0	0
			82	82		
4	L	77	Total	O	0	0
			77	77		
4	B	130	Total	O	0	0
			130	130		
4	X	100	Total	O	0	0
			100	100		
4	Y	95	Total	O	0	0
			95	95		

Chain Y:  84% 15%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.64Å 92.29Å 91.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.16 – 1.53 29.16 – 1.53	Depositor EDS
% Data completeness (in resolution range)	93.4 (29.16-1.53) 91.5 (29.16-1.53)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.53Å)	Xtrriage
Refinement program	PHENIX 1.21.2_5419+SVN	Depositor
R, R_{free}	0.206 , 0.233 0.210 , 0.237	Depositor DCC
R_{free} test set	6337 reflections (5.42%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtrriage
Anisotropy	0.271	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.258 for k,h,-l 0.044 for -h,-l,-k 0.046 for l,-k,h 0.044 for k,l,h 0.044 for l,h,k	Xtrriage
Reported twinning fraction	0.300 for k,h,-l	Depositor
Outliers	0 of 115130 reflections	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5768	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.12	0/795	0.35	0/1079
1	B	0.12	0/794	0.34	0/1078
2	H	0.10	0/932	0.28	0/1262
2	X	0.11	0/932	0.29	0/1262
3	L	0.14	0/902	0.40	0/1226
3	Y	0.12	0/913	0.37	0/1239
All	All	0.12	0/5268	0.34	0/7146

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	776	0	740	7	0
1	B	775	0	745	4	0
2	H	912	0	858	5	0
2	X	912	0	858	2	0
3	L	883	0	847	7	0
3	Y	894	0	867	12	0
4	A	132	0	0	3	0
4	B	130	0	0	0	0
4	H	82	0	0	2	0
4	L	77	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	X	100	0	0	0	0
4	Y	95	0	0	3	0
All	All	5768	0	4915	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:Y:19:THR:HG22	4:Y:277:HOH:O	1.73	0.87
2:H:78:LYS:NZ	4:H:201:HOH:O	2.24	0.71
3:Y:68:GLN:NE2	4:Y:204:HOH:O	2.27	0.66
1:A:88:LEU:HB3	1:B:88:LEU:HB3	1.78	0.65
2:H:42:LYS:NZ	2:H:95:SER:O	2.38	0.55

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/112 (84%)	93 (99%)	1 (1%)	0	100	100
1	B	94/112 (84%)	93 (99%)	1 (1%)	0	100	100
2	H	118/123 (96%)	118 (100%)	0	0	100	100
2	X	118/123 (96%)	117 (99%)	1 (1%)	0	100	100
3	L	113/117 (97%)	108 (96%)	5 (4%)	0	100	100
3	Y	114/117 (97%)	110 (96%)	4 (4%)	0	100	100
All	All	651/704 (92%)	639 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/104 (85%)	88 (100%)	0	100	100
1	B	88/104 (85%)	88 (100%)	0	100	100
2	H	100/102 (98%)	100 (100%)	0	100	100
2	X	100/102 (98%)	100 (100%)	0	100	100
3	L	100/102 (98%)	100 (100%)	0	100	100
3	Y	101/102 (99%)	101 (100%)	0	100	100
All	All	577/616 (94%)	577 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
3	Y	81	GLN
3	Y	75	HIS
3	Y	48	ASN
1	B	70	ASN
3	Y	57	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	96/112 (85%)	-0.87	0 100 100	25, 28, 37, 79	0
1	B	96/112 (85%)	-0.87	0 100 100	24, 29, 44, 78	0
2	H	120/123 (97%)	-0.44	2 (1%) 69 77	28, 37, 58, 68	0
2	X	120/123 (97%)	-0.75	0 100 100	25, 32, 46, 68	0
3	L	115/117 (98%)	0.02	4 (3%) 47 56	27, 46, 89, 117	0
3	Y	116/117 (99%)	-0.52	0 100 100	25, 36, 51, 88	0
All	All	663/704 (94%)	-0.56	6 (0%) 81 86	24, 34, 64, 117	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	36	GLY	3.1
2	H	60	PRO	2.4
3	L	94	GLY	2.4
2	H	139	SER	2.1
3	L	109	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.