

# wwPDB X-ray Structure Validation Summary Report (i)

Sep 9, 2023 – 09:38 PM EDT

PDB ID : 4HL6

Title : YfdE from Escherichia coli

Authors: Mullins, E.A.; Sullivan, K.L.; Nyffeler, K.E.; Kappock, T.J.

Deposited on : 2012-10-16

Resolution : 2.12 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467 Xtriage (Phenix): 1.13

EDS : 2.35.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
$R_{free}$	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	401	83%	10%	6%
1	В	401	86%	8%	6%
1	С	401	84%	9%	6%
1	D	401	83%	9%	7%
1	Е	401	86%	7%	7%



Continued from previous page...

Mol	Chain	Length	Quality of chain		
	_		3%		
1	F	401	82%	11%	• 6%



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 18281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Uncharacterized protein YfdE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	A	375	Total	С	N	О	S	0	0	0
1	A	313	2878	1826	493	539	20	U	0	
1	В	376	Total	С	N	О	S	0	0	0
1	Ъ	310	2884	1829	494	541	20	U	U	U
1	С	375	Total	С	N	О	S	0	0	0
1		310	2878	1826	493	539	20	U	0	
1	D	374	Total	С	N	О	S	0	0	0
1	D	314	2871	1823	492	536	20	U	0	
1	Е	374	Total	С	N	О	S	0	0	0
1	12	314	2871	1823	492	536	20	U	0	
1	F	375	Total	С	N	О	S	0	0	0
1	I'	313	2878	1826	493	539	20	U	U	

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP P76518
A	-18	GLY	-	expression tag	UNP P76518
A	-17	SER	-	expression tag	UNP P76518
A	-16	SER	-	expression tag	UNP P76518
A	-15	HIS	-	expression tag	UNP P76518
A	-14	HIS	-	expression tag	UNP P76518
A	-13	HIS	-	expression tag	UNP P76518
A	-12	HIS	-	expression tag	UNP P76518
A	-11	HIS	-	expression tag	UNP P76518
A	-10	HIS	-	expression tag	UNP P76518
A	-9	SER	-	expression tag	UNP P76518
A	-8	SER	-	expression tag	UNP P76518
A	-7	GLY	-	expression tag	UNP P76518
A	-6	LEU	=	expression tag	UNP P76518
A	-5	VAL	=	expression tag	UNP P76518
A	-4	PRO	-	expression tag	UNP P76518
A	-3	ARG	-	expression tag	UNP P76518



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P76518
A	-1	SER	_	expression tag	UNP P76518
A	0	HIS	-	expression tag	UNP P76518
В	-19	MET	_	expression tag	UNP P76518
В	-18	GLY	-	expression tag	UNP P76518
В	-17	SER	_	expression tag	UNP P76518
В	-16	SER	-	expression tag	UNP P76518
В	-15	HIS	-	expression tag	UNP P76518
В	-14	HIS	-	expression tag	UNP P76518
В	-13	HIS	_	expression tag	UNP P76518
В	-12	HIS	-	expression tag	UNP P76518
В	-11	HIS	-	expression tag	UNP P76518
В	-10	HIS	-	expression tag	UNP P76518
В	-9	SER	_	expression tag	UNP P76518
В	-8	SER	-	expression tag	UNP P76518
В	-7	GLY	_	expression tag	UNP P76518
В	-6	LEU	-	expression tag	UNP P76518
В	-5	VAL	-	expression tag	UNP P76518
В	-4	PRO	_	expression tag	UNP P76518
В	-3	ARG	-	expression tag	UNP P76518
В	-2	GLY	_	expression tag	UNP P76518
В	-1	SER	-	expression tag	UNP P76518
В	0	HIS	-	expression tag	UNP P76518
С	-19	MET	-	expression tag	UNP P76518
С	-18	GLY	-	expression tag	UNP P76518
С	-17	SER	-	expression tag	UNP P76518
С	-16	SER	-	expression tag	UNP P76518
С	-15	HIS	-	expression tag	UNP P76518
С	-14	HIS	-	expression tag	UNP P76518
С	-13	HIS	-	expression tag	UNP P76518
С	-12	HIS	-	expression tag	UNP P76518
С	-11	HIS	-	expression tag	UNP P76518
С	-10	HIS	-	expression tag	UNP P76518
С	-9	SER	-	expression tag	UNP P76518
С	-8	SER	-	expression tag	UNP P76518
С	-7	GLY	-	expression tag	UNP P76518
С	-6	LEU	-	expression tag	UNP P76518
С	-5	VAL	-	expression tag	UNP P76518
С	-4	PRO	-	expression tag	UNP P76518
С	-3	ARG	-	expression tag	UNP P76518
С	-2	GLY	-	expression tag	UNP P76518
С	-1	SER	-	expression tag	UNP P76518



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	0	HIS	-	expression tag	UNP P76518
D	-19	MET	-	expression tag	UNP P76518
D	-18	GLY	_	expression tag	UNP P76518
D	-17	SER	-	expression tag	UNP P76518
D	-16	SER	-	expression tag	UNP P76518
D	-15	HIS	-	expression tag	UNP P76518
D	-14	HIS	-	expression tag	UNP P76518
D	-13	HIS	-	expression tag	UNP P76518
D	-12	HIS	-	expression tag	UNP P76518
D	-11	HIS	-	expression tag	UNP P76518
D	-10	HIS	-	expression tag	UNP P76518
D	-9	SER	-	expression tag	UNP P76518
D	-8	SER	-	expression tag	UNP P76518
D	-7	GLY	-	expression tag	UNP P76518
D	-6	LEU	-	expression tag	UNP P76518
D	-5	VAL	-	expression tag	UNP P76518
D	-4	PRO	-	expression tag	UNP P76518
D	-3	ARG	-	expression tag	UNP P76518
D	-2	GLY	-	expression tag	UNP P76518
D	-1	SER	-	expression tag	UNP P76518
D	0	HIS	-	expression tag	UNP P76518
Е	-19	MET	-	expression tag	UNP P76518
Е	-18	GLY	-	expression tag	UNP P76518
Е	-17	SER	-	expression tag	UNP P76518
Е	-16	SER	-	expression tag	UNP P76518
Е	-15	HIS	-	expression tag	UNP P76518
Е	-14	HIS	-	expression tag	UNP P76518
Е	-13	HIS	-	expression tag	UNP P76518
Е	-12	HIS	-	expression tag	UNP P76518
Е	-11	HIS	-	expression tag	UNP P76518
Е	-10	HIS	-	expression tag	UNP P76518
Е	-9	SER	-	expression tag	UNP P76518
Е	-8	SER	-	expression tag	UNP P76518
Е	-7	GLY	-	expression tag	UNP P76518
Е	-6	LEU	-	expression tag	UNP P76518
Е	-5	VAL		expression tag	UNP P76518
Е	-4	PRO		expression tag	UNP P76518
Е	-3	ARG	-	expression tag	UNP P76518
Е	-2	GLY		expression tag	UNP P76518
Е	-1	SER	-	expression tag	UNP P76518
Е	0	HIS	-	expression tag	UNP P76518
F	-19	MET	-	expression tag	UNP P76518



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	-18	GLY	-	expression tag	UNP P76518
F	-17	SER	-	expression tag	UNP P76518
F	-16	SER	-	expression tag	UNP P76518
F	-15	HIS	-	expression tag	UNP P76518
F	-14	HIS	-	expression tag	UNP P76518
F	-13	HIS	-	expression tag	UNP P76518
F	-12	HIS	-	expression tag	UNP P76518
F	-11	HIS	-	expression tag	UNP P76518
F	-10	HIS	-	expression tag	UNP P76518
F	-9	SER	-	expression tag	UNP P76518
F	-8	SER	-	expression tag	UNP P76518
F	-7	GLY	-	expression tag	UNP P76518
F	-6	LEU	-	expression tag	UNP P76518
F	-5	VAL	-	expression tag	UNP P76518
F	-4	PRO	-	expression tag	UNP P76518
F	-3	ARG	-	expression tag	UNP P76518
F	-2	GLY	-	expression tag	UNP P76518
F	-1	SER		expression tag	UNP P76518
F	0	HIS	-	expression tag	UNP P76518

#### • Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	185	Total O 185 185	0	0
2	В	226	Total O 226 226	0	0
2	С	203	Total O 203 203	0	0
2	D	173	Total O 173 173	0	0
2	E	118	Total O 118 118	0	0
2	F	116	Total O 116 116	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein YfdE







 $\bullet$  Molecule 1: Uncharacterized protein YfdE







 $\bullet$  Molecule 1: Uncharacterized protein YfdE









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	111.42Å 79.53Å 276.19Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $99.95^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	42.48 - 2.12	Depositor
Resolution (A)	43.81 - 2.12	EDS
% Data completeness	79.7 (42.48-2.12)	Depositor
(in resolution range)	76.1 (43.81-2.12)	EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$< I/\sigma(I) > 1$	2.16 (at 2.12Å)	Xtriage
Refinement program	PHENIX 1.7.3_928	Depositor
D.D.	0.189 , 0.236	Depositor
$R, R_{free}$	0.185 , $0.230$	DCC
$R_{free}$ test set	5456 reflections $(5.06%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.9	Xtriage
Anisotropy	0.677	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.33, 36.4	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	18281	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.24	0/2943	0.42	0/3990
1	В	0.24	0/2949	0.42	0/3998
1	С	0.24	0/2943	0.42	0/3990
1	D	0.24	0/2936	0.42	0/3982
1	Е	0.23	0/2936	0.42	0/3982
1	F	0.23	0/2943	0.41	0/3990
All	All	0.24	0/17650	0.42	0/23932

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

# 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2878	0	2858	26	0
1	В	2884	0	2863	17	0
1	С	2878	0	2858	22	0
1	D	2871	0	2853	27	0
1	Е	2871	0	2853	13	0
1	F	2878	0	2858	26	0
2	A	185	0	0	1	0
2	В	226	0	0	0	0
2	С	203	0	0	1	0
2	D	173	0	0	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
2	Ε	118	0	0	2	0
2	F	116	0	0	2	0
All	All	18281	0	17143	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} &  ext{Interatomic} \ &  ext{distance} \ &  ext{(Å)} \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:A:192:GLU:HG2	1:B:192:GLU:HG2	1.53	0.90
1:A:74:VAL:HG12	1:A:372:GLY:HA2	1.60	0.81
1:D:74:VAL:HG12	1:D:372:GLY:HA2	1.66	0.77
1:F:156:GLU:OE2	2:F:441:HOH:O	2.09	0.70
1:D:339:ILE:HD12	1:D:362:MET:HG3	1.75	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	A	373/401 (93%)	360 (96%)	11 (3%)	2(0%)	29	25
1	В	374/401 (93%)	365 (98%)	7 (2%)	2 (0%)	29	25
1	С	373/401 (93%)	361 (97%)	9 (2%)	3 (1%)	19	14
1	D	372/401 (93%)	362 (97%)	9 (2%)	1 (0%)	41	40
1	E	372/401 (93%)	350 (94%)	21 (6%)	1 (0%)	41	40
1	F	373/401 (93%)	361 (97%)	11 (3%)	1 (0%)	41	40
All	All	2237/2406 (93%)	2159 (96%)	68 (3%)	10 (0%)	34	32



5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	21	VAL
1	В	21	VAL
1	A	228	ARG
1	С	21	VAL
1	С	23	ASN

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	311/334 (93%)	304 (98%)	7 (2%)	50 53
1	В	312/334 (93%)	308 (99%)	4 (1%)	69 74
1	С	311/334 (93%)	306 (98%)	5 (2%)	62 68
1	D	310/334 (93%)	304 (98%)	6 (2%)	57 61
1	E	310/334 (93%)	303 (98%)	7 (2%)	50 53
1	F	311/334 (93%)	304 (98%)	7 (2%)	50 53
All	All	1865/2004 (93%)	1829 (98%)	36 (2%)	57 61

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	296	ARG
1	F	269	THR
1	F	20	HIS
1	F	153	ILE
1	С	118	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

## 5.3.3 RNA (i)

There are no RNA molecules in this entry.



# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	375/401 (93%)	-0.13	2 (0%) 91 92	19, 29, 47, 66	0
1	В	376/401 (93%)	-0.23	0 100 100	17, 26, 41, 59	0
1	С	375/401 (93%)	-0.23	4 (1%) 80 84	18, 28, 44, 62	0
1	D	374/401 (93%)	-0.14	0 100 100	20, 30, 48, 60	0
1	E	374/401 (93%)	-0.00	6 (1%) 72 76	19, 34, 52, 72	0
1	F	375/401 (93%)	0.21	11 (2%) 51 57	20, 38, 59, 74	0
All	All	2249/2406 (93%)	-0.09	23 (1%) 82 85	17, 30, 51, 74	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	381	SER	5.3
1	F	381	SER	4.0
1	С	59	GLY	3.6
1	Е	196	ARG	3.3
1	F	245	GLN	3.2

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

# 6.4 Ligands (i)

There are no ligands in this entry.



# 6.5 Other polymers (i)

There are no such residues in this entry.

