



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 5, 2026 – 04:42 pm GMT

PDB ID : 9HT9 / pdb_00009ht9
Title : CutC in complex with inhibitor1
Authors : Petersen, J.
Deposited on : 2024-12-19
Resolution : 1.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : ?? (??), CSD ??CSD?? (????)
Xtrriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1150 (1.84-1.84)
Clashscore	180529	1248 (1.84-1.84)
Ramachandran outliers	177936	1240 (1.84-1.84)
Sidechain outliers	177891	1240 (1.84-1.84)
RSRZ outliers	164620	1149 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	800	 2% 91% 8% ..
1	B	800	 % 92% 6% ..
1	C	800	 6% 90% 9% ..
1	D	800	 2% 92% 7% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 26165 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Choline trimethylamine-lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	794	Total	C	N	O	S	0	0	0
			6263	3971	1062	1185	45			
1	B	794	Total	C	N	O	S	0	0	0
			6263	3971	1062	1185	45			
1	C	794	Total	C	N	O	S	0	0	0
			6263	3971	1062	1185	45			
1	D	794	Total	C	N	O	S	0	0	0
			6263	3971	1062	1185	45			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	47	MET	-	initiating methionine	UNP Q30W70
A	48	HIS	-	expression tag	UNP Q30W70
A	49	HIS	-	expression tag	UNP Q30W70
A	50	HIS	-	expression tag	UNP Q30W70
A	51	HIS	-	expression tag	UNP Q30W70
A	52	HIS	-	expression tag	UNP Q30W70
A	53	HIS	-	expression tag	UNP Q30W70
B	47	MET	-	initiating methionine	UNP Q30W70
B	48	HIS	-	expression tag	UNP Q30W70
B	49	HIS	-	expression tag	UNP Q30W70
B	50	HIS	-	expression tag	UNP Q30W70
B	51	HIS	-	expression tag	UNP Q30W70
B	52	HIS	-	expression tag	UNP Q30W70
B	53	HIS	-	expression tag	UNP Q30W70
C	47	MET	-	initiating methionine	UNP Q30W70
C	48	HIS	-	expression tag	UNP Q30W70
C	49	HIS	-	expression tag	UNP Q30W70
C	50	HIS	-	expression tag	UNP Q30W70
C	51	HIS	-	expression tag	UNP Q30W70
C	52	HIS	-	expression tag	UNP Q30W70
C	53	HIS	-	expression tag	UNP Q30W70

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	47	MET	-	initiating methionine	UNP Q30W70
D	48	HIS	-	expression tag	UNP Q30W70
D	49	HIS	-	expression tag	UNP Q30W70
D	50	HIS	-	expression tag	UNP Q30W70
D	51	HIS	-	expression tag	UNP Q30W70
D	52	HIS	-	expression tag	UNP Q30W70
D	53	HIS	-	expression tag	UNP Q30W70

- Molecule 2 is (3 {S})-1-azabicyclo[2.2.2]octan-3-ol (CCD ID: A1IXC) (formula: C₇H₁₃NO) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			9	7	1	1		
2	B	1	Total	C	N	O	0	0
			9	7	1	1		
2	C	1	Total	C	N	O	0	0
			9	7	1	1		
2	D	1	Total	C	N	O	0	0
			9	7	1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	328	Total	O	0	0
			328	328		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	330	Total 330	O 330	0	0
3	C	186	Total 186	O 186	0	0
3	D	233	Total 233	O 233	0	0



● Molecule 1: Choline trimethylamine-lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	81.66Å 104.76Å 119.49Å 102.43° 108.54° 104.91°	Depositor
Resolution (Å)	107.17 – 1.84 107.17 – 1.84	Depositor EDS
% Data completeness (in resolution range)	97.0 (107.17-1.84) 97.1 (107.17-1.84)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 1.83Å)	Xtrriage
Refinement program	REFMAC 5.8.0267 STEINER	Depositor
R, R_{free}	0.182 , 0.216 0.190 , 0.222	Depositor DCC
R_{free} test set	14479 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtrriage
Anisotropy	0.093	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 22.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	26165	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1IXC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.11	2/6408 (0.0%)	1.35	12/8678 (0.1%)
1	B	1.11	2/6408 (0.0%)	1.35	16/8678 (0.2%)
1	C	1.07	0/6408	1.38	6/8678 (0.1%)
1	D	1.10	3/6408 (0.0%)	1.38	6/8678 (0.1%)
All	All	1.10	7/25632 (0.0%)	1.37	40/34712 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	2
1	D	0	2
All	All	0	6

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	571	HIS	CE1-NE2	5.89	1.38	1.32
1	B	146	GLU	CD-OE2	5.63	1.36	1.25
1	A	593	ASP	C-O	5.48	1.31	1.23
1	B	570	ALA	C-O	5.39	1.30	1.24
1	D	462	HIS	CE1-NE2	5.27	1.37	1.32

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	238	ARG	CG-CD-NE	-7.57	95.35	112.00
1	B	540	ASP	CB-CA-C	-7.57	98.70	110.81
1	B	539	TYR	CA-C-N	7.28	130.37	120.54
1	B	539	TYR	C-N-CA	7.28	130.37	120.54
1	A	539	TYR	CA-C-N	7.26	130.32	120.38

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	526	GLY	Peptide
1	B	526	GLY	Peptide
1	C	526	GLY	Peptide
1	C	766	GLY	Peptide
1	D	526	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6263	0	6135	42	0
1	B	6263	0	6135	33	0
1	C	6263	0	6135	47	0
1	D	6263	0	6135	34	0
2	A	9	0	0	0	0
2	B	9	0	0	0	0
2	C	9	0	0	0	0
2	D	9	0	0	0	0
3	A	328	0	0	9	0
3	B	330	0	0	6	0
3	C	186	0	0	7	0
3	D	233	0	0	7	0
All	All	26165	0	24540	138	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 138 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:786:MET:SD	1:D:786:MET:SD	2.53	1.06
1:D:800:GLU:HG2	3:D:1219:HOH:O	1.56	1.02
1:B:674:ASP:HB2	3:B:1275:HOH:O	1.70	0.89
1:B:130:ARG:HD2	3:B:1140:HOH:O	1.74	0.87
1:A:130:ARG:CD	3:A:1058:HOH:O	2.26	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	792/800 (99%)	773 (98%)	17 (2%)	2 (0%)	37 25
1	B	792/800 (99%)	768 (97%)	21 (3%)	3 (0%)	30 18
1	C	792/800 (99%)	765 (97%)	24 (3%)	3 (0%)	30 18
1	D	792/800 (99%)	769 (97%)	21 (3%)	2 (0%)	37 25
All	All	3168/3200 (99%)	3075 (97%)	83 (3%)	10 (0%)	37 25

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	652	LEU
1	C	700	ILE
1	D	700	ILE
1	A	700	ILE
1	B	700	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	669/676 (99%)	656 (98%)	13 (2%)	52	36
1	B	669/676 (99%)	654 (98%)	15 (2%)	47	31
1	C	669/676 (99%)	657 (98%)	12 (2%)	54	39
1	D	669/676 (99%)	651 (97%)	18 (3%)	40	22
All	All	2676/2704 (99%)	2618 (98%)	58 (2%)	47	31

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	178	ASN
1	D	807	LYS
1	C	718	ARG
1	D	765	SER
1	D	551	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	408	HIS
1	C	808	HIS
1	C	636	GLN
1	D	68	ASN
1	B	68	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.



There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	794/800 (99%)	-0.23	12 (1%) 71 79	13, 20, 33, 54	0
1	B	794/800 (99%)	-0.26	7 (0%) 81 87	12, 20, 33, 52	0
1	C	794/800 (99%)	0.33	49 (6%) 28 29	17, 27, 50, 74	0
1	D	794/800 (99%)	0.04	18 (2%) 61 66	16, 24, 39, 58	0
All	All	3176/3200 (99%)	-0.03	86 (2%) 56 61	12, 22, 41, 74	0

The worst 5 of 86 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	526	GLY	5.0
1	A	285	ARG	4.5
1	D	53	HIS	4.5
1	A	526	GLY	4.4
1	A	53	HIS	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	A1IXC	C	901	9/9	0.96	0.06	20,21,22,23	0
2	A1IXC	D	901	9/9	0.96	0.06	16,17,18,19	0
2	A1IXC	A	901	9/9	0.97	0.06	14,15,15,16	0
2	A1IXC	B	901	9/9	0.97	0.05	13,14,14,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers [i](#)

There are no such residues in this entry.