

wwPDB X-ray Structure Validation Summary Report (i)

Sep 20, 2023 – 08:34 PM EDT

PDB ID : 5HUP

Title : Crystal Structure of NadC from Streptococcus pyogenes

Authors: Booth, W.T.; Chruszcz, M.

Deposited on : 2016-01-27

Resolution : 3.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.35.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	1486 (3.50-3.34)
Clashscore	141614	1572 (3.50-3.34)
Ramachandran outliers	138981	1534 (3.50-3.34)
Sidechain outliers	138945	1535 (3.50-3.34)
RSRZ outliers	127900	1395 (3.50-3.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	315	83% 7%	10%
1	В	315	80% 10%	10%
1	C	315	83% 6%	11%
1	E	315	79% 10%	10%
1	F	315	84% 6%	10%



Continued from previous page...

Mol	Chain	Length	Quality of chain		
1	Н	315	77%	13%	10%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 12751 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Nicotinate-nucleotide pyrophosphorylase (Carboxylating).

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	285	Total	С	N	О	S	0	0	0
1	A	200	2137	1352	368	408	9	0	0	
1	В	285	Total	С	N	О	S	0	0	0
1	Б	200	2146	1354	368	415	9	0	U	
1	C	280	Total	С	N	О	S	0	0	0
1		200	1994	1254	344	390	6	U	0	
1	Е	284	Total	С	N	О	S	0	0	0
1	12	204	2102	1328	363	403	8	0	0	
1	F	F 284	Total	С	N	О	S	0	0	0
1			2137	1350	366	412	9	0	0	
1	Н	285	Total	С	N	О	S	0	0	0
1	1 H	285	2150	1357	372	412	9		U	

There are 150 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-24	MET	-	initiating methionine	UNP Q1J647
A	-23	HIS	-	expression tag	UNP Q1J647
A	-22	HIS	-	expression tag	UNP Q1J647
A	-21	HIS	-	expression tag	UNP Q1J647
A	-20	HIS	-	expression tag	UNP Q1J647
A	-19	HIS	-	expression tag	UNP Q1J647
A	-18	HIS	-	expression tag	UNP Q1J647
A	-17	SER	-	expression tag	UNP Q1J647
A	-16	SER	-	expression tag	UNP Q1J647
A	-15	GLY	-	expression tag	UNP Q1J647
A	-14	VAL	-	expression tag	UNP Q1J647
A	-13	ASP	-	expression tag	UNP Q1J647
A	-12	LEU	-	expression tag	UNP Q1J647
A	-11	GLY	-	expression tag	UNP Q1J647
A	-10	THR	-	expression tag	UNP Q1J647
A	-9	GLU	-	expression tag	UNP Q1J647
A	-8	ASN	-	expression tag	UNP Q1J647



 $Continued\ from\ previous\ page...$

A		Modelled	Actual	${f Comment}$	Reference
1	-7	LEU	-	expression tag	UNP Q1J647
A	-6	TYR	-	expression tag	UNP Q1J647
A	-5	PHE	-	expression tag	UNP Q1J647
A	-4	GLN	-	expression tag	UNP Q1J647
A	-3	SER	-	expression tag	UNP Q1J647
A	-2	GLY	-	expression tag	UNP Q1J647
A	-1	SER	-	expression tag	UNP Q1J647
A	0	GLY	-	expression tag	UNP Q1J647
В	-24	MET	-	initiating methionine	UNP Q1J647
В	-23	HIS	-	expression tag	UNP Q1J647
В	-22	HIS	-	expression tag	UNP Q1J647
В	-21	HIS	-	expression tag	UNP Q1J647
В	-20	HIS	-	expression tag	UNP Q1J647
В	-19	HIS	-	expression tag	UNP Q1J647
В	-18	HIS	-	expression tag	UNP Q1J647
В	-17	SER	-	expression tag	UNP Q1J647
В	-16	SER	-	expression tag	UNP Q1J647
В	-15	GLY	-	expression tag	UNP Q1J647
В	-14	VAL	-	expression tag	UNP Q1J647
В	-13	ASP	-	expression tag	UNP Q1J647
В	-12	LEU	-	expression tag	UNP Q1J647
В	-11	GLY	-	expression tag	UNP Q1J647
В	-10	THR	-	expression tag	UNP Q1J647
В	-9	GLU	-	expression tag	UNP Q1J647
В	-8	ASN	-	expression tag	UNP Q1J647
В	-7	LEU	-	expression tag	UNP Q1J647
В	-6	TYR	-	expression tag	UNP Q1J647
В	-5	PHE	-	expression tag	UNP Q1J647
В	-4	GLN	-	expression tag	UNP Q1J647
В	-3	SER	-	expression tag	UNP Q1J647
В	-2	GLY	-	expression tag	UNP Q1J647
В	-1	SER	-	expression tag	UNP Q1J647
В	0	GLY	-	expression tag	UNP Q1J647
С	-24	MET	-	initiating methionine	UNP Q1J647
С	-23	HIS	-	expression tag	UNP Q1J647
С	-22	HIS	-	expression tag	UNP Q1J647
С	-21	HIS	-	expression tag	UNP Q1J647
С	-20	HIS	-	expression tag	UNP Q1J647
С	-19	HIS	-	expression tag	UNP Q1J647
С	-18	HIS	-	expression tag	UNP Q1J647
С	-17	SER	-	expression tag	UNP Q1J647
С	-16	SER	-	expression tag	UNP Q1J647



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
С	-15	GLY	-	expression tag	UNP Q1J647
С	-14	VAL	-	expression tag	UNP Q1J647
С	-13	ASP	-	expression tag	UNP Q1J647
С	-12	LEU	-	expression tag	UNP Q1J647
С	-11	GLY	-	expression tag	UNP Q1J647
С	-10	THR	-	expression tag	UNP Q1J647
С	-9	GLU	-	expression tag	UNP Q1J647
С	-8	ASN	-	expression tag	UNP Q1J647
С	-7	LEU	-	expression tag	UNP Q1J647
С	-6	TYR	-	expression tag	UNP Q1J647
С	-5	PHE	-	expression tag	UNP Q1J647
С	-4	GLN	-	expression tag	UNP Q1J647
С	-3	SER	-	expression tag	UNP Q1J647
С	-2	GLY	-	expression tag	UNP Q1J647
С	-1	SER	-	expression tag	UNP Q1J647
С	0	GLY	-	expression tag	UNP Q1J647
Е	-24	MET	-	initiating methionine	UNP Q1J647
Е	-23	HIS	-	expression tag	UNP Q1J647
Е	-22	HIS	-	expression tag	UNP Q1J647
Е	-21	HIS	-	expression tag	UNP Q1J647
Е	-20	HIS	-	expression tag	UNP Q1J647
Е	-19	HIS	-	expression tag	UNP Q1J647
Е	-18	HIS	-	expression tag	UNP Q1J647
Е	-17	SER	-	expression tag	UNP Q1J647
Е	-16	SER	-	expression tag	UNP Q1J647
Е	-15	GLY	-	expression tag	UNP Q1J647
Е	-14	VAL	-	expression tag	UNP Q1J647
Е	-13	ASP	-	expression tag	UNP Q1J647
Е	-12	LEU	-	expression tag	UNP Q1J647
Е	-11	GLY	-	expression tag	UNP Q1J647
Е	-10	THR	-	expression tag	UNP Q1J647
Е	-9	GLU	-	expression tag	UNP Q1J647
Е	-8	ASN	-	expression tag	UNP Q1J647
Е	-7	LEU	-	expression tag	UNP Q1J647
Е	-6	TYR	-	expression tag	UNP Q1J647
Е	-5	PHE	-	expression tag	UNP Q1J647
Е	-4	GLN		expression tag	UNP Q1J647
Е	-3	SER	-	expression tag	UNP Q1J647
Е	-2	GLY	-	expression tag	UNP Q1J647
Е	-1	SER	-	expression tag	UNP Q1J647
Е	0	GLY	-	expression tag	UNP Q1J647
F	-24	MET	-	initiating methionine	UNP Q1J647



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
F	-23	HIS	-	expression tag	UNP Q1J647
F	-22	HIS	-	expression tag	UNP Q1J647
F	-21	HIS	-	expression tag	UNP Q1J647
F	-20	HIS	-	expression tag	UNP Q1J647
F	-19	HIS	-	expression tag	UNP Q1J647
F	-18	HIS	-	expression tag	UNP Q1J647
F	-17	SER	-	expression tag	UNP Q1J647
F	-16	SER	-	expression tag	UNP Q1J647
F	-15	GLY	-	expression tag	UNP Q1J647
F	-14	VAL	-	expression tag	UNP Q1J647
F	-13	ASP	-	expression tag	UNP Q1J647
F	-12	LEU	-	expression tag	UNP Q1J647
F	-11	GLY	-	expression tag	UNP Q1J647
F	-10	THR	-	expression tag	UNP Q1J647
F	-9	GLU	-	expression tag	UNP Q1J647
F	-8	ASN	-	expression tag	UNP Q1J647
F	-7	LEU	-	expression tag	UNP Q1J647
F	-6	TYR	-	expression tag	UNP Q1J647
F	-5	PHE	-	expression tag	UNP Q1J647
F	-4	GLN	-	expression tag	UNP Q1J647
F	-3	SER	-	expression tag	UNP Q1J647
F	-2	GLY	-	expression tag	UNP Q1J647
F	-1	SER	-	expression tag	UNP Q1J647
F	0	GLY	-	expression tag	UNP Q1J647
Н	-24	MET	-	initiating methionine	UNP Q1J647
Н	-23	HIS	-	expression tag	UNP Q1J647
Н	-22	HIS	-	expression tag	UNP Q1J647
Н	-21	HIS	-	expression tag	UNP Q1J647
Н	-20	HIS	-	expression tag	UNP Q1J647
Н	-19	HIS	-	expression tag	UNP Q1J647
Н	-18	HIS	-	expression tag	UNP Q1J647
Н	-17	SER	-	expression tag	UNP Q1J647
Н	-16	SER	-	expression tag	UNP Q1J647
Н	-15	GLY	-	expression tag	UNP Q1J647
Н	-14	VAL	-	expression tag	UNP Q1J647
Н	-13	ASP	-	expression tag	UNP Q1J647
Н	-12	LEU	-	expression tag	UNP Q1J647
Н	-11	GLY	-	expression tag	UNP Q1J647
Н	-10	THR	-	expression tag	UNP Q1J647
Н	-9	GLU	-	expression tag	UNP Q1J647
Н	-8	ASN	-	expression tag	UNP Q1J647
Н	-7	LEU	-	expression tag	UNP Q1J647



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
Н	-6	TYR	-	expression tag	UNP Q1J647
Н	-5	PHE	-	expression tag	UNP Q1J647
Н	-4	GLN	-	expression tag	UNP Q1J647
Н	-3	SER	-	expression tag	UNP Q1J647
Н	-2	GLY	-	expression tag	UNP Q1J647
Н	-1	SER	-	expression tag	UNP Q1J647
Н	0	GLY	-	expression tag	UNP Q1J647

 \bullet Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	В	1	Total O S 5 4 1	0	0
2	В	1	Total O S 5 4 1	0	0
2	С	1	Total O S 5 4 1	0	0
2	С	1	Total O S 5 4 1	0	0



Continued from previous page...


Mol	Chain	Residues	Atom	ıs	ZeroOcc	AltConf
2	С	1	Total C) S	0	0
		1	5 4		0	U
2	E	1	Total () S	0	0
	Ш	1	5 4		O	0
2	E	1	Total C) S	0	0
	L	1	5 4		O	0
2	F	1	Total C) S	0	0
	I.	1	5 4		U	Ŭ
$\frac{1}{2}$	F	1	Total C		0	0
	I.	1	5 4	1 1	O	U
$\frac{1}{2}$	Н	1	Total C) S	0	0
	11	1	5 4	1 1	O	U
2	Н	1	Total C) S	0	0
	11	1	5 4	1 1	O	U
2	Н	1	Total C) S	0	0
	11	1	5 4	1 1		0
2	Н	1	Total C) S	0	0
	11	1	5 4	1 1	U	U




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Nicotinate-nucleotide pyrophosphorylase (Carboxylating)








• Molecule 1: Nicotinate-nucleotide pyrophosphorylase (Carboxylating)





• Molecule 1: Nicotinate-nucleotide pyrophosphorylase (Carboxylating)





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	108.15Å 188.82Å 222.18Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 - 3.42	Depositor
Resolution (A)	20.17 - 3.42	EDS
% Data completeness	78.9 (20.00-3.42)	Depositor
(in resolution range)	79.4 (20.17-3.42)	EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$< I/\sigma(I) > 1$	4.79 (at 3.44Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
D.D.	0.233 , 0.294	Depositor
R, R_{free}	0.233 , 0.291	DCC
R_{free} test set	1221 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	67.9	Xtriage
Anisotropy	0.386	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.27 \; , 34.4$	EDS
L-test for twinning ²	$< L > = 0.52, < L^2> = 0.35$	Xtriage
Estimated twinning fraction	0.020 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l	Xtriage
Estimated twinning fraction	0.021 for 1/2 *h + 1/2 *k, 3/2 *h - 1/2 *k, -1	Alliage
F_o, F_c correlation	0.88	EDS
Total number of atoms	12751	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Во	nd lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.71	1/2168 (0.0%)	0.84	$4/2936 \; (0.1\%)$	
1	В	0.70	0/2179	0.89	5/2954~(0.2%)	
1	С	0.69	0/2023	0.84	$3/2756 \ (0.1\%)$	
1	Е	0.70	0/2134	0.83	6/2897~(0.2%)	
1	F	0.72	0/2168	0.86	$4/2937 \ (0.1\%)$	
1	Н	0.74	1/2183 (0.0%)	0.87	2/2958~(0.1%)	
All	All	0.71	$2/12855 \ (0.0\%)$	0.86	24/17438 (0.1%)	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$Ideal(\AA)$
1	Н	245	GLU	CD-OE2	-7.14	1.17	1.25
1	A	245	GLU	CD-OE2	-6.72	1.18	1.25

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	241	ARG	NE-CZ-NH2	16.30	128.45	120.30
1	В	241	ARG	NE-CZ-NH1	-10.23	115.19	120.30
1	Е	191	ARG	NE-CZ-NH1	7.33	123.97	120.30
1	С	84	ARG	NE-CZ-NH2	7.26	123.93	120.30
1	A	245	GLU	CG-CD-OE1	6.99	132.27	118.30

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2137	0	2112	14	0
1	В	2146	0	2101	24	0
1	С	1994	0	1840	16	0
1	Е	2102	0	2038	21	1
1	F	2137	0	2108	12	0
1	Н	2150	0	2114	31	1
2	A	20	0	0	0	0
2	В	10	0	0	0	0
2	С	15	0	0	0	0
2	Ε	10	0	0	1	0
2	F	10	0	0	0	0
2	Н	20	0	0	1	0
All	All	12751	0	12313	106	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$	
1:E:109:LEU:HD11	1:E:282:MET:HG2	1.58	0.83	
1:B:37:ASP:HA	1:B:99:ARG:HH21	1.46	0.81	
1:E:109:LEU:HD11	1:E:282:MET:CG	2.12	0.78	
1:A:207:LEU:HD12	1:A:208:ALA:N	2.00	0.76	
1:H:35:ILE:HG22	1:H:35:ILE:O	1.85	0.76	

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1 Atom-2		$egin{aligned} ext{Interatomic} \ ext{distance} & (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:E:225:ASN:OD1	1:H:127:GLU:OE2[7_454]	2.11	0.09



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	ed Favoured Allowed		Outliers Per		centiles	
1	A	283/315 (90%)	270 (95%)	13 (5%)	0	100	100	
1	В	283/315 (90%)	267 (94%)	16 (6%)	0	100	100	
1	С	276/315 (88%)	264 (96%)	12 (4%)	0	100	100	
1	${ m E}$	282/315~(90%)	267 (95%)	15 (5%)	0	100	100	
1	F	$282/315 \ (90\%)$	268 (95%)	14 (5%)	0	100	100	
1	Н	283/315 (90%)	270 (95%)	13 (5%)	0	100	100	
All	All	1689/1890 (89%)	1606 (95%)	83 (5%)	0	100	100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	220/260 (85%)	219 (100%)	1 (0%)	88 95		
1	В	223/260 (86%)	221 (99%)	2 (1%)	78 90		
1	С	188/260 (72%)	186 (99%)	2 (1%)	73 87		
1	${ m E}$	212/260 (82%)	210 (99%)	2 (1%)	78 90		
1	F	222/260 (85%)	221 (100%)	1 (0%)	88 95		
1	Н	223/260 (86%)	221 (99%)	2 (1%)	78 90		
All	All	1288/1560 (83%)	1278 (99%)	10 (1%)	81 92		

5 of 10 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	F	168	ASP
1	Н	164	PHE
1	Н	168	ASP
1	С	164	PHE
1	С	199	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	26	HIS
1	С	161	ASN
1	Н	26	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	${ m Res}$	Link	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	Е	302	-	4,4,4	0.36	0	6,6,6	0.34	0
2	SO4	A	303	-	4,4,4	0.40	0	6,6,6	0.17	0



Mol	Trino	Chain	Res	Link	В	Bond lengths			Bond angles		
IVIOI	Type	Chain	Res Lilik		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	SO4	A	304	-	4,4,4	0.37	0	6,6,6	0.12	0	
2	SO4	A	301	-	4,4,4	0.36	0	6,6,6	0.22	0	
2	SO4	Н	303	-	4,4,4	0.37	0	6,6,6	0.18	0	
2	SO4	В	302	-	4,4,4	0.36	0	6,6,6	0.23	0	
2	SO4	Е	301	-	4,4,4	0.35	0	6,6,6	0.32	0	
2	SO4	F	302	-	4,4,4	0.37	0	6,6,6	0.20	0	
2	SO4	Н	301	-	4,4,4	0.34	0	6,6,6	0.50	0	
2	SO4	С	301	-	4,4,4	0.36	0	6,6,6	0.14	0	
2	SO4	F	301	-	4,4,4	0.36	0	6,6,6	0.40	0	
2	SO4	Н	302	-	4,4,4	0.40	0	6,6,6	0.13	0	
2	SO4	С	303	-	4,4,4	0.32	0	6,6,6	0.15	0	
2	SO4	Н	304	-	4,4,4	0.38	0	6,6,6	0.09	0	
2	SO4	A	302	-	4,4,4	0.41	0	6,6,6	0.28	0	
2	SO4	В	301	-	4,4,4	0.40	0	6,6,6	0.39	0	
2	SO4	С	302	-	4,4,4	0.39	0	6,6,6	0.28	0	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Е	301	SO4	1	0
2	Н	301	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	285/315~(90%)	-0.35	1 (0%) 92 91	39, 76, 116, 130	0
1	В	285/315~(90%)	-0.17	3 (1%) 80 79	53, 78, 118, 151	0
1	С	280/315 (88%)	0.11	11 (3%) 39 39	60, 104, 174, 190	0
1	E	284/315 (90%)	-0.03	4 (1%) 75 73	57, 99, 148, 182	0
1	F	284/315 (90%)	-0.38	1 (0%) 92 91	39, 68, 101, 129	0
1	Н	285/315 (90%)	-0.35	1 (0%) 92 91	43, 73, 99, 117	0
All	All	1703/1890 (90%)	-0.20	21 (1%) 79 77	39, 80, 153, 190	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	289	ASP	5.7
1	С	242	SER	3.7
1	F	289	ASP	3.3
1	С	217	GLY	3.2
1	Е	37	ASP	3.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
2	SO4	С	303	5/5	0.48	0.36	172,175,182,185	0
2	SO4	A	301	5/5	0.70	0.39	162,163,167,168	0
2	SO4	Η	304	5/5	0.71	0.31	156,167,172,174	0
2	SO4	Н	301	5/5	0.77	0.29	120,122,130,131	0
2	SO4	A	304	5/5	0.78	0.31	131,131,137,140	0
2	SO4	Н	303	5/5	0.84	0.27	117,119,125,129	0
2	SO4	F	302	5/5	0.86	0.29	124,124,134,136	0
2	SO4	С	302	5/5	0.87	0.36	129,136,144,150	0
2	SO4	F	301	5/5	0.87	0.22	110,118,125,135	0
2	SO4	Η	302	5/5	0.88	0.32	132,133,142,147	0
2	SO4	С	301	5/5	0.89	0.23	126,126,131,131	0
2	SO4	A	303	5/5	0.89	0.25	101,103,111,120	0
2	SO4	В	302	5/5	0.89	0.24	116,117,121,122	0
2	SO4	A	302	5/5	0.90	0.25	104,109,118,119	0
2	SO4	Е	302	5/5	0.90	0.23	114,125,135,137	0
2	SO4	В	301	5/5	0.92	0.18	81,90,95,100	0
2	SO4	Е	301	5/5	0.94	0.18	92,92,97,104	0

6.5 Other polymers (i)

There are no such residues in this entry.

