



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2025 – 04:03 PM EDT

PDB ID : 7I17 / pdb_00007i17
Title : PanDDA analysis group deposition – Main Protease (SARS-CoV-2) in complex with fragment C10 from the F2X-Entry Screen in orthorhombic space group
Authors : Barthel, T.; Benz, L.S.; Wollenhaupt, J.; Weiss, M.S.
Deposited on : 2025-02-03
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div><div>8%</div><div>91%</div><div>8%</div><div>.</div></div>
1	B	306	<div><div>25%</div><div>85%</div><div>10%</div><div>..</div></div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5275 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3C-like proteinase nsp5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	0	16	0
			2725	1702	470	516	37			
1	B	304	Total	C	N	O	S	0	2	0
			2386	1509	405	447	25			

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is 1,3-benzodioxole-5-carbothioamide (CCD ID: TJV) (formula: C₈H₇NO₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	1
			24	16	2	4	2		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Na	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	76	Total	O	0	5
			85	85		
6	B	45	Total	O	0	0
			45	45		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3C-like proteinase nsp5



• Molecule 1: 3C-like proteinase nsp5



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.89Å 99.84Å 103.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.00 – 1.69 46.00 – 1.69	Depositor EDS
% Data completeness (in resolution range)	98.9 (46.00-1.69) 99.0 (46.00-1.69)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 1.69Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.230 , 0.260 0.237 , 0.263	Depositor DCC
R_{free} test set	2101 reflections (2.67%)	wwPDB-VP
Wilson B-factor (Å ²)	24.8	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 25.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.007 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5275	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, DMS, NA, TJV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	3/2789 (0.1%)	1.25	3/3783 (0.1%)
1	B	1.12	4/2438 (0.2%)	1.38	7/3315 (0.2%)
All	All	1.08	7/5227 (0.1%)	1.31	10/7098 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	261	VAL	C-O	6.41	1.31	1.24
1	B	265	CYS	C-O	6.26	1.31	1.24
1	A	27	LEU	C-O	5.78	1.30	1.23
1	B	226	THR	C-O	5.76	1.31	1.23
1	A	39	PRO	C-O	-5.54	1.17	1.23
1	A	184	PRO	C-O	-5.27	1.18	1.23
1	B	260	ALA	C-O	5.14	1.30	1.23

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	263	ASP	CA-CB-CG	11.51	124.11	112.60
1	A	153	ASP	CB-CA-C	7.21	122.02	110.19
1	B	2	GLY	CA-C-O	-5.83	118.43	122.22
1	B	277	ASN	CB-CA-C	5.72	120.70	112.12
1	B	127	GLN	CB-CG-CD	5.59	122.11	112.60
1	B	262	LEU	CA-C-N	5.55	128.50	120.79
1	B	262	LEU	C-N-CA	5.55	128.50	120.79
1	A	148	VAL	CA-C-N	5.25	127.89	122.80
1	A	148	VAL	C-N-CA	5.25	127.89	122.80
1	B	196	THR	CB-CA-C	5.14	117.56	109.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2725	0	2586	15	0
1	B	2386	0	2337	34	0
2	A	1	0	0	0	0
3	A	24	0	0	0	0
4	A	4	0	6	0	0
4	B	4	0	6	3	0
5	B	1	0	0	0	0
6	A	85	0	0	0	0
6	B	45	0	0	6	0
All	All	5275	0	4935	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:SER:HB2	1:B:304:THR:HG21	1.35	1.07
1:A:121:SER:CB	1:B:304:THR:HG21	1.91	1.00
1:B:188:ARG:HD2	6:B:541:HOH:O	1.69	0.91
4:B:402:DMS:H21	6:B:531:HOH:O	1.73	0.89
1:A:121:SER:HB3	1:B:304:THR:CG2	2.02	0.87
1:A:121:SER:CB	1:B:304:THR:CG2	2.52	0.87
1:A:121:SER:HB3	1:B:304:THR:HG23	1.59	0.82
1:B:260:ALA:O	1:B:263:ASP:HB3	1.82	0.79
1:A:4:ARG:NH2	1:B:127:GLN:O	2.17	0.77
1:B:58:LEU:HD22	1:B:82:MET:HE3	1.72	0.69
1:B:188:ARG:CD	6:B:541:HOH:O	2.38	0.60
1:B:266:ALA:O	1:B:270:GLU:HG2	2.03	0.59
1:A:121:SER:HB2	1:B:304:THR:CG2	2.18	0.55
1:A:286:LEU:HD22	6:B:544:HOH:O	2.07	0.55
1:B:62:SER:HG	1:B:64:HIS:CD2	2.27	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:263:ASP:O	1:B:266:ALA:HB3	2.11	0.50
1:B:265:CYS:O	1:B:269:LYS:N	2.20	0.50
1:B:294:PHE:CE2	1:B:298:ARG:HG3	2.47	0.50
1:A:86:VAL:HG23	1:A:179:GLY:HA2	1.94	0.49
1:B:226:THR:OG1	1:B:229:ASP:OD2	2.28	0.49
1:B:225:THR:OG1	1:B:226:THR:N	2.45	0.49
1:B:225:THR:HG22	1:B:262:LEU:O	2.12	0.49
1:B:62:SER:OG	1:B:64:HIS:CD2	2.66	0.48
1:B:6:MET:HB3	4:B:402:DMS:H22	1.94	0.48
1:B:55:GLU:CD	6:B:503:HOH:O	2.57	0.47
1:A:109:GLY:HA2	1:A:200:ILE:HD13	1.97	0.47
1:A:285:ALA:HB3	1:B:285:ALA:HB3	1.96	0.47
1:B:260:ALA:O	1:B:263:ASP:CB	2.60	0.47
1:B:294:PHE:CZ	1:B:298:ARG:HG3	2.49	0.46
4:B:402:DMS:C2	6:B:531:HOH:O	2.46	0.46
1:A:31:TRP:CD2	1:A:95:ASN:HB2	2.51	0.46
1:B:294:PHE:O	1:B:298:ARG:HB2	2.16	0.46
1:B:40:ARG:HB2	1:B:82:MET:HE2	1.98	0.45
1:B:62:SER:OG	1:B:64:HIS:NE2	2.49	0.45
1:B:199:THR:HG21	1:B:239:TYR:CZ	2.52	0.44
1:A:6:MET:HE2	1:B:126:TYR:CE2	2.53	0.44
1:B:276:MET:HA	1:B:279:ARG:CZ	2.48	0.43
1:B:227:LEU:HD12	1:B:227:LEU:HA	1.90	0.42
1:A:113:SER:O	1:A:149:GLY:HA2	2.19	0.42
1:A:101:TYR:HA	1:A:157:VAL:O	2.20	0.41
1:B:113:SER:O	1:B:149:GLY:HA2	2.21	0.41
1:B:225:THR:O	1:B:262:LEU:HD13	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/306 (113%)	336 (97%)	10 (3%)	0	100	100
1	B	308/306 (101%)	297 (96%)	9 (3%)	2 (1%)	22	10
All	All	654/612 (107%)	633 (97%)	19 (3%)	2 (0%)	37	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	276	MET
1	B	301	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/263 (116%)	303 (99%)	3 (1%)	73	64
1	B	267/263 (102%)	253 (95%)	14 (5%)	19	7
All	All	573/526 (109%)	556 (97%)	17 (3%)	40	19

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	LEU
1	A	69	GLN
1	A	154	TYR
1	B	46	SER
1	B	59	ILE
1	B	86	VAL
1	B	121	SER
1	B	125[A]	VAL
1	B	125[B]	VAL
1	B	125[C]	VAL
1	B	125[D]	VAL
1	B	165	MET
1	B	216	ASP
1	B	227	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	279	ARG
1	B	301	SER
1	B	304	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	HIS
1	B	51	ASN
1	B	277	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DMS	B	402	-	3,3,3	0.42	0	3,3,3	0.18	0
4	DMS	A	403	-	3,3,3	0.37	0	3,3,3	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TJV	A	402[D]	-	13,13,13	3.20	4 (30%)	18,18,18	1.58	3 (16%)
3	TJV	A	402[C]	-	13,13,13	3.20	4 (30%)	18,18,18	1.58	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TJV	A	402[D]	-	-	0/4/10/10	0/2/2/2
3	TJV	A	402[C]	-	-	0/4/10/10	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402[C]	TJV	C6-N	9.41	1.42	1.32
3	A	402[D]	TJV	C6-N	9.41	1.42	1.32
3	A	402[C]	TJV	C6-S	-4.50	1.59	1.66
3	A	402[D]	TJV	C6-S	-4.50	1.59	1.66
3	A	402[C]	TJV	C4-C5	3.15	1.44	1.39
3	A	402[D]	TJV	C4-C5	3.15	1.44	1.39
3	A	402[C]	TJV	C1-C	2.53	1.42	1.38
3	A	402[D]	TJV	C1-C	2.53	1.42	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402[C]	TJV	C5-C6-N	4.60	122.86	116.25
3	A	402[D]	TJV	C5-C6-N	4.60	122.86	116.25
3	A	402[C]	TJV	S-C6-N	-2.91	117.62	122.68
3	A	402[D]	TJV	S-C6-N	-2.91	117.62	122.68
3	A	402[C]	TJV	C1-C-C5	2.30	123.26	120.80
3	A	402[D]	TJV	C1-C-C5	2.30	123.26	120.80

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	402	DMS	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	300/306 (98%)	0.51	25 (8%)	19 19	5, 27, 44, 90	16 (5%)
1	B	304/306 (99%)	1.11	76 (25%)	2 2	5, 32, 59, 73	2 (0%)
All	All	604/612 (98%)	0.81	101 (16%)	5 4	5, 29, 55, 90	18 (2%)

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	154	TYR	6.2
1	A	154	TYR	6.0
1	B	275	GLY	5.7
1	B	218	TRP	5.3
1	B	276	MET	5.2
1	B	220	LEU	4.6
1	B	303	VAL	4.6
1	B	266	ALA	4.5
1	B	59	ILE	4.5
1	B	300	CYS	4.4
1	B	232	LEU	4.3
1	A	153	ASP	4.2
1	B	302	GLY	4.1
1	A	67	LEU	4.0
1	B	223	PHE	4.0
1	B	271	LEU	4.0
1	B	269	LYS	4.0
1	B	233	VAL	3.9
1	B	259	ILE	3.8
1	B	257	THR	3.8
1	B	224	THR	3.6
1	B	236	LYS	3.6
1	B	262	LEU	3.6
1	B	227	LEU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	219	PHE	3.4
1	B	225	THR	3.4
1	B	278	GLY	3.4
1	B	294	PHE	3.4
1	B	304	THR	3.4
1	B	64	HIS	3.3
1	B	261	VAL	3.3
1	B	188	ARG	3.3
1	B	280	THR	3.3
1	A	294	PHE	3.2
1	B	237	TYR	3.2
1	B	73	VAL	3.2
1	B	50	LEU	3.2
1	B	301	SER	3.2
1	B	273	GLN	3.2
1	A	73	VAL	3.1
1	A	47	GLU	3.1
1	B	209	TYR	3.1
1	B	226	THR	3.0
1	B	60	ARG	3.0
1	B	274	ASN	2.9
1	B	277	ASN	2.8
1	B	249	ILE	2.8
1	B	260	ALA	2.8
1	B	67	LEU	2.8
1	A	94	ALA	2.8
1	B	72	ASN	2.7
1	B	142	ASN	2.7
1	A	286	LEU	2.7
1	B	75	LEU	2.7
1	B	45	THR	2.7
1	A	64	HIS	2.7
1	B	74	GLN	2.7
1	B	272	LEU	2.7
1	B	62	SER	2.6
1	B	265	CYS	2.6
1	B	279	ARG	2.6
1	B	270	GLU	2.6
1	A	46	SER	2.6
1	A	78	ILE	2.6
1	B	252	PRO	2.6
1	B	24	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	23	GLY	2.5
1	A	155	ASP	2.5
1	B	267	SER	2.5
1	B	213	ILE	2.4
1	B	247	VAL	2.4
1	B	155	ASP	2.4
1	B	230	PHE	2.3
1	B	285	ALA	2.3
1	A	102	LYS	2.3
1	B	46	SER	2.3
1	B	235	MET	2.3
1	B	286	LEU	2.3
1	B	245	ASP	2.3
1	A	79	GLY	2.3
1	B	57	LEU	2.3
1	A	74	GLN	2.2
1	A	222	ARG	2.2
1	B	47	GLU	2.2
1	A	300	CYS	2.2
1	A	49[A]	MET	2.1
1	A	75	LEU	2.1
1	A	297	VAL	2.1
1	B	212	VAL	2.1
1	B	76	ARG	2.1
1	B	58	LEU	2.1
1	B	268	LEU	2.1
1	B	229	ASP	2.1
1	B	298	ARG	2.1
1	A	66	PHE	2.1
1	B	211	ALA	2.0
1	A	89	LEU	2.0
1	B	217	ARG	2.0
1	A	62	SER	2.0
1	A	278	GLY	2.0
1	A	283	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	TJV	A	402[C]	12/12	0.87	0.23	29,35,38,38	12
3	TJV	A	402[D]	12/12	0.87	0.23	29,35,38,38	12
5	NA	B	401	1/1	0.88	0.14	46,46,46,46	0
4	DMS	B	402	4/4	0.94	0.13	39,42,44,45	0
4	DMS	A	403	4/4	0.94	0.14	34,35,40,41	0
2	CL	A	401	1/1	0.98	0.07	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around TJV A 402 (C):

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TJV A 402 (D):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.