



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 20, 2024 – 02:26 PM JST

PDB ID : 8IGZ
Title : Xcc NAMPT Quadruple mutant
Authors : Xu, G.L.; Ming, Z.H.
Deposited on : 2023-02-21
Resolution : 3.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1292 (3.14-3.10)
Clashscore	141614	1389 (3.14-3.10)
Ramachandran outliers	138981	1337 (3.14-3.10)
Sidechain outliers	138945	1337 (3.14-3.10)
RSRZ outliers	127900	1260 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	 2% (Poor fit) 79% (Green) 14% (Yellow) 6% (Grey)
1	B	482	 3% (Poor fit) 77% (Green) 11% (Yellow) 12% (Grey)

2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 6837 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-B cell enhancing factor related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	454	Total	C	N	O	S	0	0	0
			3544	2248	628	660	8			
1	B	423	Total	C	N	O	S	0	0	0
			3293	2093	579	614	7			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP A0A0H2X5R2
A	-12	GLY	-	expression tag	UNP A0A0H2X5R2
A	-11	SER	-	expression tag	UNP A0A0H2X5R2
A	-10	SER	-	expression tag	UNP A0A0H2X5R2
A	-9	HIS	-	expression tag	UNP A0A0H2X5R2
A	-8	HIS	-	expression tag	UNP A0A0H2X5R2
A	-7	HIS	-	expression tag	UNP A0A0H2X5R2
A	-6	HIS	-	expression tag	UNP A0A0H2X5R2
A	-5	HIS	-	expression tag	UNP A0A0H2X5R2
A	-4	HIS	-	expression tag	UNP A0A0H2X5R2
A	-3	SER	-	expression tag	UNP A0A0H2X5R2
A	-2	GLN	-	expression tag	UNP A0A0H2X5R2
A	-1	GLY	-	expression tag	UNP A0A0H2X5R2
A	0	SER	-	expression tag	UNP A0A0H2X5R2
A	175	PHE	HIS	engineered mutation	UNP A0A0H2X5R2
A	224	PHE	ILE	engineered mutation	UNP A0A0H2X5R2
A	291	PHE	VAL	engineered mutation	UNP A0A0H2X5R2
A	332	PHE	ILE	engineered mutation	UNP A0A0H2X5R2
B	-13	MET	-	initiating methionine	UNP A0A0H2X5R2
B	-12	GLY	-	expression tag	UNP A0A0H2X5R2
B	-11	SER	-	expression tag	UNP A0A0H2X5R2
B	-10	SER	-	expression tag	UNP A0A0H2X5R2
B	-9	HIS	-	expression tag	UNP A0A0H2X5R2
B	-8	HIS	-	expression tag	UNP A0A0H2X5R2
B	-7	HIS	-	expression tag	UNP A0A0H2X5R2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	HIS	-	expression tag	UNP A0A0H2X5R2
B	-5	HIS	-	expression tag	UNP A0A0H2X5R2
B	-4	HIS	-	expression tag	UNP A0A0H2X5R2
B	-3	SER	-	expression tag	UNP A0A0H2X5R2
B	-2	GLN	-	expression tag	UNP A0A0H2X5R2
B	-1	GLY	-	expression tag	UNP A0A0H2X5R2
B	0	SER	-	expression tag	UNP A0A0H2X5R2
B	175	PHE	HIS	engineered mutation	UNP A0A0H2X5R2
B	224	PHE	ILE	engineered mutation	UNP A0A0H2X5R2
B	291	PHE	VAL	engineered mutation	UNP A0A0H2X5R2
B	332	PHE	ILE	engineered mutation	UNP A0A0H2X5R2

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pre-B cell enhancing factor related protein

Chain A: 



- Molecule 1: Pre-B cell enhancing factor related protein

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	115.76Å 115.76Å 317.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.72 – 3.11 35.72 – 3.11	Depositor EDS
% Data completeness (in resolution range)	99.9 (35.72-3.11) 100.0 (35.72-3.11)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 3.12Å)	Xtrriage
Refinement program	PHENIX 1.15_3459	Depositor
R, R_{free}	0.213 , 0.255 0.213 , 0.255	Depositor DCC
R_{free} test set	1201 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	31.1	Xtrriage
Anisotropy	0.002	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6837	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/3624	0.48	0/4930
1	B	0.27	0/3364	0.48	0/4574
All	All	0.27	0/6988	0.48	0/9504

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	282	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3544	0	3475	40	0
1	B	3293	0	3236	31	0
All	All	6837	0	6711	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:7:LEU:HD21	1:B:95:LEU:HD11	1.71	0.73
1:B:264:SER:HB2	1:B:302:VAL:HG23	1.72	0.72
1:A:95:LEU:HD13	1:A:131:LEU:HD11	1.74	0.68
1:A:281:GLU:OE2	1:A:282:GLU:HB2	1.94	0.68
1:B:95:LEU:HD13	1:B:131:LEU:HD11	1.78	0.65
1:A:415:ARG:NH1	1:A:422:TYR:OH	2.30	0.64
1:B:265:TYR:HA	1:B:268:TYR:HB3	1.79	0.64
1:A:268:TYR:CE1	1:A:272:ARG:HD3	2.35	0.60
1:A:182:VAL:HG11	1:A:188:ALA:HB2	1.82	0.59
1:A:278:THR:O	1:A:281:GLU:HB3	2.05	0.56
1:B:65:ILE:HD13	1:B:85:ARG:HG2	1.88	0.56
1:A:7:LEU:HD21	1:A:95:LEU:HD11	1.89	0.55
1:B:182:VAL:HG11	1:B:188:ALA:HB2	1.88	0.55
1:A:17:SER:HA	1:A:19:TRP:CZ3	2.43	0.54
1:A:75:HIS:NE2	1:B:223:SER:HB3	2.23	0.53
1:A:444:VAL:O	1:A:451:LEU:N	2.41	0.52
1:A:443:THR:HG22	1:A:451:LEU:HD12	1.91	0.51
1:B:164:ASP:OD2	1:B:319:ASN:HB2	2.11	0.50
1:B:316:HIS:CE1	1:B:324:LYS:HE3	2.47	0.50
1:A:393:ASP:HB3	1:A:408:ARG:HB2	1.94	0.50
1:B:290:VAL:HG23	1:B:329:VAL:HG12	1.92	0.50
1:A:316:HIS:CE1	1:A:324:LYS:HE3	2.47	0.50
1:A:17:SER:HA	1:A:19:TRP:HZ3	1.77	0.50
1:B:300:VAL:HA	1:B:345:ILE:HG12	1.94	0.49
1:B:355:ALA:O	1:B:358:ASN:ND2	2.46	0.49
1:A:71:LEU:HD11	1:B:208:LEU:HD23	1.96	0.48
1:A:247:MET:HG2	1:A:260:VAL:HG12	1.96	0.48
1:A:282:GLU:HA	1:A:285:ALA:HB3	1.95	0.48
1:A:301:ASP:OD1	1:A:348:ARG:NH1	2.36	0.48
1:B:282:GLU:HG2	1:B:286:SER:HB3	1.95	0.48
1:A:136:LEU:HD23	1:B:180:ARG:HD3	1.96	0.48
1:B:412:THR:OG1	1:B:439:ASP:OD1	2.32	0.47
1:B:432:THR:O	1:B:432:THR:OG1	2.33	0.47
1:A:452:ARG:HA	1:A:452:ARG:HD2	1.65	0.46
1:B:263:ASP:O	1:B:264:SER:HB3	2.15	0.46
1:A:265:TYR:CE2	1:A:269:ARG:HD2	2.50	0.46
1:A:333:GLN:HG3	1:A:337:ILE:HD11	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:298:ASP:O	1:B:302:VAL:HG12	2.16	0.46
1:B:310:LEU:HB3	1:B:329:VAL:HG21	1.98	0.46
1:A:4:LEU:HD22	1:B:185:LEU:HD21	1.98	0.45
1:A:115:MET:HE2	1:A:117:ILE:HD11	1.98	0.45
1:B:283:ILE:HG12	1:B:290:VAL:HG11	1.99	0.45
1:B:162:THR:HG22	1:B:323:TYR:CZ	2.52	0.44
1:B:264:SER:HB2	1:B:302:VAL:CG2	2.44	0.44
1:A:273:GLU:O	1:A:273:GLU:HG2	2.17	0.44
1:A:467:ARG:O	1:A:468:LEU:HB2	2.18	0.43
1:A:349:ILE:HG13	1:A:350:THR:N	2.33	0.43
1:A:263:ASP:OD1	1:A:266:ASP:N	2.44	0.43
1:A:137:ARG:HA	1:A:139:TRP:CD1	2.54	0.43
1:A:34:GLU:HG3	1:A:36:ARG:HG2	2.00	0.43
1:A:31:PHE:CE2	1:A:382:CYS:HB2	2.54	0.42
1:A:301:ASP:HB3	1:A:305:GLN:NE2	2.35	0.42
1:B:443:THR:HG22	1:B:451:LEU:HD12	2.02	0.42
1:A:412:THR:OG1	1:A:439:ASP:OD1	2.36	0.42
1:A:11:THR:HA	1:A:133:THR:HB	2.01	0.42
1:A:180:ARG:HD3	1:B:136:LEU:HD23	2.01	0.42
1:A:241:VAL:HG23	1:A:275:TRP:CE2	2.55	0.42
1:B:316:HIS:NE2	1:B:324:LYS:HE3	2.35	0.42
1:B:98:ARG:HD3	1:B:446:GLU:OE2	2.19	0.42
1:A:13:SER:HB3	1:A:77:GLU:HG3	2.00	0.41
1:B:310:LEU:HD22	1:B:329:VAL:HG11	2.02	0.41
1:A:100:ARG:HH22	1:A:439:ASP:CG	2.24	0.41
1:A:151:VAL:HG11	1:A:174:LEU:HD22	2.03	0.41
1:B:212:ARG:HD3	1:B:218:PRO:HA	2.03	0.41
1:B:467:ARG:O	1:B:468:LEU:HB2	2.20	0.41
1:A:316:HIS:NE2	1:A:324:LYS:HE3	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	448/482 (93%)	431 (96%)	17 (4%)	0	100	100
1	B	413/482 (86%)	399 (97%)	13 (3%)	1 (0%)	47	79
All	All	861/964 (89%)	830 (96%)	30 (4%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	264	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/391 (94%)	356 (97%)	11 (3%)	41	70
1	B	342/391 (88%)	333 (97%)	9 (3%)	46	74
All	All	709/782 (91%)	689 (97%)	20 (3%)	43	72

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	42	GLN
1	A	176	ASP
1	A	229	HIS
1	A	274	HIS
1	A	281	GLU
1	A	335	ASP
1	A	357	ASP
1	A	378	PHE
1	A	417	ARG
1	A	452	ARG
1	B	1	MET
1	B	19	TRP
1	B	42	GLN
1	B	129	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	268	TYR
1	B	335	ASP
1	B	357	ASP
1	B	378	PHE
1	B	415	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	316	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	454/482 (94%)	-0.31	9 (1%) 65 45	7, 18, 54, 101	0
1	B	423/482 (87%)	-0.27	15 (3%) 44 23	8, 17, 60, 95	0
All	All	877/964 (90%)	-0.29	24 (2%) 54 31	7, 17, 58, 101	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	282	GLU	4.9
1	B	281	GLU	4.6
1	B	285	ALA	4.4
1	A	406	SER	3.8
1	B	286	SER	3.6
1	A	275	TRP	3.5
1	B	279	LEU	3.4
1	B	268	TYR	3.3
1	B	274	HIS	3.2
1	B	390	ALA	3.2
1	A	277	THR	2.7
1	B	388	ASP	2.6
1	B	395	TYR	2.4
1	B	226	ALA	2.3
1	A	272	ARG	2.3
1	A	387	VAL	2.3
1	B	225	PRO	2.3
1	A	25	GLY	2.2
1	A	428	ASP	2.2
1	B	406	SER	2.1
1	B	283	ILE	2.1
1	A	282	GLU	2.1
1	B	273	GLU	2.1
1	A	237	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.