

Full wwPDB X-ray Structure Validation Report (i)

Oct 29, 2024 – 12:11 AM JST

PDB ID : 9J56

Title: Functional Investigation of the SAM-Dependent Methyltransferases Rdmb in

Anthracycline Biosynthesis

Authors: Yang, Q.Y.; Sang, M.L.; Zhang, W.

Deposited on : 2024-08-11

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	374	81%	10%	9%
1	В	374	78%	11%	12%



2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 5270 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Aclacinomycin 10-hydroxylase RdmB.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	339	Total	С	N	О	S	1.4	0	0
1	Λ	009	2505	1580	443	474	8	14	U	0
1	D	330	Total	С	N	О	S	4	0	0
1	Б	330	2410	1525	423	455	7	4	U	

• Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	S	0	0
	A	1	26	14	6	5	1	0	
2	D	1	Total	С	N	О	S	0	0
	Б	1	26	14	6	5	1	0	

• Molecule 3 is $(7 \{S\},9 \{S\})-7-[(2 \{R\},4 \{S\},5 \{S\},6 \{S\})-4-azanyl-6-methyl-5-oxidanyl-ox an-2-yl]oxy-9-ethyl-4-methoxy-6,9,11-tris(oxidanyl)-8,10-dihydro-7 {H}-tetracene-5,12-d ione (three-letter code: A1L3V) (formula: <math>C_{27}H_{31}NO_9$) (labeled as "Ligand of Interest" by



depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total 37	_	_	0	0
3	В	1	Total 37	_	_	0	0

• Molecule 4 is water.


\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	136	Total O 136 136	0	0
4	В	93	Total O 93 93	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aclacinomycin 10-hydroxylase RdmB





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants	80.00Å 80.00Å 234.20Å	Donogitor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.72 - 2.10	Depositor
Resolution (A)	44.72 - 2.10	EDS
% Data completeness	95.5 (44.72-2.10)	Depositor
(in resolution range)	95.5 (44.72-2.10)	EDS
R_{merge}	0.02	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.08 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.20.1	Depositor
D.D.	0.214 , 0.259	Depositor
R, R_{free}	0.227 , 0.263	DCC
R_{free} test set	49767 reflections (4.03%)	wwPDB-VP
Wilson B-factor (Å ²)	40.6	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 43.0	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5270	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: A1L3V, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.38	0/2549	0.65	0/3473	
1	В	0.35	0/2451	0.62	0/3340	
All	All	0.36	0/5000	0.63	0/6813	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2505	0	2458	27	0
1	В	2410	0	2352	26	0
2	A	26	0	19	0	0
2	В	26	0	18	0	0
3	A	37	0	0	3	0
3	В	37	0	0	0	0
4	A	136	0	0	4	0
4	В	93	0	0	1	0
All	All	5270	0	4847	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

	A	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:A:165:CYS:SG	4:A:626:HOH:O	2.38	0.80
1:B:286:LEU:HG	1:B:350:ILE:HG12	1.68	0.75
1:A:176:ASP:OD1	1:A:203:ARG:NH1	2.23	0.72
1:B:92:THR:HG23	1:B:95:GLY:H	1.56	0.71
1:A:208:ARG:NH2	4:A:501:HOH:O	2.22	0.69
1:A:183:VAL:HG12	1:A:207:LEU:HD21	1.77	0.66
1:A:204:ALA:HB3	1:A:207:LEU:HD12	1.83	0.60
1:B:77:GLY:O	1:B:92:THR:HG21	2.04	0.57
1:B:92:THR:CG2	1:B:95:GLY:H	2.18	0.56
1:B:258:LEU:O	1:B:317:ARG:NH2	2.40	0.54
1:A:140:ARG:NH1	4:A:502:HOH:O	2.24	0.53
1:A:159:PHE:HZ	3:A:402:A1L3V:C32	2.24	0.51
1:A:75:VAL:HG11	1:B:299:PHE:HA	1.93	0.51
1:A:12:PRO:HG3	1:B:96:MET:HB2	1.93	0.51
1:A:183:VAL:HG13	1:A:250:ASP:OD1	2.11	0.50
1:A:204:ALA:HB1	1:A:206:HIS:CE1	2.46	0.50
1:A:68:ARG:NH2	1:B:309:LEU:HB2	2.28	0.49
1:B:40:ARG:HG2	1:B:43:ASP:HB2	1.95	0.48
1:A:127:LEU:HD12	1:B:35:VAL:HG13	1.96	0.48
1:A:186:VAL:HG23	1:A:207:LEU:HD22	1.96	0.47
1:B:204:ALA:HB1	1:B:206:HIS:CE1	2.50	0.47
1:A:331:LEU:HB3	1:A:353:PHE:HB3	1.97	0.47
1:A:11:GLU:N	1:A:12:PRO:HD3	2.30	0.46
1:B:331:LEU:HB3	1:B:353:PHE:HB3	1.97	0.46
1:B:274:VAL:HG22	1:B:331:LEU:HD21	1.97	0.46
1:B:219:GLU:OE1	1:B:222:ARG:NH1	2.50	0.45
1:A:187:LEU:HD11	1:A:212:VAL:HG23	1.99	0.45
1:A:292:GLU:H	1:A:292:GLU:CD	2.21	0.44
1:A:42:VAL:HG13	1:A:91:PRO:HG3	1.99	0.44
1:B:42:VAL:HG13	1:B:91:PRO:HG3	1.99	0.44
1:A:149:LEU:HD23	1:A:155:LEU:HD23	2.00	0.44
1:A:159:PHE:CZ	3:A:402:A1L3V:C32	3.01	0.44
1:A:309:LEU:HB2	1:B:68:ARG:NH2	2.32	0.43
1:B:152:ASP:OD2	1:B:155:LEU:HD13	2.18	0.43
1:A:282:ARG:HD2	1:A:354:THR:HG22	2.00	0.42
1:B:59:THR:O	1:B:60:ASP:HB3	2.18	0.42
1:A:178:TYR:CE1	1:A:284:LEU:HD21	2.54	0.42
1:B:282:ARG:HD3	1:B:354:THR:HG22	2.01	0.42
1:B:255:SER:HA	1:B:286:LEU:HB2	2.02	0.42
1:B:208:ARG:NH2	4:B:507:HOH:O	2.53	0.42



I 'omtamalod	tmom	mmonia	maaa
Continued	110111	DIEUIUU	DUJUE
00,000,000	.,	p	p = 9 =

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance } (ext{Å}) \end{array}$	Clash overlap (Å)
1:B:330:GLY:O	1:B:356:VAL:HG22	2.19	0.42
1:B:149:LEU:HD23	1:B:155:LEU:HB3	2.02	0.41
1:B:208:ARG:HE	1:B:208:ARG:HB3	1.80	0.41
1:A:71:ARG:NH1	4:A:516:HOH:O	2.52	0.41
1:A:282:ARG:CD	1:A:354:THR:HG22	2.51	0.41
1:B:71:ARG:O	1:B:75:VAL:HG23	2.21	0.41
3:A:402:A1L3V:O37	3:A:402:A1L3V:O06	2.38	0.40
1:B:315:ARG:HE	1:B:315:ARG:HB2	1.72	0.40
1:A:173:ALA:HB3	1:A:174:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	rs Percentile	
1	A	333/374 (89%)	327 (98%)	6 (2%)	0	100	100
1	В	320/374 (86%)	311 (97%)	9 (3%)	0	100	100
All	All	653/748 (87%)	638 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	250/287~(87%)	246 (98%)	4 (2%)	58 65		
1	В	238/287 (83%)	235 (99%)	3 (1%)	65 72		
All	All	488/574 (85%)	481 (99%)	7 (1%)	62 70		

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	241	PHE
1	A	250	ASP
1	A	287	ASP
1	A	319	ARG
1	В	117	SER
1	В	166	ASP
1	В	241	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Chain	Res	s Link	Bond lengths			Bond angles			
		nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2	
2	SAH	A	401	-	24,28,28	1.76	6 (25%)	25,40,40	2.12	6 (24%)
2	SAH	В	401	-	24,28,28	1.90	6 (25%)	25,40,40	2.07	6 (24%)
3	A1L3V	В	402	-	40,41,41	1.71	12 (30%)	55,64,64	1.28	7 (12%)
3	A1L3V	A	402	-	40,41,41	1.89	11 (27%)	55,64,64	1.61	10 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	A	401	-	-	3/11/31/31	0/3/3/3
2	SAH	В	401	-	-	3/11/31/31	0/3/3/3
3	A1L3V	В	402	-	-	8/9/55/55	0/5/5/5
3	A1L3V	A	402	-	-	5/9/55/55	0/5/5/5

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
2	В	401	SAH	O4'-C4'	-6.18	1.31	1.45
2	A	401	SAH	O4'-C4'	-5.37	1.33	1.45
3	A	402	A1L3V	O01-C02	-4.58	1.14	1.22
3	A	402	A1L3V	C36-C29	-4.56	1.34	1.41
3	В	402	A1L3V	C24-C25	4.36	1.56	1.51
3	A	402	A1L3V	O37-C28	-4.16	1.15	1.22
3	A	402	A1L3V	C04-C03	-3.19	1.33	1.41
3	В	402	A1L3V	C18-C16	-3.17	1.46	1.53
3	A	402	A1L3V	O23-C20	-3.09	1.38	1.44
2	A	401	SAH	C6-N6	3.07	1.45	1.34
2	В	401	SAH	C6-N6	3.01	1.45	1.34
3	A	402	A1L3V	O11-C10	2.96	1.49	1.42
3	В	402	A1L3V	C29-C28	2.94	1.54	1.47
2	В	401	SAH	O2'-C2'	2.88	1.49	1.43
3	A	402	A1L3V	C18-C16	-2.80	1.47	1.53
2	В	401	SAH	O3'-C3'	-2.76	1.36	1.43
3	В	402	A1L3V	O11-C10	2.76	1.49	1.42
3	В	402	A1L3V	O01-C02	-2.69	1.17	1.22
2	В	401	SAH	C5-C4	-2.61	1.34	1.40



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	A	402	A1L3V	C03-C26	-2.56	1.36	1.41
3	В	402	A1L3V	C36-C02	2.51	1.53	1.48
3	В	402	A1L3V	O37-C28	-2.50	1.18	1.22
3	В	402	A1L3V	C04-C28	2.50	1.53	1.47
3	В	402	A1L3V	O23-C20	-2.45	1.39	1.44
2	A	401	SAH	C2-N3	2.41	1.36	1.32
2	A	401	SAH	O3'-C3'	-2.38	1.37	1.43
2	A	401	SAH	O2'-C2'	2.37	1.48	1.43
2	A	401	SAH	C5-C4	-2.35	1.34	1.40
3	A	402	A1L3V	C07-C25	-2.30	1.35	1.39
3	В	402	A1L3V	C03-C02	2.28	1.53	1.47
3	В	402	A1L3V	C36-C29	-2.18	1.38	1.41
2	В	401	SAH	C2-N3	2.12	1.35	1.32
3	A	402	A1L3V	C29-C28	2.12	1.52	1.47
3	A	402	A1L3V	C14-C16	-2.07	1.49	1.53
3	В	402	A1L3V	O09-C10	-2.05	1.36	1.41

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	401	SAH	N3-C2-N1	-5.73	119.73	128.68
2	A	401	SAH	N3-C2-N1	-5.16	120.61	128.68
2	A	401	SAH	C5-C6-N6	4.71	127.51	120.35
2	В	401	SAH	C5-C6-N6	4.28	126.86	120.35
2	A	401	SAH	O4'-C1'-C2'	-4.21	100.77	106.93
3	A	402	A1L3V	C21-C20-C24	-4.15	105.31	111.67
2	A	401	SAH	CB-CG-SD	-4.06	104.21	113.31
2	В	401	SAH	CB-CG-SD	-4.04	104.25	113.31
3	A	402	A1L3V	C24-C25-C26	-3.98	111.83	119.23
3	В	402	A1L3V	O31-C30-C29	3.64	120.92	115.85
3	A	402	A1L3V	O11-C12-C14	3.47	115.75	109.52
3	A	402	A1L3V	O09-C10-C18	3.43	115.03	108.41
3	A	402	A1L3V	O15-C14-C12	3.41	117.23	109.67
3	В	402	A1L3V	O31-C30-C33	-3.12	119.03	124.37
2	В	401	SAH	O4'-C1'-C2'	-3.01	102.52	106.93
2	A	401	SAH	N6-C6-N1	-2.95	112.44	118.57
3	В	402	A1L3V	C32-O31-C30	-2.95	113.08	117.53
2	В	401	SAH	N6-C6-N1	-2.93	112.49	118.57
3	A	402	A1L3V	O37-C28-C04	-2.85	116.82	121.43
2	A	401	SAH	C5'-SD-CG	2.79	110.63	102.27
2	В	401	SAH	C5'-SD-CG	2.75	110.50	102.27
3	A	402	A1L3V	O31-C30-C33	-2.68	119.77	124.37



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	402	A1L3V	C18-C16-C14	2.62	113.99	110.04
3	A	402	A1L3V	O31-C30-C29	2.56	119.41	115.85
3	В	402	A1L3V	C10-O11-C12	-2.44	107.26	113.84
3	В	402	A1L3V	O09-C10-C18	2.43	113.09	108.41
3	В	402	A1L3V	C18-C16-C14	2.32	113.54	110.04
3	A	402	A1L3V	C26-C25-C07	2.26	122.67	119.17
3	В	402	A1L3V	C05-C07-C08	2.02	121.75	118.59

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	SAH	N-CA-CB-CG
2	A	401	SAH	C-CA-CB-CG
2	В	401	SAH	N-CA-CB-CG
2	В	401	SAH	C-CA-CB-CG
3	A	402	A1L3V	O11-C10-O09-C08
3	A	402	A1L3V	C18-C10-O09-C08
3	A	402	A1L3V	C19-C20-C21-C22
3	A	402	A1L3V	O23-C20-C21-C22
3	В	402	A1L3V	C18-C10-O09-C08
3	В	402	A1L3V	C19-C20-C21-C22
3	В	402	A1L3V	O23-C20-C21-C22
3	В	402	A1L3V	C24-C20-C21-C22
3	A	402	A1L3V	C24-C20-C21-C22
3	В	402	A1L3V	C33-C30-O31-C32
3	В	402	A1L3V	C29-C30-O31-C32
2	В	401	SAH	C4'-C5'-SD-CG
3	В	402	A1L3V	C19-C08-O09-C10
2	A	401	SAH	C4'-C5'-SD-CG
3	В	402	A1L3V	C07-C08-O09-C10

There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	A1L3V	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will




also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	339/374 (90%)	0.15	11 (3%) 50 52	24, 41, 61, 73	4 (1%)
1	В	330/374 (88%)	0.49	20 (6%) 28 30	30, 50, 70, 85	1 (0%)
All	All	669/748 (89%)	0.32	31 (4%) 38 40	24, 45, 67, 85	5 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	50	ASP	4.5
1	В	165	CYS	4.2
1	В	55	LEU	4.1
1	В	88	PRO	3.9
1	В	47	ALA	3.5
1	В	172	GLU	3.4
1	В	9	PRO	3.3
1	В	46	LEU	3.2
1	В	54	GLY	3.0
1	A	88	PRO	2.9
1	A	293	GLY	2.9
1	В	81	GLY	2.9
1	A	178	TYR	2.8
1	В	44	HIS	2.8
1	В	53	ALA	2.7
1	В	166	ASP	2.7
1	A	292	GLU	2.7
1	A	11	GLU	2.6
1	В	45	LEU	2.6
1	В	342	THR	2.6
1	A	48	GLY	2.5
1	A	207	LEU	2.5
1	В	357	SER	2.5
1	В	93	ARG	2.4



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	176	ASP	2.4
1	В	41	LEU	2.4
1	В	99	ALA	2.3
1	A	53	ALA	2.3
1	A	82	GLY	2.2
1	A	14	ASP	2.1
1	В	42	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	A1L3V	В	402	37/37	0.71	0.17	20,20,20,20	0
3	A1L3V	A	402	37/37	0.77	0.16	20,20,20,20	0
2	SAH	В	401	26/26	0.93	0.09	44,46,51,54	0
2	SAH	A	401	26/26	0.96	0.06	33,37,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














Electron density around SAH B 401:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SAH A 401:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

