



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:27 AM UTC

PDB ID : 9JZE / pdb_00009jze
Title : PfDXR - Mn²⁺ - NADPH - MAMK431 quaternary complex
Authors : Takada, S.; Sakamoto, Y.; Tanaka, N.
Deposited on : 2024-10-14
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	
1	B	488	

2 Entry composition i


There are 6 unique types of molecules in this entry. The entry contains 7070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-deoxy-D-xylulose 5-phosphate reductoisomerase, apicoplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	407	Total	C	N	O	S	0	0	0
			3245	2081	531	613	20			
1	B	407	Total	C	N	O	S	0	0	0
			3245	2081	531	613	20			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mn 1 1	0	0
3	B	1	Total Mn 1 1	0	0

- Molecule 4 is [(1 {S})-4-[3-(4-methoxyphenyl)propyl-oxidanyl-amino]-4-oxidanylidene-1-phenyl-butyl]phosphonic acid (CCD ID: A1L45) (formula: C₂₀H₂₆NO₆P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 28 20 1 6 1	0	0
4	B	1	Total C N O P 28 20 1 6 1	0	0

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	3	Total Ca 3 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	198	Total O 198 198	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	225	Total 225	O 225	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.56Å 77.27Å 111.43Å 90.00° 92.46° 90.00°	Depositor
Resolution (Å)	63.48 – 1.84 63.48 – 1.84	Depositor EDS
% Data completeness (in resolution range)	99.7 (63.48-1.84) 99.9 (63.48-1.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 1.84Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.167 , 0.200 0.179 , 0.208	Depositor DCC
R_{free} test set	3800 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtrriage
Anisotropy	0.241	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7070	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, A1L45, MN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/3303	1.13	13/4457 (0.3%)
1	B	0.64	0/3303	1.11	3/4457 (0.1%)
All	All	0.63	0/6606	1.12	16/8914 (0.2%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	286	THR	CA-CB-OG1	-7.07	99.00	109.60
1	A	275	GLN	N-CA-CB	-6.73	99.85	110.28
1	A	424	GLU	CB-CA-C	-5.86	101.44	110.81
1	A	234	HIS	CA-CB-CG	-5.60	108.20	113.80
1	A	399	PHE	CA-CB-CG	-5.51	108.29	113.80
1	A	137	ASP	CA-CB-CG	5.49	118.09	112.60
1	A	305	ASP	CA-CB-CG	5.42	118.02	112.60
1	A	213	PHE	CB-CA-C	-5.36	101.89	110.79
1	A	358	PRO	N-CA-CB	5.35	108.08	103.27
1	B	458	GLU	N-CA-CB	5.29	118.44	110.30
1	A	424	GLU	N-CA-CB	5.23	117.74	109.94
1	A	167	GLU	N-CA-CB	5.19	117.93	110.20
1	A	273	PRO	N-CA-C	5.17	120.39	114.03
1	B	234	HIS	CA-CB-CG	-5.15	108.65	113.80
1	A	200	VAL	N-CA-CB	-5.14	104.99	111.46
1	B	104	GLU	CB-CA-C	-5.10	101.38	109.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3245	0	3296	39	0
1	B	3245	0	3296	34	0
2	A	48	0	26	6	0
2	B	48	0	26	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	28	0	0	0	0
4	B	28	0	0	0	0
5	B	3	0	0	0	0
6	A	198	0	0	13	0
6	B	225	0	0	12	0
All	All	7070	0	6644	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328:ASN:HB2	6:A:737:HOH:O	1.40	1.19
1:B:372:ASP:HB3	6:B:747:HOH:O	1.67	0.95
1:B:480:ASP:HB2	6:B:653:HOH:O	1.66	0.94
1:A:217:LYS:HE2	1:A:458:GLU:O	1.74	0.88
1:A:95:ASN:HB3	6:A:772:HOH:O	1.77	0.85
1:A:480:ASP:HB2	6:A:628:HOH:O	1.76	0.84
1:A:143:GLU:O	1:A:147:LEU:HD23	1.87	0.74
1:B:217:LYS:HE3	6:B:703:HOH:O	1.88	0.72
1:B:449:GLU:HG2	6:B:780:HOH:O	1.92	0.70
1:A:186:GLY:HA2	6:A:626:HOH:O	1.93	0.68
1:B:449:GLU:CG	6:B:780:HOH:O	2.43	0.67
1:B:131:GLU:HB2	6:B:708:HOH:O	1.95	0.66
1:B:480:ASP:CB	6:B:653:HOH:O	2.29	0.66
1:A:141:TYR:CE2	1:A:145:LYS:HD2	2.31	0.65
1:A:122:TYR:OH	1:A:126:ARG:NH2	2.31	0.61
1:B:254:ASP:O	1:B:255:ASN:HB2	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:403:LYS:HE3	1:B:407:GLN:NE2	2.20	0.56
1:B:411:LYS:HE2	1:B:415:TYR:CE2	2.42	0.55
1:B:279:MET:HE1	1:B:438:PHE:CZ	2.44	0.53
1:B:459:ASN:OD1	1:B:462:ASP:OD2	2.27	0.53
1:A:184:PHE:HA	6:A:760:HOH:O	2.09	0.52
1:B:372:ASP:CB	6:B:747:HOH:O	2.41	0.52
2:A:501:NDP:H42N	6:A:637:HOH:O	2.08	0.52
1:B:135:ILE:HD11	1:B:141:TYR:HA	1.91	0.52
1:B:166:LYS:HE2	1:B:192:TYR:CZ	2.46	0.51
1:A:106:VAL:HG23	1:A:107:PHE:CD2	2.46	0.51
1:A:113:TYR:CE1	2:A:501:NDP:H2A	2.46	0.50
1:A:480:ASP:CB	6:A:628:HOH:O	2.48	0.50
1:A:138:LYS:HD3	1:A:160:CYS:SG	2.53	0.49
1:A:221:ILE:N	1:A:221:ILE:HD13	2.28	0.49
1:B:217:LYS:NZ	1:B:458:GLU:O	2.27	0.49
1:B:230:VAL:O	1:B:360:MET:HE1	2.13	0.48
1:B:135:ILE:CD1	1:B:141:TYR:HA	2.43	0.48
1:A:115:ASN:HB3	2:A:501:NDP:O3X	2.14	0.47
1:B:216:LYS:NZ	1:B:254:ASP:OD1	2.39	0.47
1:A:204:ASN:N	6:A:608:HOH:O	2.46	0.47
1:B:393:LYS:HG2	6:B:681:HOH:O	2.13	0.47
1:B:115:ASN:HD22	1:B:136:HIS:HB3	1.80	0.47
1:B:403:LYS:HE3	1:B:407:GLN:HE21	1.80	0.47
1:A:116:LYS:HA	1:A:135:ILE:HD11	1.97	0.46
1:A:115:ASN:HB3	2:A:501:NDP:P2B	2.55	0.46
1:B:299:GLY:N	6:B:601:HOH:O	2.44	0.46
2:A:501:NDP:C5N	6:A:637:HOH:O	2.64	0.46
1:B:202:LEU:HD23	1:B:208:ILE:HD11	1.98	0.46
1:A:138:LYS:HA	1:A:160:CYS:SG	2.56	0.46
1:B:191:MET:HE3	1:B:192:TYR:CE2	2.51	0.45
1:B:461:GLU:O	1:B:465:LYS:HG3	2.16	0.45
1:A:203:ALA:O	2:A:501:NDP:H1D	2.17	0.45
1:A:340:ILE:HD12	1:A:391:PHE:HZ	1.81	0.45
1:B:120:GLU:H	1:B:120:GLU:CD	2.25	0.45
1:A:283:LYS:HE3	6:A:783:HOH:O	2.16	0.45
1:A:197:ASN:HA	1:A:225:ALA:HB2	1.98	0.44
1:A:106:VAL:CG2	1:A:107:PHE:CD2	3.01	0.43
1:A:191:MET:HE2	1:A:214:PHE:CE2	2.53	0.43
1:B:135:ILE:HD11	1:B:141:TYR:HB2	1.98	0.43
1:A:76:LYS:O	1:A:76:LYS:HD3	2.19	0.43
6:A:769:HOH:O	1:B:375:LYS:HE2	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:299:GLY:HA3	6:B:601:HOH:O	2.19	0.43
1:A:135:ILE:O	1:A:160:CYS:HA	2.19	0.43
1:A:136:HIS:ND1	1:A:161:GLY:HA2	2.34	0.43
1:A:141:TYR:CD1	1:A:160:CYS:HB2	2.53	0.43
1:A:191:MET:HG3	1:A:218:LEU:HD21	2.01	0.42
1:A:191:MET:HE2	1:A:214:PHE:CZ	2.55	0.42
1:B:415:TYR:N	1:B:416:PRO:CD	2.82	0.42
1:A:187:LEU:HD23	1:A:464:MET:HE1	2.02	0.42
1:A:349:LYS:HA	1:A:349:LYS:HD3	1.86	0.42
1:B:119:ASN:ND2	1:B:147:LEU:HD11	2.35	0.41
1:A:259:ILE:HD12	6:A:662:HOH:O	2.21	0.41
1:A:203:ALA:HB3	6:A:608:HOH:O	2.19	0.41
1:A:131:GLU:OE1	1:A:131:GLU:N	2.53	0.41
1:A:152:LYS:O	1:A:153:ASP:HB2	2.21	0.41
1:B:355:MET:HE2	6:B:616:HOH:O	2.21	0.40
1:B:480:ASP:O	1:B:484:LYS:HD2	2.21	0.40
1:A:135:ILE:HD13	1:A:140:VAL:HG23	2.04	0.40
1:A:141:TYR:CE1	1:A:160:CYS:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/488 (83%)	392 (97%)	11 (3%)	0	100	100
1	B	403/488 (83%)	391 (97%)	12 (3%)	0	100	100
All	All	806/976 (83%)	783 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/449 (83%)	365 (98%)	9 (2%)	43	25
1	B	374/449 (83%)	367 (98%)	7 (2%)	50	34
All	All	748/898 (83%)	732 (98%)	16 (2%)	47	30

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	LYS
1	A	131	GLU
1	A	167	GLU
1	A	210	SER
1	A	218	LEU
1	A	221	ILE
1	A	359	ASP
1	A	458	GLU
1	A	469	GLN
1	B	116	LYS
1	B	135	ILE
1	B	252	LEU
1	B	387	SER
1	B	455	LYS
1	B	459	ASN
1	B	484	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	105	ASN
1	A	108	ASN
1	A	115	ASN
1	A	150	ASN
1	A	255	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	260	ASN
1	A	284	ASN
1	A	361	GLN
1	A	407	GLN
1	A	446	GLN
1	A	452	ASN
1	A	469	GLN
1	B	105	ASN
1	B	115	ASN
1	B	119	ASN
1	B	172	ASN
1	B	253	GLN
1	B	255	ASN
1	B	261	ASN
1	B	361	GLN
1	B	377	ASN
1	B	407	GLN
1	B	428	ASN
1	B	432	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 5 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NDP	A	501	-	51,52,52	0.77	3 (5%)	71,80,80	0.86	3 (4%)
2	NDP	B	501	-	51,52,52	1.15	2 (3%)	71,80,80	0.88	2 (2%)
4	A1L45	B	503	3	28,29,29	2.02	5 (17%)	30,39,39	1.22	3 (10%)
4	A1L45	A	503	3	28,29,29	1.64	4 (14%)	30,39,39	1.39	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	501	-	-	12/34/77/77	0/5/5/5
2	NDP	B	501	-	-	4/34/77/77	0/5/5/5
4	A1L45	B	503	3	-	7/27/27/27	0/2/2/2
4	A1L45	A	503	3	-	5/27/27/27	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	A1L45	P1-C7	6.05	1.90	1.81
4	B	503	A1L45	P1-O5	-5.90	1.45	1.54
4	B	503	A1L45	O1-N1	-5.81	1.35	1.40
2	B	501	NDP	P2B-O2B	4.90	1.68	1.59
2	B	501	NDP	PN-O3	4.57	1.64	1.59
2	A	501	NDP	PN-O3	2.72	1.62	1.59
4	A	503	A1L45	P1-O5	-2.69	1.50	1.54
2	A	501	NDP	P2B-O2B	2.53	1.64	1.59
4	B	503	A1L45	P1-O3	2.41	1.53	1.49
4	A	503	A1L45	C5-C4	2.26	1.43	1.38
4	A	503	A1L45	P1-O3	2.24	1.53	1.49
4	B	503	A1L45	C5-C6	2.22	1.42	1.38
4	B	503	A1L45	C18-C17	2.12	1.42	1.38
2	A	501	NDP	PA-O3	2.00	1.61	1.59

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	A1L45	C12-C11-N1	3.85	119.04	111.11
2	A	501	NDP	P2B-O2B-C2B	-3.28	114.67	123.43
4	B	503	A1L45	C11-C12-C13	-2.83	105.62	113.05
4	A	503	A1L45	O5-P1-O3	-2.65	106.80	113.45
2	A	501	NDP	C3B-C2B-C1B	-2.39	98.23	102.81
2	B	501	NDP	O3X-P2B-O2X	2.33	116.53	107.80
4	A	503	A1L45	C2-C1-C7	-2.31	116.94	120.73
2	B	501	NDP	C3B-C2B-C1B	-2.27	98.46	102.81
4	A	503	A1L45	C20-O6-C17	2.17	122.16	117.50
4	A	503	A1L45	C8-C9-C10	-2.12	110.60	113.39
2	A	501	NDP	C3N-C7N-N7N	2.08	121.37	117.67
4	B	503	A1L45	C3-C4-C5	2.03	122.64	119.87
4	B	503	A1L45	C13-C14-C15	-2.00	116.07	121.18

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	NDP	C5B-O5B-PA-O2A
2	A	501	NDP	C5B-O5B-PA-O3
2	A	501	NDP	PN-O3-PA-O5B
2	A	501	NDP	O4B-C4B-C5B-O5B
2	A	501	NDP	C5D-O5D-PN-O3
2	A	501	NDP	C5D-O5D-PN-O1N
2	A	501	NDP	C5D-O5D-PN-O2N
4	A	503	A1L45	C16-C17-O6-C20
4	A	503	A1L45	C18-C17-O6-C20
4	B	503	A1L45	C16-C17-O6-C20
4	B	503	A1L45	C18-C17-O6-C20
2	A	501	NDP	O4D-C1D-N1N-C6N
2	A	501	NDP	C3B-C4B-C5B-O5B
2	B	501	NDP	O4D-C1D-N1N-C6N
4	A	503	A1L45	C11-C12-C13-C14
2	A	501	NDP	PA-O3-PN-O1N
2	B	501	NDP	PN-O3-PA-O1A
4	B	503	A1L45	C12-C13-C14-C19
4	B	503	A1L45	C12-C13-C14-C15
2	A	501	NDP	C1B-C2B-O2B-P2B
2	B	501	NDP	C1B-C2B-O2B-P2B
4	A	503	A1L45	N1-C10-C9-C8
4	B	503	A1L45	N1-C10-C9-C8
4	B	503	A1L45	C12-C11-N1-O1
2	B	501	NDP	PN-O3-PA-O2A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	503	A1L45	O2-C10-N1-O1
4	B	503	A1L45	O2-C10-N1-O1
2	A	501	NDP	PA-O3-PN-O2N



There are no ring outliers.


1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	NDP	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/488 (83%)	0.39	46 (11%) 10 10	14, 27, 82, 107	0
1	B	407/488 (83%)	-0.01	11 (2%) 56 61	14, 24, 48, 86	0
All	All	814/976 (83%)	0.19	57 (7%) 22 23	14, 26, 71, 107	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	184	PHE	4.8
1	A	122	TYR	4.6
1	A	118	VAL	4.5
1	A	144	LEU	4.4
1	A	135	ILE	4.3
1	A	147	LEU	4.3
1	B	485	HIS	4.2
1	B	188	TYR	4.1
1	A	183	SER	4.0
1	A	113	TYR	3.9
1	A	140	VAL	3.9
1	A	151	ILE	3.9
1	A	141	TYR	3.8
1	A	160	CYS	3.7
1	A	161	GLY	3.7
1	A	148	VAL	3.7
1	A	114	VAL	3.6
1	A	154	TYR	3.5
1	A	192	TYR	3.5
1	A	158	ILE	3.5
1	B	486	ASN	3.4
1	A	157	ILE	3.3
1	A	185	GLN	3.3
1	B	298	MET	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	159	LEU	3.2
1	B	297	LYS	3.0
1	A	134	CYS	3.0
1	A	136	HIS	2.9
1	A	182	ASP	2.8
1	A	133	LEU	2.8
1	A	181	ILE	2.7
1	A	485	HIS	2.7
1	A	188	TYR	2.6
1	A	129	LEU	2.6
1	A	76	LYS	2.6
1	A	292	LYS	2.5
1	A	119	ASN	2.4
1	A	155	LYS	2.4
1	B	76	LYS	2.4
1	B	299	GLY	2.4
1	A	186	GLY	2.3
1	A	213	PHE	2.3
1	A	115	ASN	2.3
1	B	484	LYS	2.3
1	A	165	MET	2.2
1	B	449	GLU	2.2
1	A	168	ILE	2.2
1	A	120	GLU	2.2
1	A	469	GLN	2.2
1	A	85	SER	2.2
1	B	147	LEU	2.2
1	A	172	ASN	2.1
1	A	116	LYS	2.1
1	A	152	LYS	2.1
1	A	145	LYS	2.1
1	A	125	ALA	2.1
1	B	459	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NDP	A	501	48/48	0.79	0.19	42,56,68,80	48
2	NDP	B	501	48/48	0.92	0.09	24,31,39,41	0
4	A1L45	A	503	28/28	0.96	0.12	20,27,69,88	0
4	A1L45	B	503	28/28	0.97	0.10	17,23,70,76	0
5	CA	B	505	1/1	0.98	0.10	29,29,29,29	0
5	CA	B	504	1/1	0.99	0.06	20,20,20,20	0
3	MN	B	502	1/1	0.99	0.02	16,16,16,16	0
5	CA	B	506	1/1	0.99	0.07	31,31,31,31	0
3	MN	A	502	1/1	1.00	0.01	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1L45 A 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1L45 B 503:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MN B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.