

wwPDB X-ray Structure Validation Summary Report (i)

Feb 10, 2025 - 01:26 PM JST

PDB ID	:	9K2D
Title	:	Structure of ClpP from Staphylococcus aureus in complex with ZY39
Authors	:	Wei, B.Y.; Wang, P.Y.; Wu, W.; Zhang, T.; Yang, CG.
Deposited on		
Resolution	:	1.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{l} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	164625	1356 (1.98-1.98)
Clashscore	180529	1437 (1.98-1.98)
Ramachandran outliers	177936	1426 (1.98-1.98)
Sidechain outliers	177891	1426 (1.98-1.98)
RSRZ outliers	164620	1356 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	201	6% 	• 8%
1	В	201	<u>5%</u> 92%	• 5%
1	С	201	2% 93%	• 5%
1	D	201	4%	• 8%
1	Е	201	5% 91%	• 7%
1	F	201	<u>8%</u> 90%	• 8%



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	G	201	4% 90%	• 8%
1	Н	201	90%	• 7%
1	Ι	201	88%	• 9%
1	J	201	5% 87%	• 10%
1	Κ	201	5% 89%	• 9%
1	L	201	4% 93%	• 5%
1	М	201	2% 91%	• 6%
1	Ν	201	4% 89%	• 10%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 21643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	184	Total	С	Ν	0	S	0	1	0
	11	104	1402	885	237	274	6	0	1	0
1	В	190	Total	С	Ν	Ο	\mathbf{S}	0	0	0
-	D	100	1433	901	242	284	6	Ŭ	0	0
1	С	191	Total	С	Ν	Ο	\mathbf{S}	0	1	0
		101	1447	912	245	284	6	Ŭ	-	
1	D	184	Total	С	Ν	Ο	\mathbf{S}	0	0	0
1		101	1398	881	237	274	6	0	0	0
1	Е	187	Total	С	Ν	Ο	S	0	0	0
-		101	1423	897	240	280	6	Ŭ		0
1	F	185	Total	С	Ν	Ο	S	0	1	0
	1	100	1412	891	238	277	6	0	1	0
1	G	185	Total	С	Ν	Ο	S	0	1	1
	<u> </u>	100	1397	883	237	271	6	0	1	1
1	Н	186	Total	С	Ν	Ο	S	0	0	0
	11	100	1415	894	240	275	6	0	0	0
1	Ι	183	Total	С	Ν	Ο	S	0	0	0
	1	100	1393	879	237	271	6	0	0	0
1	J	180	Total	С	Ν	Ο	S	0	0	0
	0	100	1372	866	232	268	6	0	0	0
1	Κ	183	Total	С	Ν	Ο	\mathbf{S}	0	1	0
	11	100	1393	878	237	272	6	0	Ĩ	0
1	L	190	Total	С	Ν	Ο	S	0	1	0
	Ц	150	1451	913	248	284	6		1	0
1	М	189	Total	С	Ν	Ο	S	0	0	0
	IVI	105	1456	914	250	286	6	0	0	U
1	Ν	181	Total	С	Ν	Ο	S	0	1	0
	11	101	1391	879	235	271	6		L	0

• Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit.

There are 84 discrepancies between the modelled and reference sequences:

A 196 HIS - expression tag UNP A7WZR9	Chain	Residue	esidue Modelled		Comment	Reference
	А	196	HIS	-	expression tag	UNP A7WZR9



Chain	Residue	Modelled	Actual	Comment	Reference
А	197	HIS	-	expression tag	UNP A7WZR9
А	198	HIS	-	expression tag	UNP A7WZR9
А	199	HIS	-	expression tag	UNP A7WZR9
А	200	HIS	_	expression tag	UNP A7WZR9
А	201	HIS	_	expression tag	UNP A7WZR9
В	196	HIS	-	expression tag	UNP A7WZR9
В	197	HIS	_	expression tag	UNP A7WZR9
В	198	HIS	_	expression tag	UNP A7WZR9
В	199	HIS	_	expression tag	UNP A7WZR9
В	200	HIS	-	expression tag	UNP A7WZR9
В	201	HIS	-	expression tag	UNP A7WZR9
С	196	HIS	_	expression tag	UNP A7WZR9
С	197	HIS	-	expression tag	UNP A7WZR9
С	198	HIS	-	expression tag	UNP A7WZR9
С	199	HIS	-	expression tag	UNP A7WZR9
С	200	HIS	_	expression tag	UNP A7WZR9
С	201	HIS	_	expression tag	UNP A7WZR9
D	196	HIS	-	expression tag	UNP A7WZR9
D	197	HIS	_	expression tag	UNP A7WZR9
D	198	HIS	_	expression tag	UNP A7WZR9
D	199	HIS	_	expression tag	UNP A7WZR9
D	200	HIS	-	expression tag	UNP A7WZR9
D	201	HIS	-	expression tag	UNP A7WZR9
Е	196	HIS	-	expression tag	UNP A7WZR9
Е	197	HIS	-	expression tag	UNP A7WZR9
Е	198	HIS	-	expression tag	UNP A7WZR9
Е	199	HIS	-	expression tag	UNP A7WZR9
Е	200	HIS	-	expression tag	UNP A7WZR9
Е	201	HIS	-	expression tag	UNP A7WZR9
F	196	HIS	-	expression tag	UNP A7WZR9
F	197	HIS	-	expression tag	UNP A7WZR9
F	198	HIS	-	expression tag	UNP A7WZR9
F	199	HIS	-	expression tag	UNP A7WZR9
F	200	HIS	-	expression tag	UNP A7WZR9
F	201	HIS	-	expression tag	UNP A7WZR9
G	196	HIS	-	expression tag	UNP A7WZR9
G	197	HIS	-	expression tag	UNP A7WZR9
G	198	HIS	-	expression tag	UNP A7WZR9
G	199	HIS	-	expression tag	UNP A7WZR9
G	200	HIS	-	expression tag	UNP A7WZR9
G	201	HIS	-	expression tag	UNP A7WZR9
Н	196	HIS	-	expression tag	UNP A7WZR9

Continued from previous page...




	ed from pre	vious page			
Chain	Residue	Modelled	Actual	Comment	Reference
Н	197	HIS	-	expression tag	UNP A7WZR9
Н	198	HIS	-	expression tag	UNP A7WZR9
Н	199	HIS	-	expression tag	UNP A7WZR9
Н	200	HIS	-	expression tag	UNP A7WZR9
Н	201	HIS	-	expression tag	UNP A7WZR9
Ι	196	HIS	_	expression tag	UNP A7WZR9
Ι	197	HIS	_	expression tag	UNP A7WZR9
Ι	198	HIS	-	expression tag	UNP A7WZR9
Ι	199	HIS	-	expression tag	UNP A7WZR9
Ι	200	HIS	-	expression tag	UNP A7WZR9
Ι	201	HIS	-	expression tag	UNP A7WZR9
J	196	HIS	-	expression tag	UNP A7WZR9
J	197	HIS	-	expression tag	UNP A7WZR9
J	198	HIS	-	expression tag	UNP A7WZR9
J	199	HIS	-	expression tag	UNP A7WZR9
J	200	HIS	-	expression tag	UNP A7WZR9
J	201	HIS	-	expression tag	UNP A7WZR9
K	196	HIS	-	expression tag	UNP A7WZR9
K	197	HIS	-	expression tag	UNP A7WZR9
K	198	HIS	-	expression tag	UNP A7WZR9
K	199	HIS	-	expression tag	UNP A7WZR9
K	200	HIS	-	expression tag	UNP A7WZR9
K	201	HIS	-	expression tag	UNP A7WZR9
L	196	HIS	-	expression tag	UNP A7WZR9
L	197	HIS	-	expression tag	UNP A7WZR9
L	198	HIS	-	expression tag	UNP A7WZR9
L	199	HIS	-	expression tag	UNP A7WZR9
L	200	HIS	-	expression tag	UNP A7WZR9
L	201	HIS	-	expression tag	UNP A7WZR9
М	196	HIS	-	expression tag	UNP A7WZR9
М	197	HIS	-	expression tag	UNP A7WZR9
М	198	HIS	-	expression tag	UNP A7WZR9
М	199	HIS	-	expression tag	UNP A7WZR9
М	200	HIS	_	expression tag	UNP A7WZR9
М	201	HIS	-	expression tag	UNP A7WZR9
N	196	HIS	-	expression tag	UNP A7WZR9
N	197	HIS	-	expression tag	UNP A7WZR9
N	198	HIS	-	expression tag	UNP A7WZR9
N	199	HIS	-	expression tag	UNP A7WZR9
N	200	HIS	-	expression tag	UNP A7WZR9
N	201	HIS	-	expression tag	UNP A7WZR9
L	1	1	1		1

Continued from previous page...

• Molecule 2 is (6 {S},9 {a} {S})-6-[(2 {S})-butan-2-yl]-8-[(4-methoxynaphthalen-1-yl)methyl



]-4,7-bis (oxidanylidene)- {N}-[4,4,4-tris(fluoranyl)butyl]-3,6,9,9 {a}-tetrahydro-2 {H}-pyra zino [1,2-a]pyrimidine-1-carboxamide (three-letter code: A1EEK) (formula: $\rm C_{28}H_{35}F_3N_4O_4)$ (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues		Ato	\mathbf{ms}			ZeroOcc	AltConf	
2	А	1	Total	С	F	Ν	0	0	0	
	A	1	39	28	3	4	4	0	U	
2	В	1	Total	С	F	Ν	Ο	0	0	
	D	1	39	28	3	4	4	0	0	
2	С	1	Total	С	F	Ν	0	0	0	
	U	1	39	28	3	4	4	0	0	
2	С	1	Total	С	F	Ν	Ο	0	0	
	C	1	39	28	3	4	4	0	0	
2	D	1	Total	С	F	Ν	Ο	0	0	
	D	1	39	28	3	4	4	0	0	
2	F	1	Total	С	F	Ν	Ο	0	0	
	Г	1	39	28	3	4	4	0	0	
2	G	1	Total	С	F	Ν	Ο	0	0	
2	G	1	39	28	3	4	4	0	0	
2	Н	1	Total	С	F	Ν	Ο	0	0	
2	11	1	39	28	3	4	4	0	0	
2	Ι	1	Total	С	F	Ν	Ο	0	0	
	1	I	39	28	3	4	4	0	0	
2	J	1	Total	С	F	Ν	Ο	0	0	
	0	I	39	28	3	4	4	0	0	
2	Κ	1	Total	\mathbf{C}	F	Ν	0	0	0	
	17	1	39	28	3	4	4	0	0	
2	L	1	Total	\mathbf{C}	F	Ν	Ο	0	0	
-		1	39	28	3	4	4		U U	



$\alpha \cdot \cdot \cdot$	C		
Continued	from	previous	page
	9	1	1 0

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
0	М	1	Total	С	F	Ν	Ο	0	0
Z	101	L	39	28	3	4	4	0	0
0	N	1	Total	С	F	Ν	0	0	0
Z	IN	L	39	28	3	4	4	0	0

• Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	F	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	G	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	Н	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 8 6 2 \end{array}$	0	0
3	Ι	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	J	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	K	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	L	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	М	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	Ν	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	101	Total O 101 101	0	0
4	В	97	Total O 97 97	0	0
4	С	98	Total O 98 98	0	0
4	D	98	Total O 98 98	0	0
4	Е	91	Total O 91 91	0	0
4	F	74	TotalO7474	0	0
4	G	86	Total O 86 86	0	0
4	Н	72	Total O 72 72	0	0
4	Ι	78	Total O 78 78	0	0
4	J	81	Total O 81 81	0	0
4	K	72	TotalO7272	0	0
4	L	79	Total O 79 79	0	0
4	М	85	Total O 85 85	0	0
4	Ν	90	Total O 90 90	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ATP-dependent Clp protease proteolytic subunit




• Molecule 1: ATP-dependent Clp protease proteolytic subunit



• Molecule 1: ATP-dependent Clp protease proteolytic subunit



• Molecule 1: ATP-dependent Clp protease proteolytic subunit





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	94.66Å 190.44Å 96.56Å	Depositor
a, b, c, α , β , γ	90.00° 117.40° 90.00°	Depositor
Resolution (Å)	95.22 - 1.98	Depositor
Resolution (A)	95.22 - 1.98	EDS
% Data completeness	96.4 (95.22 - 1.98)	Depositor
(in resolution range)	$96.4 \ (95.22 - 1.98)$	EDS
R _{merge}	0.10	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.16 (at 1.98 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.208 , 0.228	Depositor
n, n_{free}	0.212 , 0.230	DCC
R_{free} test set	10179 reflections (4.87%)	wwPDB-VP
Wilson B-factor $(Å^2)$	32.3	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.36, 30.9	EDS
L-test for twinning ²	$< L > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21643	wwPDB-VP
Average B, all atoms $(Å^2)$	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: A1EEK, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.33	0/1423	0.52	0/1925
1	В	0.37	0/1451	0.54	0/1966
1	С	0.33	0/1469	0.52	0/1990
1	D	0.32	0/1416	0.52	0/1915
1	Е	0.33	0/1441	0.52	0/1949
1	F	0.33	0/1433	0.52	0/1938
1	G	0.37	0/1418	0.51	0/1918
1	Н	0.33	0/1433	0.53	0/1937
1	Ι	0.33	0/1411	0.51	0/1907
1	J	0.36	0/1390	0.51	0/1880
1	Κ	0.36	0/1414	0.51	0/1913
1	L	0.36	0/1473	0.53	0/1992
1	М	0.34	0/1475	0.54	0/1993
1	Ν	0.39	0/1412	0.53	0/1908
All	All	0.35	0/20059	0.52	0/27131

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1402	0	1404	2	0
1	В	1433	0	1403	1	0
1	С	1447	0	1436	0	0
1	D	1398	0	1389	1	0
1	Е	1423	0	1415	1	0
1	F	1412	0	1406	1	0
1	G	1397	0	1393	1	0
1	Н	1415	0	1418	3	0
1	Ι	1393	0	1394	4	0
1	J	1372	0	1370	3	0
1	Κ	1393	0	1387	2	0
1	L	1451	0	1455	1	0
1	М	1456	0	1462	2	0
1	Ν	1391	0	1403	0	0
2	А	39	0	0	0	0
2	В	39	0	0	0	0
2	С	78	0	0	0	0
2	D	39	0	0	0	0
2	F	39	0	0	0	0
2	G	39	0	0	0	0
2	Н	39	0	0	0	0
2	Ι	39	0	0	0	0
2	J	39	0	0	0	0
2	Κ	39	0	0	0	0
2	L	39	0	0	0	0
2	М	39	0	0	0	0
2	Ν	39	0	0	0	0
3	А	8	0	14	0	0
3	В	8	0	14	0	0
3	С	8	0	14	0	0
3	D	8	0	14	0	0
3	Ε	8	0	14	0	0
3	F	8	0	14	0	0
3	G	8	0	14	0	0
3	Н	8	0	14	0	0
3	Ι	8	0	14	0	0
3	J	8	0	14	0	0
3	K	8	0	14	0	0
3	L	8	0	14	0	0
3	М	8	0	14	0	0
3	N	8	0	14	0	0
4	А	101	0	0	0	0
4	В	97	0	0	0	0



Mol	Chain	-	H(model)	H(added)	Clashes	Symm-Clashes
4	С	98	0	0	0	0
4	D	98	0	0	0	0
4	Е	91	0	0	0	0
4	F	74	0	0	0	0
4	G	86	0	0	0	0
4	Н	72	0	0	0	0
4	Ι	78	0	0	0	0
4	J	81	0	0	0	0
4	Κ	72	0	0	0	0
4	L	79	0	0	0	0
4	М	85	0	0	0	0
4	Ν	90	0	0	0	0
All	All	21643	0	19931	12	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

The worst 5 of 12 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:42:ASN:HD21	1:I:65:ASN:HD22	1.57	0.53
1:M:67:PRO:HA	1:M:97:ALA:HB3	1.95	0.48
1:D:106:ALA:HA	1:D:157:ARG:HD2	1.96	0.47
1:I:149:LYS:HE3	1:J:117:ASN:HD22	1.79	0.47
1:J:79:ASP:HB3	1:K:115:LEU:HD13	2.00	0.44

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



9K2D

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	181/201~(90%)	179~(99%)	2(1%)	0	100	100
1	В	186/201~(92%)	182 (98%)	4(2%)	0	100	100
1	\mathbf{C}	190/201~(94%)	187~(98%)	3~(2%)	0	100	100
1	D	180/201~(90%)	177~(98%)	3~(2%)	0	100	100
1	Ε	183/201~(91%)	179~(98%)	4 (2%)	0	100	100
1	F	182/201~(90%)	178 (98%)	4 (2%)	0	100	100
1	G	182/201~(90%)	180 (99%)	2(1%)	0	100	100
1	Н	182/201~(90%)	179 (98%)	3(2%)	0	100	100
1	Ι	179/201~(89%)	175 (98%)	4 (2%)	0	100	100
1	J	176/201~(88%)	172 (98%)	4 (2%)	0	100	100
1	Κ	180/201~(90%)	177 (98%)	3~(2%)	0	100	100
1	L	189/201~(94%)	186 (98%)	3(2%)	0	100	100
1	М	187/201 (93%)	183 (98%)	4 (2%)	0	100	100
1	Ν	178/201~(89%)	176 (99%)	2 (1%)	0	100	100
All	All	2555/2814 (91%)	2510 (98%)	45 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	149/169~(88%)	147 (99%)	2(1%)	65 62
1	В	149/169~(88%)	144 (97%)	5(3%)	32 21
1	С	151/169~(89%)	146 (97%)	5(3%)	33 22
1	D	147/169~(87%)	142 (97%)	5(3%)	32 21
1	Ε	150/169~(89%)	146 (97%)	4 (3%)	40 30
1	F	149/169~(88%)	146 (98%)	3 (2%)	50 43
1	G	146/169~(86%)	143 (98%)	3~(2%)	48 42



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	Η	149/169~(88%)	146~(98%)	3~(2%)	50	43
1	Ι	147/169~(87%)	144~(98%)	3~(2%)	50	43
1	J	145/169~(86%)	143~(99%)	2(1%)	62	60
1	Κ	147/169~(87%)	145~(99%)	2(1%)	62	60
1	L	154/169~(91%)	152~(99%)	2(1%)	65	62
1	М	155/169~(92%)	152 (98%)	3~(2%)	52	46
1	Ν	149/169~(88%)	146~(98%)	3~(2%)	50	43
All	All	2087/2366~(88%)	2042~(98%)	45 (2%)	47	40

Continued from previous page...

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Н	123	HIS
1	Κ	95	MET
1	Н	195	LYS
1	Ι	123	HIS
1	L	27	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	Ι	160	GLN
1	J	117	ASN
1	J	89	GLN
1	J	160	GLN
1	F	151	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

28 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	Bond lengths		В	ond ang	les
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	A1EEK	N	301	-	40,42,42	0.26	0	$53,\!61,\!61$	0.89	3 (5%)
3	MPD	J	302	-	7,7,7	0.25	0	9,10,10	0.38	0
3	MPD	G	302	-	$7,\!7,\!7$	0.25	0	9,10,10	0.35	0
2	A1EEK	М	301	-	40,42,42	0.26	0	$53,\!61,\!61$	0.84	3(5%)
3	MPD	D	302	-	7,7,7	0.23	0	9,10,10	0.41	0
3	MPD	Н	302	-	$7,\!7,\!7$	0.26	0	$9,\!10,\!10$	0.43	0
3	MPD	I	302	-	7,7,7	0.27	0	9,10,10	0.33	0
2	A1EEK	А	301	-	40,42,42	0.29	0	$53,\!61,\!61$	0.68	1 (1%)
2	A1EEK	J	301	-	40,42,42	0.26	0	$53,\!61,\!61$	0.75	1 (1%)
3	MPD	К	302	-	7,7,7	0.27	0	9,10,10	0.46	0
2	A1EEK	С	301	-	40,42,42	0.26	0	$53,\!61,\!61$	0.86	4 (7%)
2	A1EEK	F	301	-	40,42,42	0.24	0	$53,\!61,\!61$	0.76	2 (3%)
2	A1EEK	С	302	-	40,42,42	0.25	0	$53,\!61,\!61$	0.77	3 (5%)
3	MPD	N	302	-	7,7,7	0.25	0	9,10,10	0.43	0
2	A1EEK	K	301	-	40,42,42	0.27	0	$53,\!61,\!61$	0.74	2 (3%)
3	MPD	А	302	-	7,7,7	0.20	0	9,10,10	0.26	0
2	A1EEK	L	301	-	40,42,42	0.23	0	$53,\!61,\!61$	0.69	2 (3%)
2	A1EEK	G	301	-	40,42,42	0.26	0	53,61,61	0.82	2 (3%)
3	MPD	М	302	_	7,7,7	0.24	0	9,10,10	0.26	0
3	MPD	В	302	-	7,7,7	0.26	0	9,10,10	0.37	0
2	A1EEK	D	301	-	40,42,42	0.26	0	$53,\!61,\!61$	0.70	1 (1%)
2	A1EEK	Н	301	-	40,42,42	0.24	0	53,61,61	0.70	2 (3%)
2	A1EEK	Ι	301	-	40,42,42	0.23	0	53,61,61	0.72	2(3%)
3	MPD	F	302	_	7, 7, 7	0.27	0	9,10,10	0.33	0



Mol	Mol Type Chain Res Link		Link	Bond lengths			Bond angles			
	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	MPD	С	303	-	7,7,7	0.25	0	9,10,10	0.42	0
3	MPD	Е	301	-	7,7,7	0.27	0	$9,\!10,\!10$	0.43	0
2	A1EEK	В	301	-	40,42,42	0.28	0	$53,\!61,\!61$	0.64	1 (1%)
3	MPD	L	302	-	7,7,7	0.26	0	9,10,10	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1EEK	N	301	-	-	0/24/57/57	0/4/4/4
3	MPD	J	302	-	-	0/5/5/5	-
3	MPD	G	302	-	-	0/5/5/5	-
2	A1EEK	М	301	-	-	6/24/57/57	0/4/4/4
3	MPD	D	302	-	-	0/5/5/5	-
3	MPD	Н	302	-	-	0/5/5/5	-
3	MPD	Ι	302	-	-	0/5/5/5	-
2	A1EEK	А	301	-	-	1/24/57/57	0/4/4/4
2	A1EEK	J	301	-	-	5/24/57/57	0/4/4/4
3	MPD	К	302	-	-	0/5/5/5	-
2	A1EEK	С	301	-	-	6/24/57/57	0/4/4/4
2	A1EEK	F	301	-	-	5/24/57/57	0/4/4/4
2	A1EEK	С	302	-	-	1/24/57/57	0/4/4/4
3	MPD	Ν	302	-	-	0/5/5/5	-
2	A1EEK	К	301	-	-	6/24/57/57	0/4/4/4
3	MPD	А	302	-	-	0/5/5/5	-
2	A1EEK	L	301	-	-	3/24/57/57	0/4/4/4
2	A1EEK	G	301	-	-	2/24/57/57	0/4/4/4
3	MPD	М	302	-	-	1/5/5/5	-
3	MPD	В	302	-	-	0/5/5/5	-
2	A1EEK	D	301	-	-	1/24/57/57	0/4/4/4
2	A1EEK	Η	301	-	-	2/24/57/57	0/4/4/4
2	A1EEK	Ι	301	-	-	5/24/57/57	0/4/4/4
3	MPD	F	302	-	-	0/5/5/5	-
3	MPD	С	303	-	-	0/5/5/5	-
3	MPD	Е	301	-	-	0/5/5/5	-
2	A1EEK	В	301	-	-	1/24/57/57	0/4/4/4
3	MPD	L	302	-	-	0/5/5/5	-



There are no bond length outliers.

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Ν	301	A1EEK	C19-C18-N	-4.63	105.12	111.02
2	J	301	A1EEK	C19-C18-N	-3.69	106.32	111.02
2	Н	301	A1EEK	C19-C18-N	-3.47	106.60	111.02
2	С	302	A1EEK	C19-C18-N	-3.46	106.61	111.02
2	G	301	A1EEK	C19-C18-N	-3.44	106.64	111.02

There are no chirality outliers.

5 of 45 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	301	A1EEK	C-C1-C2-C3
2	G	301	A1EEK	C-C1-C2-C3
2	Н	301	A1EEK	C1-C2-C4-N3
2	Ι	301	A1EEK	C-C1-C2-C3
2	Κ	301	A1EEK	C-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.















































































5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2		$OWAB(Å^2)$	Q<0.9
1	А	184/201~(91%)	0.47	12 (6%) 26	37	19, 32, 51, 61	1 (0%)
1	В	190/201~(94%)	0.52	10 (5%) 33	44	25, 30, 48, 60	0
1	С	191/201~(95%)	0.43	5 (2%) 57	67	18, 31, 41, 51	1 (0%)
1	D	184/201~(91%)	0.34	9 (4%) 36	46	23, 29, 42, 55	0
1	Е	187/201~(93%)	0.42	11 (5%) 29	41	25, 31, 44, 61	0
1	F	185/201~(92%)	0.75	16 (8%) 18	26	22, 35, 47, 60	1 (0%)
1	G	185/201~(92%)	0.64	9 (4%) 36	46	22, 35, 48, 54	1 (0%)
1	Н	186/201~(92%)	0.47	9 (4%) 36	47	25, 31, 43, 56	0
1	Ι	183/201~(91%)	0.66	9 (4%) 36	46	28, 34, 48, 58	0
1	J	180/201~(89%)	0.71	11 (6%) 28	39	29, 35, 46, 54	0
1	Κ	183/201~(91%)	0.65	11 (6%) 29	39	19, 33, 47, 71	1 (0%)
1	L	190/201~(94%)	0.50	9 (4%) 37	48	19, 32, 46, 56	1 (0%)
1	М	189/201~(94%)	0.36	5 (2%) 57	67	24, 30, 39, 46	0
1	Ν	181/201~(90%)	0.38	9 (4%) 35	46	18, 29, 41, 53	1 (0%)
All	All	2598/2814~(92%)	0.52	135 (5%) 34	45	18, 32, 46, 71	7~(0%)

The worst 5 of 135 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Ν	8	ILE	4.7
1	F	3	LEU	4.4
1	В	11	THR	4.4
1	В	18	TYR	4.3
1	Κ	3	LEU	4.3



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	М	302	8/8	0.83	0.19	41,42,42,43	0
3	MPD	Ι	302	8/8	0.84	0.16	40,41,41,42	0
3	MPD	J	302	8/8	0.86	0.16	46,47,47,47	0
3	MPD	G	302	8/8	0.86	0.18	48,49,49,49	0
3	MPD	L	302	8/8	0.87	0.14	40,40,41,41	0
3	MPD	С	303	8/8	0.87	0.15	41,41,42,42	0
3	MPD	D	302	8/8	0.88	0.14	39,40,40,41	0
2	A1EEK	М	301	39/39	0.90	0.11	29,34,39,40	0
3	MPD	В	302	8/8	0.90	0.15	43,44,44,44	0
2	A1EEK	F	301	39/39	0.90	0.11	34,38,42,43	0
3	MPD	N	302	8/8	0.90	0.14	36,36,36,36	0
2	A1EEK	В	301	39/39	0.91	0.10	29,33,38,38	0
3	MPD	K	302	8/8	0.91	0.14	40,40,41,41	0
2	A1EEK	G	301	39/39	0.91	0.10	34,38,41,42	0
3	MPD	Н	302	8/8	0.91	0.15	43,43,43,43	0
2	A1EEK	Ι	301	39/39	0.91	0.10	32,38,43,44	0
2	A1EEK	С	301	39/39	0.92	0.09	29,34,38,39	0
3	MPD	Е	301	8/8	0.92	0.13	44,44,45,45	0
3	MPD	А	302	8/8	0.92	0.14	47,47,48,48	0
2	A1EEK	J	301	39/39	0.92	0.10	33,39,41,41	0
2	A1EEK	L	301	39/39	0.92	0.09	30,36,38,38	0
2	A1EEK	Н	301	39/39	0.93	0.09	26,29,34,35	0
2	A1EEK	N	301	39/39	0.93	0.09	27,33,38,39	0
2	A1EEK	А	301	39/39	0.93	0.09	29,33,35,36	0
2	A1EEK	К	301	39/39	0.94	0.09	35,39,43,44	0
2	A1EEK	D	301	39/39	0.94	0.07	24,26,29,29	0
2	A1EEK	С	302	39/39	0.94	0.08	27,32,37,38	0


Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
3	MPD	F	302	8/8	0.94	0.11	42,43,43,43	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






























































































































6.5 Other polymers (i)

There are no such residues in this entry.

