



# wwPDB X-ray Structure Validation Summary Report i

Nov 16, 2023 – 01:49 AM JST

PDB ID : 6KEZ  
Title : Crystal structure of GAPDH/CP12/PRK complex from *Arabidopsis thaliana*  
Authors : Yu, A.; Xie, Y.; Li, M.  
Deposited on : 2019-07-05  
Resolution : 3.50 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain				
1	G	339		80%	19%	1%	.
1	H	339		76%	22%	1%	..
2	I	352		75%	21%	3%	..
2	J	352		76%	20%	3%	..
2	K	352		76%	20%	2%	..
2	L	352		72%	24%	4%	..
3	M	80		68%	22%	11%	• 9%
3	N	80		62%	24%	10%	5% 9%
3	O	80		75%	14%	2%	• 10%
3	P	80		78%	11%	2%	• 10%

## 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 33826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase GAPA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total 2552	C 1612	N 444	O 487	S 9	0	0	0
1	B	336	Total 2552	C 1612	N 444	O 487	S 9	0	0	0
1	C	336	Total 2552	C 1612	N 444	O 487	S 9	0	0	0
1	D	335	Total 2542	C 1606	N 442	O 485	S 9	0	0	0
1	E	336	Total 2551	C 1612	N 444	O 486	S 9	0	0	0
1	F	336	Total 2552	C 1612	N 444	O 487	S 9	0	0	0
1	G	336	Total 2552	C 1612	N 444	O 487	S 9	0	0	0
1	H	336	Total 2552	C 1612	N 444	O 487	S 9	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP P25856
A	-1	ASN	-	expression tag	UNP P25856
A	0	ALA	-	expression tag	UNP P25856
B	-2	SER	-	expression tag	UNP P25856
B	-1	ASN	-	expression tag	UNP P25856
B	0	ALA	-	expression tag	UNP P25856
C	-2	SER	-	expression tag	UNP P25856
C	-1	ASN	-	expression tag	UNP P25856
C	0	ALA	-	expression tag	UNP P25856
D	-2	SER	-	expression tag	UNP P25856
D	-1	ASN	-	expression tag	UNP P25856
D	0	ALA	-	expression tag	UNP P25856
E	-2	SER	-	expression tag	UNP P25856

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	ASN	-	expression tag	UNP P25856
E	0	ALA	-	expression tag	UNP P25856
F	-2	SER	-	expression tag	UNP P25856
F	-1	ASN	-	expression tag	UNP P25856
F	0	ALA	-	expression tag	UNP P25856
G	-2	SER	-	expression tag	UNP P25856
G	-1	ASN	-	expression tag	UNP P25856
G	0	ALA	-	expression tag	UNP P25856
H	-2	SER	-	expression tag	UNP P25856
H	-1	ASN	-	expression tag	UNP P25856
H	0	ALA	-	expression tag	UNP P25856

- Molecule 2 is a protein called Phosphoribulokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	342	Total	C	N	O	S			
			2713	1730	452	520	11	0	0	0
2	K	342	Total	C	N	O	S			
			2713	1730	452	520	11	0	0	0
2	J	342	Total	C	N	O	S			
			2713	1730	452	520	11	0	0	0
2	L	342	Total	C	N	O	S			
			2710	1729	452	518	11	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	0	SER	-	expression tag	UNP P25697
I	1	ASN	-	expression tag	UNP P25697
I	2	ALA	-	expression tag	UNP P25697
K	0	SER	-	expression tag	UNP P25697
K	1	ASN	-	expression tag	UNP P25697
K	2	ALA	-	expression tag	UNP P25697
J	0	SER	-	expression tag	UNP P25697
J	1	ASN	-	expression tag	UNP P25697
J	2	ALA	-	expression tag	UNP P25697
L	0	SER	-	expression tag	UNP P25697
L	1	ASN	-	expression tag	UNP P25697
L	2	ALA	-	expression tag	UNP P25697


- Molecule 3 is a protein called Calvin cycle protein CP12-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	73	Total	C	N	O	S	0	0	0
			550	328	92	126	4			
3	N	73	Total	C	N	O	S	0	0	0
			550	328	92	126	4			
3	O	72	Total	C	N	O	S	0	0	0
			540	324	91	121	4			
3	P	72	Total	C	N	O	S	0	0	0
			543	325	91	123	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	-1	SER	-	expression tag	UNP Q9LZP9
M	0	ASN	-	expression tag	UNP Q9LZP9
M	1	ALA	-	expression tag	UNP Q9LZP9
N	-1	SER	-	expression tag	UNP Q9LZP9
N	0	ASN	-	expression tag	UNP Q9LZP9
N	1	ALA	-	expression tag	UNP Q9LZP9
O	-1	SER	-	expression tag	UNP Q9LZP9
O	0	ASN	-	expression tag	UNP Q9LZP9
O	1	ALA	-	expression tag	UNP Q9LZP9
P	-1	SER	-	expression tag	UNP Q9LZP9
P	0	ASN	-	expression tag	UNP Q9LZP9
P	1	ALA	-	expression tag	UNP Q9LZP9

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total C N O P					0	0
			44	21	7	14	2		
4	B	1	Total C N O P					0	0
			44	21	7	14	2		
4	C	1	Total C N O P					0	0
			44	21	7	14	2		
4	D	1	Total C N O P					0	0
			44	21	7	14	2		
4	E	1	Total C N O P					0	0
			44	21	7	14	2		
4	F	1	Total C N O P					0	0
			44	21	7	14	2		
4	G	1	Total C N O P					0	0
			44	21	7	14	2		
4	H	1	Total C N O P					0	0
			44	21	7	14	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total O		0	0
			2	2		
5	B	9	Total O		0	0
			9	9		
5	C	7	Total O		0	0
			7	7		
5	D	1	Total O		0	0
			1	1		
5	E	4	Total O		0	0
			4	4		
5	F	2	Total O		0	0
			2	2		
5	G	6	Total O		0	0
			6	6		
5	H	2	Total O		0	0
			2	2		
5	I	1	Total O		0	0
			1	1		
5	K	1	Total O		0	0
			1	1		
5	N	2	Total O		0	0
			2	2		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase GAPA1




- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase GAPA1



- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase GAPA1






- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase GAPA1

Chain D:



- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase GAPA1

Chain E:



- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase GAPA1


Chain F:



- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase GAPA1


Chain G:






- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase GAPA1

Chain H: 76% 22% ..



- Molecule 2: Phosphoribulokinase

Chain I: 75% 21% ..



L273 E274 R281 L295 S296 Y300 V303 Q306 D312 Q324 V327 T331 R332 R346 ALA GLU ALA LYS ALA

- Molecule 2: Phosphoribulokinase

Chain K: 76% 20% ..



L108 R309 H310 F313 S316 R318 G319 T320 I331 I334 A343 T344 A345 B346 ALA GLU ALA LYS ALA

- Molecule 2: Phosphoribulokinase

Chain J: 76% 20% ..




- Molecule 2: Phosphoribulokinase




- Molecule 3: Calvin cycle protein CP12-2



- Molecule 3: Calvin cycle protein CP12-2



- Molecule 3: Calvin cycle protein CP12-2



- Molecule 3: Calvin cycle protein CP12-2





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.81Å 204.73Å 157.04Å 90.00° 106.88° 90.00°	Depositor
Resolution (Å)	50.00 – 3.50 50.09 – 3.50	Depositor EDS
% Data completeness (in resolution range)	87.3 (50.00-3.50) 86.9 (50.09-3.50)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	3.02 (at 3.48Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
$R$ , $R_{free}$	0.246 , 0.287 0.247 , 0.284	Depositor DCC
$R_{free}$ test set	2956 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.0	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , -10.0	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.45$ , $< L^2 > = 0.27$	Xtriage
Estimated twinning fraction	0.118 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.83	EDS
Total number of atoms	33826	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/2596	0.45	0/3523
1	B	0.25	0/2596	0.45	0/3523
1	C	0.25	0/2596	0.44	0/3523
1	D	0.24	0/2586	0.44	0/3512
1	E	0.38	0/2595	0.48	0/3523
1	F	0.25	0/2596	0.45	0/3523
1	G	0.25	0/2596	0.45	0/3523
1	H	0.26	0/2596	0.45	0/3523
2	I	0.27	0/2772	0.50	0/3754
2	J	0.26	0/2772	0.45	0/3754
2	K	0.25	0/2772	0.44	0/3754
2	L	0.25	0/2769	0.45	0/3750
3	M	0.26	0/557	0.43	0/753
3	N	0.24	0/557	0.42	0/753
3	O	0.24	0/547	0.38	0/740
3	P	0.26	0/550	0.43	0/744
All	All	0.27	0/34053	0.45	0/46175

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2552	0	2593	54	0
1	B	2552	0	2593	61	0
1	C	2552	0	2593	50	0
1	D	2542	0	2580	42	0
1	E	2551	0	2593	57	0
1	F	2552	0	2593	61	0
1	G	2552	0	2593	46	0
1	H	2552	0	2593	57	0
2	I	2713	0	2687	50	0
2	J	2713	0	2687	42	0
2	K	2713	0	2689	44	0
2	L	2710	0	2685	58	0
3	M	550	0	496	14	0
3	N	550	0	496	19	0
3	O	540	0	489	16	0
3	P	543	0	491	7	0
4	A	44	0	26	1	0
4	B	44	0	26	2	0
4	C	44	0	26	5	0
4	D	44	0	26	1	0
4	E	44	0	26	1	0
4	F	44	0	26	1	0
4	G	44	0	26	2	0
4	H	44	0	26	2	0
5	A	2	0	0	1	0
5	B	9	0	0	1	0
5	C	7	0	0	2	0
5	D	1	0	0	0	0
5	E	4	0	0	1	0
5	F	2	0	0	1	0
5	G	6	0	0	0	0
5	H	2	0	0	0	0
5	I	1	0	0	0	0
5	K	1	0	0	0	0
5	N	2	0	0	0	0
All	All	33826	0	33659	588	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 588 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:240:VAL:HG11	1:B:283:SER:HB3	1.54	0.87
1:F:10:GLY:HA2	5:F:501:HOH:O	1.73	0.86
1:G:187:ARG:HH22	3:N:74:ARG:HH12	1.24	0.82
1:C:183:THR:HG22	4:C:401:NAD:H72N	1.43	0.80
3:O:12:GLU:CG	3:O:16:LYS:HE3	2.14	0.78

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	334/339 (98%)	319 (96%)	15 (4%)	0	100 100
1	B	334/339 (98%)	320 (96%)	14 (4%)	0	100 100
1	C	334/339 (98%)	326 (98%)	8 (2%)	0	100 100
1	D	333/339 (98%)	324 (97%)	9 (3%)	0	100 100
1	E	334/339 (98%)	321 (96%)	13 (4%)	0	100 100
1	F	334/339 (98%)	325 (97%)	9 (3%)	0	100 100
1	G	334/339 (98%)	324 (97%)	10 (3%)	0	100 100
1	H	334/339 (98%)	324 (97%)	10 (3%)	0	100 100
2	I	340/352 (97%)	324 (95%)	16 (5%)	0	100 100
2	J	340/352 (97%)	326 (96%)	14 (4%)	0	100 100
2	K	340/352 (97%)	333 (98%)	7 (2%)	0	100 100
2	L	340/352 (97%)	332 (98%)	8 (2%)	0	100 100
3	M	71/80 (89%)	69 (97%)	2 (3%)	0	100 100
3	N	71/80 (89%)	70 (99%)	1 (1%)	0	100 100
3	O	70/80 (88%)	69 (99%)	1 (1%)	0	100 100
3	P	70/80 (88%)	70 (100%)	0	0	100 100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	4313/4440 (97%)	4176 (97%)	137 (3%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	279/281 (99%)	266 (95%)	13 (5%)	26 60
1	B	279/281 (99%)	268 (96%)	11 (4%)	32 64
1	C	279/281 (99%)	265 (95%)	14 (5%)	24 58
1	D	278/281 (99%)	269 (97%)	9 (3%)	39 69
1	E	279/281 (99%)	268 (96%)	11 (4%)	32 64
1	F	279/281 (99%)	268 (96%)	11 (4%)	32 64
1	G	279/281 (99%)	272 (98%)	7 (2%)	47 75
1	H	279/281 (99%)	273 (98%)	6 (2%)	52 78
2	I	300/305 (98%)	287 (96%)	13 (4%)	29 62
2	J	300/305 (98%)	289 (96%)	11 (4%)	34 65
2	K	300/305 (98%)	292 (97%)	8 (3%)	44 73
2	L	299/305 (98%)	285 (95%)	14 (5%)	26 60
3	M	60/64 (94%)	58 (97%)	2 (3%)	38 68
3	N	60/64 (94%)	53 (88%)	7 (12%)	5 26
3	O	58/64 (91%)	57 (98%)	1 (2%)	60 82
3	P	59/64 (92%)	54 (92%)	5 (8%)	10 39
All	All	3667/3724 (98%)	3524 (96%)	143 (4%)	32 64

5 of 143 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	116	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	L	219	LEU
3	N	21	THR
1	E	99	THR
1	E	74	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
3	M	78	ASN
3	N	78	ASN
1	D	41	GLN
1	D	111	HIS
1	F	316	ASN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAD	B	401	-	42,48,48	0.80	1 (2%)	50,73,73	1.19	5 (10%)
4	NAD	E	401	-	42,48,48	0.81	1 (2%)	50,73,73	1.25	5 (10%)
4	NAD	F	401	-	42,48,48	0.81	1 (2%)	50,73,73	1.23	5 (10%)
4	NAD	A	401	-	42,48,48	0.82	1 (2%)	50,73,73	1.26	5 (10%)
4	NAD	C	401	-	42,48,48	2.59	24 (57%)	50,73,73	2.57	18 (36%)
4	NAD	H	401	-	42,48,48	0.81	1 (2%)	50,73,73	1.08	4 (8%)
4	NAD	G	401	-	42,48,48	0.81	1 (2%)	50,73,73	1.20	4 (8%)
4	NAD	D	401	-	42,48,48	0.81	1 (2%)	50,73,73	1.25	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAD	B	401	-	-	6/26/62/62	0/5/5/5
4	NAD	E	401	-	-	8/26/62/62	0/5/5/5
4	NAD	F	401	-	-	10/26/62/62	0/5/5/5
4	NAD	A	401	-	-	11/26/62/62	0/5/5/5
4	NAD	C	401	-	-	11/26/62/62	0/5/5/5
4	NAD	H	401	-	-	3/26/62/62	0/5/5/5
4	NAD	G	401	-	-	11/26/62/62	0/5/5/5
4	NAD	D	401	-	-	11/26/62/62	0/5/5/5

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	401	NAD	C2D-C1D	-6.90	1.43	1.53
4	C	401	NAD	O7N-C7N	-4.39	1.15	1.24
4	C	401	NAD	O4D-C4D	-4.14	1.35	1.45
4	C	401	NAD	PN-O1N	-3.99	1.36	1.50
4	C	401	NAD	C4A-N3A	-3.96	1.30	1.35

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	401	NAD	O4B-C1B-C2B	-7.86	95.45	106.93
4	C	401	NAD	O3D-C3D-C4D	-7.24	90.13	111.05

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	C	401	NAD	O4D-C1D-C2D	-6.98	96.73	106.93
4	C	401	NAD	O7N-C7N-N7N	-4.65	115.97	122.58
4	C	401	NAD	N3A-C2A-N1A	-4.10	122.27	128.68

There are no chirality outliers.

5 of 71 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	401	NAD	C5B-O5B-PA-O3
4	A	401	NAD	C5D-O5D-PN-O1N
4	A	401	NAD	O4D-C1D-N1N-C2N
4	A	401	NAD	O4D-C1D-N1N-C6N
4	A	401	NAD	C2D-C1D-N1N-C2N


There are no ring outliers.


8 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symmm-Clashes
4	B	401	NAD	2	0
4	E	401	NAD	1	0
4	F	401	NAD	1	0
4	A	401	NAD	1	0
4	C	401	NAD	5	0
4	H	401	NAD	2	0
4	G	401	NAD	2	0
4	D	401	NAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	336/339 (99%)	-0.22	1 (0%)	94 91	21, 35, 52, 79	0
1	B	336/339 (99%)	-0.17	0	100 100	20, 32, 48, 72	0
1	C	336/339 (99%)	-0.16	0	100 100	22, 36, 55, 66	0
1	D	335/339 (98%)	-0.11	0	100 100	19, 39, 60, 83	0
1	E	336/339 (99%)	-0.04	1 (0%)	94 91	22, 50, 78, 92	0
1	F	336/339 (99%)	-0.17	0	100 100	19, 37, 52, 69	0
1	G	336/339 (99%)	-0.17	0	100 100	17, 31, 46, 64	0
1	H	336/339 (99%)	-0.15	0	100 100	23, 40, 59, 79	0
2	I	342/352 (97%)	0.02	3 (0%)	84 79	37, 64, 91, 155	0
2	J	342/352 (97%)	0.27	11 (3%)	47 42	35, 82, 124, 142	0
2	K	342/352 (97%)	0.09	7 (2%)	65 60	33, 68, 99, 112	0
2	L	342/352 (97%)	0.26	13 (3%)	40 36	38, 87, 127, 168	3 (0%)
3	M	73/80 (91%)	0.66	9 (12%)	4 5	42, 148, 174, 188	0
3	N	73/80 (91%)	0.69	8 (10%)	5 6	30, 138, 169, 174	0
3	O	72/80 (90%)	0.22	2 (2%)	53 47	26, 91, 121, 176	0
3	P	72/80 (90%)	0.21	2 (2%)	53 47	42, 101, 135, 162	2 (2%)
All	All	4345/4440 (97%)	-0.01	57 (1%)	77 71	17, 45, 114, 188	5 (0%)

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	63	LEU	5.7
3	N	37	GLU	5.6
2	I	19	LYS	4.9
2	J	73	VAL	4.2
3	N	29	GLY	4.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAD	D	401	44/44	0.89	0.33	45,52,61,65	0
4	NAD	E	401	44/44	0.91	0.30	58,66,72,76	0
4	NAD	G	401	44/44	0.92	0.26	31,31,39,43	0
4	NAD	B	401	44/44	0.94	0.20	32,32,38,45	0
4	NAD	A	401	44/44	0.94	0.26	34,34,41,44	0
4	NAD	H	401	44/44	0.94	0.27	37,40,50,53	0
4	NAD	C	401	44/44	0.95	0.21	30,30,43,47	0
4	NAD	F	401	44/44	0.95	0.21	32,32,32,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.