

Full wwPDB X-ray Structure Validation Report (i)

May 6, 2025 – 01:02 PM JST

PDB ID : 9KRQ / pdb_00009krq

Title : Crystal structure of ZXC21 RBD

Authors : Lan, J.; Wang, C.H.

Deposited on : 2024-11-28

Resolution : 3.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 2.0rc1

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.006 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.43.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 3.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	164625	1842 (3.10-3.06)
Clashscore	180529	1965 (3.10-3.06)
Ramachandran outliers	177936	1859 (3.10-3.06)
Sidechain outliers	177891	1858 (3.10-3.06)
RSRZ outliers	164620	1842 (3.10-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	181	5% 65%	30%	6%
1	В	181	65%	31%	



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2942 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Spike glycoprotein.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	D	181	Total	С	N	О	S	0	0	0
1		101	1464	938	249	271	6	U	U	U
1	Λ	101	Total	С	N	О	S	0	0	0
1	I A	181	1464	938	249	271	6	0	U	U

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	439	THR	VAL	conflict	UNP A0A8A0R1U2
В	441	HIS	SER	conflict	UNP A0A8A0R1U2
В	504	HIS	-	expression tag	UNP A0A8A0R1U2
В	505	HIS	-	expression tag	UNP A0A8A0R1U2
В	506	HIS	-	expression tag	UNP A0A8A0R1U2
В	507	HIS	-	expression tag	UNP A0A8A0R1U2
В	508	HIS	-	expression tag	UNP A0A8A0R1U2
В	509	HIS	-	expression tag	UNP A0A8A0R1U2
A	439	THR	VAL	conflict	UNP A0A8A0R1U2
A	441	HIS	SER	conflict	UNP A0A8A0R1U2
A	504	HIS	-	expression tag	UNP A0A8A0R1U2
A	505	HIS	-	expression tag	UNP A0A8A0R1U2
A	506	HIS	-	expression tag	UNP A0A8A0R1U2
A	507	HIS	-	expression tag	UNP A0A8A0R1U2
A	508	HIS	-	expression tag	UNP A0A8A0R1U2
A	509	HIS	-	expression tag	UNP A0A8A0R1U2

• Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	В	1	Total	C 8	N 1	O 5	0	0
			14	8	1	5		



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Spike glycoprotein





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	44.16Å 73.77Å 117.83Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.87 - 3.08	Depositor
Resolution (A)	31.87 - 3.08	EDS
% Data completeness	95.3 (31.87-3.08)	Depositor
(in resolution range)	95.1 (31.87-3.08)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.95 (at 3.06Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
D D.	0.236 , 0.269	Depositor
R, R_{free}	0.244 , 0.273	DCC
R_{free} test set	6830 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	53.8	Xtriage
Anisotropy	0.610	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 52.9	EDS
L-test for twinning ²	$ < L > = 0.47, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	2942	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.68	0/1510	1.06	5/2059~(0.2%)	
1	В	0.56	0/1510	0.92	5/2059 (0.2%)	
All	All	0.62	0/3020	0.99	10/4118 (0.2%)	

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	368	THR	N-CA-C	-7.31	103.53	114.64
1	В	369	SER	CA-C-N	-6.89	113.88	122.84
1	В	369	SER	C-N-CA	-6.89	113.88	122.84
1	A	480	LEU	N-CA-C	-6.31	102.65	110.41
1	A	472	TYR	CA-CB-CG	5.88	124.48	113.90
1	В	472	TYR	CA-CB-CG	5.48	123.77	113.90
1	В	368	THR	CB-CA-C	5.31	117.31	109.29
1	A	508	HIS	N-CA-C	-5.22	104.50	111.02
1	A	482	TYR	N-CA-C	-5.21	105.79	113.61
1	A	481	GLU	N-CA-C	-5.03	105.39	113.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1464	0	1382	44	0
1	В	1464	0	1381	42	0
2	В	14	0	13	1	0
All	All	2942	0	2776	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:494:LEU:HD22	1:A:495:LEU:HG	1.56	0.88
1:A:389:THR:HG21	1:A:497:ALA:HB3	1.56	0.87
1:B:407:ALA:HB3	1:B:410:GLN:HG3	1.65	0.78
1:A:406:VAL:O	1:A:421:LEU:HD12	1.87	0.75
1:B:336:LYS:HD2	1:B:336:LYS:H	1.50	0.74
1:A:502:CYS:HB3	1:A:507:HIS:HA	1.70	0.72
1:A:376:TYR:CE2	1:A:408:PRO:HD2	2.33	0.64
1:A:351:ARG:HG3	1:A:394:ASP:OD1	1.98	0.63
1:B:500:THR:HG23	1:B:501:VAL:HG23	1.80	0.62
1:A:494:LEU:CD2	1:A:495:LEU:HG	2.27	0.62
1:B:332:CYS:HB2	1:B:359:ALA:HA	1.81	0.61
1:B:457:ARG:HH11	1:B:459:LEU:HD11	1.66	0.61
1:A:357:CYS:SG	1:A:358:ILE:N	2.70	0.61
1:A:503:GLY:O	1:A:507:HIS:HB3	2.00	0.61
1:B:366:ASN:OD1	1:B:368:THR:HG22	2.01	0.60
1:B:421:LEU:HD21	1:B:489:VAL:HG11	1.83	0.60
1:A:336:LYS:HD2	1:A:336:LYS:H	1.67	0.57
1:A:422:PRO:HG3	1:A:454:PRO:HB3	1.86	0.57
1:A:446:SER:HB2	1:A:447:HIS:ND1	2.19	0.57
1:B:351:ARG:HG3	1:B:394:ASP:OD1	2.04	0.57
1:B:437:GLN:HG3	1:B:438:ASP:H	1.69	0.57
1:B:343:PHE:CD1	1:B:486:ARG:HD3	2.41	0.56
1:A:415:ALA:HA	1:A:419:TYR:O	2.06	0.56
1:B:373:PHE:O	1:B:374:LYS:HD3	2.04	0.56
1:B:333:PRO:HA	1:B:336:LYS:NZ	2.21	0.56
1:A:333:PRO:HA	1:A:336:LYS:HD3	1.87	0.55
1:B:343:PHE:CE2	1:B:395:THR:HB	2.41	0.55
1:A:358:ILE:HD12	1:A:507:HIS:HB2	1.89	0.55
1:B:367:SER:HA	1:A:342:ARG:NH1	2.23	0.54
1:A:412:GLY:O	1:A:416:ASP:HB2	2.05	0.54
1:A:405:GLN:OE1	1:A:412:GLY:HA3	2.08	0.53

Continued on next page...



 $Continued\ from\ previous\ page...$

Continued from previ		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	overlap(Å)
1:A:331:VAL:HG22	1:A:332:CYS:H	1.74	0.53
1:B:437:GLN:HG3	1:B:438:ASP:N	2.25	0.52
1:A:386:LEU:HD13	1:A:494:LEU:HD11	1.92	0.52
1:B:480:LEU:HA	1:B:483:GLN:HG3	1.92	0.52
1:B:329:THR:HB	1:B:356:ASP:O	2.10	0.51
1:A:336:LYS:HD2	1:A:336:LYS:N	2.26	0.50
1:B:336:LYS:HD2	1:B:336:LYS:N	2.24	0.50
1:A:480:LEU:O	1:A:481:GLU:HB2	2.10	0.49
1:A:427:GLY:HA2	1:A:492:PHE:CD2	2.46	0.49
1:B:348:ALA:HA	1:B:457:ARG:HD3	1.95	0.49
1:A:445:ARG:NH2	1:A:458:ASP:O	2.46	0.48
1:B:382:LYS:O	1:B:386:LEU:HD13	2.12	0.48
1:A:332:CYS:O	1:A:334:PHE:N	2.46	0.48
1:B:422:PRO:HG3	1:B:454:PRO:HB2	1.96	0.48
2:B:601:NAG:H5	1:A:441:HIS:CE1	2.49	0.48
1:B:434:THR:HB	1:B:437:GLN:CG	2.44	0.47
1:A:376:TYR:O	1:A:426:THR:HA	2.14	0.47
1:B:333:PRO:HA	1:B:336:LYS:HZ3	1.80	0.47
1:B:405:GLN:OE1	1:B:412:GLY:HA3	2.13	0.47
1:A:416:ASP:CG	1:A:451:LYS:HD3	2.40	0.47
1:B:346:VAL:HA	1:B:396:PHE:HB2	1.96	0.46
1:B:395:THR:HG22	1:B:488:VAL:HG22	1.98	0.46
1:A:361:TYR:CE2	1:A:383:LEU:HB3	2.51	0.46
1:B:373:PHE:C	1:B:373:PHE:CD1	2.91	0.46
1:A:357:CYS:O	1:A:501:VAL:HA	2.16	0.46
1:B:343:PHE:CE1	1:B:486:ARG:HD3	2.51	0.45
1:B:362:THR:O	1:B:363:VAL:C	2.60	0.45
1:A:391:VAL:HG21	1:A:501:VAL:HG21	1.98	0.45
1:A:406:VAL:O	1:A:421:LEU:CD1	2.60	0.45
1:B:355:SER:HA	1:B:501:VAL:HG22	1.99	0.44
1:A:480:LEU:C	1:A:482:TYR:H	2.25	0.44
1:A:334:PHE:O	1:A:338:PHE:HB2	2.18	0.44
1:A:431:ALA:HA	1:A:486:ARG:O	2.18	0.43
1:B:346:VAL:HG22	1:B:418:ASN:HB3	2.01	0.42
1:B:414:ILE:HA	1:B:418:ASN:HB2	2.01	0.42
1:A:387:CYS:HA	1:A:502:CYS:HA	2.00	0.42
1:B:357:CYS:SG	1:B:501:VAL:HG13	2.59	0.41
1:B:373:PHE:C	1:B:373:PHE:HD1	2.28	0.41
1:B:387:CYS:SG	1:B:506:HIS:CG	3.13	0.41
1:A:490:LEU:HD23	1:A:490:LEU:HA	1.80	0.41
1:B:467:ARG:HH11	1:B:469:LEU:HB2	1.86	0.41

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap $(ext{Å})$
1:B:495:LEU:HD22	1:B:495:LEU:HA	1.76	0.41
1:A:452:LEU:HD23	1:A:452:LEU:HA	1.73	0.41
1:B:400:PHE:O	1:B:403:VAL:HG23	2.19	0.41
1:B:427:GLY:HA2	1:B:492:PHE:CZ	2.56	0.41
1:A:420:LYS:HB2	1:A:420:LYS:HE3	1.78	0.41
1:B:355:SER:HA	1:B:501:VAL:CG2	2.50	0.41
1:B:376:TYR:CD1	1:B:376:TYR:N	2.88	0.41
1:A:390:SER:O	1:A:492:PHE:HA	2.21	0.41
1:A:507:HIS:O	1:A:507:HIS:CG	2.73	0.41
1:A:334:PHE:HD1	1:A:334:PHE:HA	1.72	0.40
1:A:399:ARG:HD3	1:A:482:TYR:HA	2.03	0.40
1:A:462:ASP:OD1	1:A:462:ASP:N	2.51	0.40
1:B:389:THR:HG21	1:B:496:ASN:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	179/181 (99%)	165 (92%)	14 (8%)	0	100	100
1	В	179/181 (99%)	155 (87%)	23 (13%)	1 (1%)	22	51
All	All	358/362 (99%)	320 (89%)	37 (10%)	1 (0%)	37	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type	
1	В	364	PHE	



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	163/163 (100%)	149 (91%)	14 (9%)		8	30
1	В	163/163 (100%)	154 (94%)	9 (6%)		18	45
All	All	326/326 (100%)	303 (93%)	23 (7%)		12	37

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	332	CYS
1	В	363	VAL
1	В	364	PHE
1	В	368	THR
1	В	436	LYS
1	В	493	GLU
1	В	495	LEU
1	В	496	ASN
1	В	504	HIS
1	A	332	CYS
1	A	357	CYS
1	A	362	THR
1	A	405	GLN
1	A	424	ASP
1	A	450	THR
1	A	452	LEU
1	A	477	ASN
1	A	493	GLU
1	A	494	LEU
1	A	496	ASN
1	A	500	THR
1	A	501	VAL
1	A	507	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:



Mol	Chain	Res	Type
1	В	505	HIS
1	A	339	ASN
1	A	366	ASN
1	A	441	HIS
1	A	496	ASN
1	A	508	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Type Chain		Link	Bond lengths			Bond angles		
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	В	601	1	14,14,15	0.37	0	17,19,21	1.04	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	В	601	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	601	NAG	C2-N2-C7	3.69	128.16	122.90

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	601	NAG	C3-C2-N2-C7
2	В	601	NAG	O7-C7-N2-C2
2	В	601	NAG	C8-C7-N2-C2
2	В	601	NAG	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	601	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	181/181 (100%)	0.08	9 (4%) 35	20	10, 37, 71, 137	0
1	В	181/181 (100%)	0.16	8 (4%) 39	23	10, 46, 93, 135	0
All	All	362/362 (100%)	0.12	17 (4%) 37	21	10, 41, 85, 137	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	496	ASN	4.4
1	A	495	LEU	3.8
1	A	496	ASN	3.6
1	В	497	ALA	3.3
1	В	494	LEU	3.1
1	В	495	LEU	2.8
1	В	506	HIS	2.7
1	A	503	GLY	2.6
1	A	464	ASN	2.4
1	A	498	PRO	2.4
1	A	502	CYS	2.4
1	В	508	HIS	2.3
1	В	498	PRO	2.2
1	A	497	ALA	2.2
1	A	449	SER	2.2
1	A	475	ASN	2.2
1	В	437	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)


There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	NAG	В	601	14/15	0.62	0.23	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers (i)

There are no such residues in this entry.

