



wwPDB EM Validation Summary Report i

Jun 1, 2025 – 12:09 AM JST

PDB ID : 9L3Y / pdb_00009l3y
EMDB ID : EMD-62798
Title : Cryo-EM structure of the G-protein coupled receptor 1 (GPR1) in complex with chemerin and Gi1
Authors : Zhu, Y.; He, M.; Wu, B.; Zhao, Q.
Deposited on : 2024-12-19
Resolution : 3.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

EMDB validation analysis : **FAILED**
MolProbitY : 4-5-2 with Phenix2.0rc1
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.



2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7383 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chemerin-like receptor 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	R	298	2391	1620	380	380	11	0	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	323	GLU	-	expression tag	UNP P46091
R	324	PHE	-	expression tag	UNP P46091
R	325	LEU	-	expression tag	UNP P46091
R	326	GLU	-	expression tag	UNP P46091
R	327	VAL	-	expression tag	UNP P46091
R	328	LEU	-	expression tag	UNP P46091
R	329	PHE	-	expression tag	UNP P46091
R	330	GLN	-	expression tag	UNP P46091
R	331	GLY	-	expression tag	UNP P46091
R	332	PRO	-	expression tag	UNP P46091
R	333	TRP	-	expression tag	UNP P46091
R	334	SER	-	expression tag	UNP P46091
R	335	HIS	-	expression tag	UNP P46091
R	336	PRO	-	expression tag	UNP P46091
R	337	GLN	-	expression tag	UNP P46091
R	338	PHE	-	expression tag	UNP P46091
R	339	GLU	-	expression tag	UNP P46091
R	340	LYS	-	expression tag	UNP P46091
R	341	GLY	-	expression tag	UNP P46091
R	342	GLY	-	expression tag	UNP P46091
R	343	GLY	-	expression tag	UNP P46091
R	344	SER	-	expression tag	UNP P46091
R	345	GLY	-	expression tag	UNP P46091
R	346	GLY	-	expression tag	UNP P46091
R	347	GLY	-	expression tag	UNP P46091
R	348	SER	-	expression tag	UNP P46091
R	349	GLY	-	expression tag	UNP P46091
R	350	GLY	-	expression tag	UNP P46091

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	351	SER	-	expression tag	UNP P46091
R	352	ALA	-	expression tag	UNP P46091
R	353	TRP	-	expression tag	UNP P46091
R	354	SER	-	expression tag	UNP P46091
R	355	HIS	-	expression tag	UNP P46091
R	356	PRO	-	expression tag	UNP P46091
R	357	GLN	-	expression tag	UNP P46091
R	358	PHE	-	expression tag	UNP P46091
R	359	GLU	-	expression tag	UNP P46091
R	360	LYS	-	expression tag	UNP P46091
R	361	ASP	-	expression tag	UNP P46091
R	362	TYR	-	expression tag	UNP P46091
R	363	LYS	-	expression tag	UNP P46091
R	364	ASP	-	expression tag	UNP P46091
R	365	ASP	-	expression tag	UNP P46091
R	366	ASP	-	expression tag	UNP P46091
R	367	ASP	-	expression tag	UNP P46091
R	368	LYS	-	expression tag	UNP P46091

- Molecule 2 is a protein called Retinoic acid receptor responder protein 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	L	103	Total	C	N	O	S	0	0
			669	421	118	124	6		

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(i) subunit alpha-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	A	211	Total	C	N	O	S	0	0
			1587	1019	267	290	11		

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	CYS	SER	engineered mutation	UNP P63096
A	202	THR	GLY	engineered mutation	UNP P63096
A	203	ALA	GLY	engineered mutation	UNP P63096
A	245	ALA	GLU	engineered mutation	UNP P63096
A	326	SER	ALA	engineered mutation	UNP P63096

- Molecule 4 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	B	330	Total	C	N	O	S	0	0


- Molecule 5 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	C	51	Total	C	N	O	S	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chemerin-like receptor 2



- Molecule 2: Retinoic acid receptor responder protein 2



- Molecule 3: Guanine nucleotide-binding protein G(i) subunit alpha-1






- Molecule 4: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1

Chain B: 



- Molecule 5: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2

Chain C: 



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	59593	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	70	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	R	0.13	0/2471	0.37	0/3380
2	L	0.17	0/677	0.37	0/921
3	A	0.11	0/1615	0.30	0/2187
4	B	0.08	0/2454	0.26	0/3343
5	C	0.09	0/334	0.23	0/462
All	All	0.11	0/7551	0.32	0/10293

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	2391	0	2376	49	0
2	L	669	0	532	20	0
3	A	1587	0	1487	24	0
4	B	2408	0	2237	42	0
5	C	328	0	274	2	0
All	All	7383	0	6906	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:75:THR:HG22	2:L:76:SER:H	1.54	0.71
4:B:71:VAL:HG12	4:B:81:ILE:HG12	1.71	0.71
3:A:185:VAL:HB	3:A:200:ASP:HB3	1.73	0.70
1:R:142:PRO:HB3	3:A:344:ILE:HD11	1.73	0.70
4:B:262:MET:HE3	4:B:264:TYR:HE1	1.57	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	R	294/368 (80%)	262 (89%)	32 (11%)	0	100 100
2	L	91/137 (66%)	81 (89%)	9 (10%)	1 (1%)	12 45
3	A	203/354 (57%)	185 (91%)	18 (9%)	0	100 100
4	B	326/340 (96%)	315 (97%)	11 (3%)	0	100 100
5	C	49/71 (69%)	49 (100%)	0	0	100 100
All	All	963/1270 (76%)	892 (93%)	70 (7%)	1 (0%)	50 79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	60	PHE

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	R	255/333 (77%)	255 (100%)	0	100	100
2	L	52/121 (43%)	52 (100%)	0	100	100
3	A	157/306 (51%)	157 (100%)	0	100	100
4	B	243/283 (86%)	243 (100%)	0	100	100
5	C	24/58 (41%)	24 (100%)	0	100	100
All	All	731/1101 (66%)	731 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	134	GLN
2	L	153	GLN
3	A	269	ASN
3	A	347	ASN
4	B	311	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.