



wwPDB EM Validation Summary Report ⓘ

Dec 22, 2025 – 08:29 PM JST

PDB ID : 9L79 / pdb_00009L79
EMDB ID : EMD-62871
Title : Structure of GPR119-Gs complex bound to AR231453
Authors : Sun, Q.; He, G.; Xu, X.; Liu, X.
Deposited on : 2024-12-26
Resolution : 2.98 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev129
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	13236 (2.48 - 3.48)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	359	10% 71% 25%
2	B	351	92%
3	G	68	13% 82% 16%
4	A	245	7% 76% 13% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	S	259	
6	N	138	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 9461 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucose-dependent insulinotropic receptor.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	270	Total	C	N	O	S	0	0
			2040	1355	328	340	17		

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	-23	MET	-	initiating methionine	UNP Q8TDV5
R	-22	LYS	-	expression tag	UNP Q8TDV5
R	-21	THR	-	expression tag	UNP Q8TDV5
R	-20	ILE	-	expression tag	UNP Q8TDV5
R	-19	ILE	-	expression tag	UNP Q8TDV5
R	-18	ALA	-	expression tag	UNP Q8TDV5
R	-17	LEU	-	expression tag	UNP Q8TDV5
R	-16	SER	-	expression tag	UNP Q8TDV5
R	-15	TYR	-	expression tag	UNP Q8TDV5
R	-14	ILE	-	expression tag	UNP Q8TDV5
R	-13	PHE	-	expression tag	UNP Q8TDV5
R	-12	CYS	-	expression tag	UNP Q8TDV5
R	-11	LEU	-	expression tag	UNP Q8TDV5
R	-10	VAL	-	expression tag	UNP Q8TDV5
R	-9	PHE	-	expression tag	UNP Q8TDV5
R	-8	ALA	-	expression tag	UNP Q8TDV5
R	-7	ASP	-	expression tag	UNP Q8TDV5
R	-6	TYR	-	expression tag	UNP Q8TDV5
R	-5	LYS	-	expression tag	UNP Q8TDV5
R	-4	ASP	-	expression tag	UNP Q8TDV5
R	-3	ASP	-	expression tag	UNP Q8TDV5
R	-2	ASP	-	expression tag	UNP Q8TDV5
R	-1	ASP	-	expression tag	UNP Q8TDV5
R	0	ALA	-	expression tag	UNP Q8TDV5

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	336	Total	C	N	O	S	0	0
			2482	1545	442	475	20		

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	MET	-	initiating methionine	UNP P54311
B	-9	HIS	-	expression tag	UNP P54311
B	-8	HIS	-	expression tag	UNP P54311
B	-7	HIS	-	expression tag	UNP P54311
B	-6	HIS	-	expression tag	UNP P54311
B	-5	HIS	-	expression tag	UNP P54311
B	-4	HIS	-	expression tag	UNP P54311
B	-3	GLY	-	expression tag	UNP P54311
B	-2	SER	-	expression tag	UNP P54311
B	-1	LEU	-	expression tag	UNP P54311
B	0	LEU	-	expression tag	UNP P54311
B	1	GLN	-	expression tag	UNP P54311

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	57	Total	C	N	O	S	0	0
			430	270	74	83	3		

- Molecule 4 is a protein called Guanine nucleotide-binding protein G(s) subunit alpha isoforms short.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	A	219	Total	C	N	O	S	0	0
			1752	1117	319	309	7		

There are 79 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	GLY	-	expression tag	UNP P63092
A	10	CYS	-	expression tag	UNP P63092
A	11	THR	-	expression tag	UNP P63092
A	12	LEU	-	expression tag	UNP P63092
A	13	SER	-	expression tag	UNP P63092
A	14	ALA	-	expression tag	UNP P63092
A	15	GLU	-	expression tag	UNP P63092

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	ASP	-	expression tag	UNP P63092
A	17	LYS	-	expression tag	UNP P63092
A	18	ALA	-	expression tag	UNP P63092
A	19	ALA	-	expression tag	UNP P63092
A	20	VAL	-	expression tag	UNP P63092
A	21	GLU	-	expression tag	UNP P63092
A	22	ARG	-	expression tag	UNP P63092
A	23	SER	-	expression tag	UNP P63092
A	24	LYS	-	expression tag	UNP P63092
A	25	MET	-	expression tag	UNP P63092
A	26	ILE	-	expression tag	UNP P63092
A	27	GLU	-	expression tag	UNP P63092
A	28	LYS	-	expression tag	UNP P63092
A	29	GLN	-	expression tag	UNP P63092
A	30	LEU	-	expression tag	UNP P63092
A	31	GLN	-	expression tag	UNP P63092
A	32	LYS	-	expression tag	UNP P63092
A	33	ASP	-	expression tag	UNP P63092
A	34	LYS	-	expression tag	UNP P63092
A	35	GLN	-	expression tag	UNP P63092
A	36	VAL	-	expression tag	UNP P63092
A	37	TYR	-	expression tag	UNP P63092
A	38	ARG	-	expression tag	UNP P63092
A	39	ALA	-	expression tag	UNP P63092
A	40	THR	-	expression tag	UNP P63092
A	41	HIS	-	expression tag	UNP P63092
A	42	ARG	-	expression tag	UNP P63092
A	43	LEU	-	expression tag	UNP P63092
A	44	LEU	-	expression tag	UNP P63092
A	45	LEU	-	expression tag	UNP P63092
A	46	LEU	-	expression tag	UNP P63092
A	47	GLY	-	expression tag	UNP P63092
A	48	ALA	-	expression tag	UNP P63092
A	49	ASP	-	expression tag	UNP P63092
A	50	ASN	-	expression tag	UNP P63092
A	51	SER	-	expression tag	UNP P63092
A	52	GLY	-	expression tag	UNP P63092
A	53	LYS	-	expression tag	UNP P63092
A	54	SER	-	expression tag	UNP P63092
A	55	THR	-	expression tag	UNP P63092
A	56	ILE	-	expression tag	UNP P63092
A	57	VAL	-	expression tag	UNP P63092

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	189	LYS	-	expression tag	UNP P63092
A	190	GLN	-	expression tag	UNP P63092
A	191	MET	-	expression tag	UNP P63092
A	192	ARG	-	expression tag	UNP P63092
A	193	ILE	-	expression tag	UNP P63092
A	194	LEU	-	expression tag	UNP P63092
A	195	HIS	-	expression tag	UNP P63092
A	196	GLY	-	expression tag	UNP P63092
A	197	GLY	-	expression tag	UNP P63092
A	198	SER	-	expression tag	UNP P63092
A	199	GLY	-	expression tag	UNP P63092
A	200	GLY	-	expression tag	UNP P63092
A	201	SER	-	expression tag	UNP P63092
A	202	GLY	-	expression tag	UNP P63092
A	203	GLY	-	expression tag	UNP P63092
A	249	ASP	ALA	engineered mutation	UNP P63092
A	252	ASP	SER	engineered mutation	UNP P63092
A	?	-	MET	deletion	UNP P63092
A	?	-	VAL	deletion	UNP P63092
A	?	-	ILE	deletion	UNP P63092
A	?	-	ARG	deletion	UNP P63092
A	?	-	GLU	deletion	UNP P63092
A	?	-	ASP	deletion	UNP P63092
A	?	-	ASN	deletion	UNP P63092
A	?	-	GLN	deletion	UNP P63092
A	?	-	THR	deletion	UNP P63092
A	?	-	ASN	deletion	UNP P63092
A	262	ASP	LEU	engineered mutation	UNP P63092
A	362	ALA	ILE	engineered mutation	UNP P63092
A	365	ILE	VAL	engineered mutation	UNP P63092


- Molecule 5 is a protein called scFv16.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	S	233	Total	C	N	O	S	0	0
			1774	1127	293	344	10		

- Molecule 6 is a protein called NB35.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	N	128	Total	C	N	O	S	0	0
			948	593	165	184	6		

- Molecule 7 is N-(2-fluoranyl-4-methylsulfonyl-phenyl)-5-nitro-6-[4-(3-propan-2-yl-1,2,4-oxadiazol-5-yl)piperidin-1-yl]pyrimidin-4-amine (CCD ID: 8WL) (formula: C₂₁H₂₄FN₇O₅S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						AltConf
			Total	C	F	N	O	S	
7	R	1	35	21	1	7	5	1	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucose-dependent insulinotropic receptor




- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1




- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2






- Molecule 4: Guanine nucleotide-binding protein G(s) subunit alpha isoforms short







• Molecule 5: scFv16



• Molecule 6: NB35



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	322758	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1300	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	6.429	Depositor
Minimum map value	-4.005	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.180	Depositor
Recommended contour level	0.65	Depositor
Map size ($\text{\AA}$)	220.16, 220.16, 220.16	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.86, 0.86, 0.86	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8WL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	R	0.17	1/2087 (0.0%)	0.34	0/2842
2	B	0.10	0/2529	0.28	0/3442
3	G	0.05	0/436	0.18	0/590
4	A	0.16	0/1789	0.36	0/2417
5	S	0.10	0/1818	0.30	0/2466
6	N	0.10	0/968	0.30	0/1313
All	All	0.13	1/9627 (0.0%)	0.31	0/13070

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	R	276	PRO	C-O	-5.31	1.17	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	2040	0	2064	8	0
2	B	2482	0	2341	8	0
3	G	430	0	437	1	0
4	A	1752	0	1696	23	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	S	1774	0	1700	14	0
6	N	948	0	884	7	0
7	R	35	0	0	1	0
All	All	9461	0	9122	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 59 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:S:130:GLN:NE2	5:S:217:CYS:SG	2.60	0.74
5:S:40:ALA:HB3	5:S:43:LYS:HB2	1.70	0.73
2:B:42:ARG:NH1	2:B:43:ILE:O	2.22	0.72
2:B:37:ILE:HD11	3:G:38:MET:HE3	1.72	0.70
4:A:250:SER:HB2	4:A:331:ILE:HD13	1.73	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	R	264/359 (74%)	242 (92%)	22 (8%)	0	100	100
2	B	334/351 (95%)	323 (97%)	11 (3%)	0	100	100
3	G	55/68 (81%)	55 (100%)	0	0	100	100
4	A	215/245 (88%)	192 (89%)	23 (11%)	0	100	100
5	S	229/259 (88%)	217 (95%)	12 (5%)	0	100	100
6	N	126/138 (91%)	115 (91%)	11 (9%)	0	100	100
All	All	1223/1420 (86%)	1144 (94%)	79 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	R	212/306 (69%)	211 (100%)	1 (0%)	86	94
2	B	254/293 (87%)	254 (100%)	0	100	100
3	G	45/56 (80%)	45 (100%)	0	100	100
4	A	178/214 (83%)	176 (99%)	2 (1%)	70	86
5	S	193/209 (92%)	193 (100%)	0	100	100
6	N	96/115 (84%)	96 (100%)	0	100	100
All	All	978/1193 (82%)	975 (100%)	3 (0%)	90	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	R	154	GLN
4	A	261	ASN
4	A	262	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
4	A	239	ASN
4	A	257	GLN
6	N	120	GLN
6	N	31	ASN
4	A	41	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	8WL	R	401	-	32,38,38	2.70	9 (28%)	45,56,56	2.93	19 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	8WL	R	401	-	-	1/18/36/36	0/4/4/4

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	R	401	8WL	O32-N30	10.38	1.40	1.22
7	R	401	8WL	C16-N17	5.36	1.46	1.36
7	R	401	8WL	C21-S22	4.44	1.81	1.77
7	R	401	8WL	C10-N11	4.17	1.53	1.46
7	R	401	8WL	C33-N11	3.63	1.52	1.46

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	R	401	8WL	O25-S22-O24	-9.57	101.43	117.92
7	R	401	8WL	C10-C9-C8	7.52	119.93	111.04
7	R	401	8WL	C14-N13-C12	4.86	123.23	111.75
7	R	401	8WL	C29-C12-N13	-4.77	116.14	121.50
7	R	401	8WL	N15-C14-N13	-4.65	121.32	128.60

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	R	401	8WL	C29-C12-N11-C10

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	R	401	8WL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-62871. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 121




Y Index: 133




Z Index: 105


6.3.2 Raw map



X Index: 122



Y Index: 132




Z Index: 102


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.65. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 60 nm³; this corresponds to an approximate mass of 55 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.336 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.336 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.98	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.52	4.08	3.59

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.52 differs from the reported value 2.98 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-62871 and PDB model 9L79. Per-residue inclusion information can be found in section [3](#) on page [9](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.65 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.65).

9.4 Atom inclusion [i](#)



At the recommended contour level, 85% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.65) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8350	0.5490
A	0.8330	0.5180
B	0.9350	0.5990
G	0.6710	0.5240
N	0.8660	0.5620
R	0.7630	0.5190
S	0.8030	0.5460

