



Full wwPDB X-ray Structure Validation Report i

Apr 29, 2025 – 06:16 PM JST

PDB ID : 9LFV / pdb_00009lfv
Title : Crystal structure of mouse RIP3 kinase domain complexed with LK01004
Authors : Xie, H.; Su, H.X.; Li, M.J.; Xu, Y.C.
Deposited on : 2025-01-09
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4924 (2.80-2.76)
Clashscore	180529	5458 (2.80-2.76)
Ramachandran outliers	177936	5386 (2.80-2.76)
Sidechain outliers	177891	5388 (2.80-2.76)
RSRZ outliers	164620	4926 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor-interacting serine/threonine-protein kinase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total 1955	C 1261	N 329	O 357	S 8	0	0	0
1	B	256	Total 1916	C 1232	N 331	O 345	S 8	0	0	0
1	C	250	Total 1753	C 1123	N 302	O 320	S 8	0	0	0
1	D	255	Total 1772	C 1138	N 305	O 323	S 6	0	0	0

There are 60 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	111	ALA	CYS	conflict	UNP Q9QZL0
A	136	ASP	ASN	conflict	UNP Q9QZL0
A	194	LYS	ASP	conflict	UNP Q9QZL0
A	310	LEU	-	expression tag	UNP Q9QZL0
A	311	GLU	-	expression tag	UNP Q9QZL0
A	312	HIS	-	expression tag	UNP Q9QZL0
A	313	HIS	-	expression tag	UNP Q9QZL0
A	314	HIS	-	expression tag	UNP Q9QZL0
A	315	HIS	-	expression tag	UNP Q9QZL0
A	316	HIS	-	expression tag	UNP Q9QZL0
A	317	HIS	-	expression tag	UNP Q9QZL0
A	318	HIS	-	expression tag	UNP Q9QZL0
A	319	HIS	-	expression tag	UNP Q9QZL0
A	320	HIS	-	expression tag	UNP Q9QZL0
A	321	HIS	-	expression tag	UNP Q9QZL0
B	111	ALA	CYS	conflict	UNP Q9QZL0
B	136	ASP	ASN	conflict	UNP Q9QZL0
B	194	LYS	ASP	conflict	UNP Q9QZL0
B	310	LEU	-	expression tag	UNP Q9QZL0
B	311	GLU	-	expression tag	UNP Q9QZL0
B	312	HIS	-	expression tag	UNP Q9QZL0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	313	HIS	-	expression tag	UNP Q9QZL0
B	314	HIS	-	expression tag	UNP Q9QZL0
B	315	HIS	-	expression tag	UNP Q9QZL0
B	316	HIS	-	expression tag	UNP Q9QZL0
B	317	HIS	-	expression tag	UNP Q9QZL0
B	318	HIS	-	expression tag	UNP Q9QZL0
B	319	HIS	-	expression tag	UNP Q9QZL0
B	320	HIS	-	expression tag	UNP Q9QZL0
B	321	HIS	-	expression tag	UNP Q9QZL0
C	111	ALA	CYS	conflict	UNP Q9QZL0
C	136	ASP	ASN	conflict	UNP Q9QZL0
C	194	LYS	ASP	conflict	UNP Q9QZL0
C	310	LEU	-	expression tag	UNP Q9QZL0
C	311	GLU	-	expression tag	UNP Q9QZL0
C	312	HIS	-	expression tag	UNP Q9QZL0
C	313	HIS	-	expression tag	UNP Q9QZL0
C	314	HIS	-	expression tag	UNP Q9QZL0
C	315	HIS	-	expression tag	UNP Q9QZL0
C	316	HIS	-	expression tag	UNP Q9QZL0
C	317	HIS	-	expression tag	UNP Q9QZL0
C	318	HIS	-	expression tag	UNP Q9QZL0
C	319	HIS	-	expression tag	UNP Q9QZL0
C	320	HIS	-	expression tag	UNP Q9QZL0
C	321	HIS	-	expression tag	UNP Q9QZL0
D	111	ALA	CYS	conflict	UNP Q9QZL0
D	136	ASP	ASN	conflict	UNP Q9QZL0
D	194	LYS	ASP	conflict	UNP Q9QZL0
D	310	LEU	-	expression tag	UNP Q9QZL0
D	311	GLU	-	expression tag	UNP Q9QZL0
D	312	HIS	-	expression tag	UNP Q9QZL0
D	313	HIS	-	expression tag	UNP Q9QZL0
D	314	HIS	-	expression tag	UNP Q9QZL0
D	315	HIS	-	expression tag	UNP Q9QZL0
D	316	HIS	-	expression tag	UNP Q9QZL0
D	317	HIS	-	expression tag	UNP Q9QZL0
D	318	HIS	-	expression tag	UNP Q9QZL0
D	319	HIS	-	expression tag	UNP Q9QZL0
D	320	HIS	-	expression tag	UNP Q9QZL0
D	321	HIS	-	expression tag	UNP Q9QZL0

- Molecule 2 is 3-[2-[(2 {S})-butan-2-yl]-1,3-benzothiazol-5-yl]-1- {tert}-butyl-pyrazolo[3,4-d]pyrimidin-4-amine (CCD ID: A1EJO) (formula: C₂₀H₂₄N₆S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	S		
2	A	1	27	20	6	1	0	1

- Molecule 3 is 3-[2-[(2 {R})-butan-2-yl]-1,3-benzothiazol-5-yl]-1- {tert}-butyl-pyrazolo[3,4-d]pyrimidin-4-amine (CCD ID: A1EJP) (formula: C₂₀H₂₄N₆S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	S		
3	A	1	27	20	6	1	0	1
3	B	1	27	20	6	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total C N S 27 20 6 1	0	0
3	D	1	Total C N S 27 20 6 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	7	Total O 7 7	0	0
4	B	5	Total O 5 5	0	0
4	D	2	Total O 2 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Receptor-interacting serine/threonine-protein kinase 3




- Molecule 1: Receptor-interacting serine/threonine-protein kinase 3



- Molecule 1: Receptor-interacting serine/threonine-protein kinase 3



- Molecule 1: Receptor-interacting serine/threonine-protein kinase 3



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.66Å 94.66Å 122.21Å 90.00° 90.98° 90.00°	Depositor
Resolution (Å)	61.10 – 2.78 61.10 – 2.78	Depositor EDS
% Data completeness (in resolution range)	99.0 (61.10-2.78) 99.0 (61.10-2.78)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.50 (at 2.77Å)	Xtriage
Refinement program	PHENIX (1.17.1_3660: ????)	Depositor
R , R_{free}	0.247 , 0.280 0.248 , 0.285	Depositor DCC
R_{free} test set	1537 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	69.9	Xtriage
Anisotropy	0.344	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 73.7	EDS
L-test for twinning ²	$< L > = 0.52$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	0.013 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7545	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1EJP, A1EJO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.33	0/2002	0.61	0/2733
1	B	0.32	1/1960 (0.1%)	0.57	0/2676
1	C	0.29	0/1794	0.55	0/2465
1	D	0.28	0/1816	0.55	0/2501
All	All	0.31	1/7572 (0.0%)	0.57	0/10375

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	152	ASP	C-N	5.33	1.39	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	136	ASP	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	136	ASP	Peptide
1	C	136	ASP	Peptide
1	D	136	ASP	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1955	0	1892	6	0
1	B	1916	0	1857	4	0
1	C	1753	0	1567	4	0
1	D	1772	0	1569	9	0
2	A	27	0	0	0	0
3	A	27	0	0	0	0
3	B	27	0	0	0	0
3	C	27	0	0	0	0
3	D	27	0	0	1	0
4	A	7	0	0	0	0
4	B	5	0	0	0	0
4	D	2	0	0	0	0
All	All	7545	0	6885	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:120:ARG:HD3	1:D:295:VAL:HG11	1.87	0.57
1:C:120:ARG:HD3	1:C:295:VAL:HG11	1.86	0.56
1:A:113:ARG:HB3	1:A:118:LEU:HD11	1.88	0.54
1:C:116:PRO:HB2	1:C:295:VAL:HG22	1.89	0.54
1:D:35:VAL:HG21	1:D:88:VAL:HG11	1.95	0.47
1:C:73:VAL:HG13	1:C:162:PHE:HZ	1.79	0.47
1:A:154:GLU:HG3	1:B:154:GLU:HG3	1.96	0.47
1:A:102:SER:HA	1:A:150:LEU:HA	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:33:PHE:H	1:D:165:SER:HB3	1.81	0.45
1:D:73:VAL:HG13	1:D:162:PHE:HZ	1.81	0.45
1:D:141:HIS:NE2	1:D:160:ALA:O	2.41	0.45
1:C:202:ALA:HB1	1:C:274:ARG:HD2	1.98	0.44
1:D:36:VAL:HG21	3:D:401:A1EJP:N4	2.32	0.44
1:A:156:HIS:NE2	1:B:154:GLU:OE2	2.34	0.44
1:D:79:VAL:HG12	1:D:93:LEU:HD23	1.99	0.44
1:D:103:LEU:HD13	1:D:121:LEU:HD13	1.99	0.43
1:A:28:VAL:HG21	1:A:97:PHE:HE1	1.84	0.43
1:D:288:TYR:HE1	1:D:295:VAL:HG21	1.84	0.42
1:A:103:LEU:HD13	1:A:121:LEU:HD13	2.04	0.40
1:B:124:GLU:OE2	1:B:156:HIS:ND1	2.54	0.40
1:B:212:VAL:O	1:B:216:LEU:HG	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	251/325 (77%)	238 (95%)	12 (5%)	1 (0%)	30 58
1	B	248/325 (76%)	237 (96%)	10 (4%)	1 (0%)	30 58
1	C	240/325 (74%)	229 (95%)	10 (4%)	1 (0%)	30 58
1	D	247/325 (76%)	238 (96%)	8 (3%)	1 (0%)	30 58
All	All	986/1300 (76%)	942 (96%)	40 (4%)	4 (0%)	30 58

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	86	ASP
1	B	86	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	86	ASP
1	C	86	ASP

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	204/283 (72%)	202 (99%)	2 (1%)	73 89
1	B	198/283 (70%)	197 (100%)	1 (0%)	86 95
1	C	162/283 (57%)	160 (99%)	2 (1%)	67 87
1	D	160/283 (56%)	159 (99%)	1 (1%)	84 94
All	All	724/1132 (64%)	718 (99%)	6 (1%)	79 92

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LEU
1	A	152	ASP
1	B	152	ASP
1	C	35	VAL
1	C	152	ASP
1	D	152	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	100	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	A1EJP	D	401	-	24,30,30	0.57	0	27,46,46	1.16	3 (11%)
3	A1EJP	C	401	-	24,30,30	0.62	0	27,46,46	1.06	2 (7%)
3	A1EJP	B	401	-	24,30,30	0.53	0	27,46,46	1.26	5 (18%)
3	A1EJP	A	402[B]	-	24,30,30	0.58	0	27,46,46	1.30	5 (18%)
2	A1EJO	A	401[A]	-	24,30,30	0.65	0	27,46,46	1.06	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1EJP	D	401	-	-	3/13/16/16	0/4/4/4
3	A1EJP	C	401	-	-	2/13/16/16	0/4/4/4
3	A1EJP	B	401	-	-	3/13/16/16	0/4/4/4
3	A1EJP	A	402[B]	-	-	5/13/16/16	0/4/4/4
2	A1EJO	A	401[A]	-	-	2/13/16/16	0/4/4/4

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402[B]	A1EJP	C11-C6-C5	-2.68	116.27	120.65
2	A	401[A]	A1EJO	C9-C8-N5	2.49	113.84	108.04
3	D	401	A1EJP	C9-C8-N5	2.48	113.80	108.04
2	A	401[A]	A1EJO	C6-C7-C8	-2.42	119.32	121.44
3	C	401	A1EJP	C9-C8-N5	2.38	113.57	108.04
3	A	402[B]	A1EJP	C7-C6-C5	2.35	124.67	120.15
3	A	402[B]	A1EJP	C9-C8-N5	2.34	113.49	108.04
3	B	401	A1EJP	C9-C8-N5	2.28	113.34	108.04
3	B	401	A1EJP	C11-C6-C5	-2.27	116.93	120.65
3	C	401	A1EJP	C6-C7-C8	-2.23	119.49	121.44
3	D	401	A1EJP	C6-C7-C8	-2.22	119.49	121.44
3	A	402[B]	A1EJP	C6-C7-C8	-2.22	119.50	121.44
3	B	401	A1EJP	C7-C6-C5	2.21	124.40	120.15
3	B	401	A1EJP	C13-C12-N5	2.19	129.24	125.08
3	B	401	A1EJP	C6-C5-N4	2.16	124.47	120.78
3	D	401	A1EJP	C11-C6-C5	-2.08	117.24	120.65
3	A	402[B]	A1EJP	C18-C17-N3	2.01	111.98	108.80

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401[A]	A1EJO	N4-C5-C6-C7
2	A	401[A]	A1EJO	N4-C5-C6-C11
3	A	402[B]	A1EJP	C20-C17-N3-N4
3	A	402[B]	A1EJP	N4-C5-C6-C7
3	A	402[B]	A1EJP	N4-C5-C6-C11
3	B	401	A1EJP	N4-C5-C6-C7
3	B	401	A1EJP	N4-C5-C6-C11
3	C	401	A1EJP	N4-C5-C6-C7
3	C	401	A1EJP	N4-C5-C6-C11
3	D	401	A1EJP	N4-C5-C6-C7
3	D	401	A1EJP	N4-C5-C6-C11
3	A	402[B]	A1EJP	C16-C13-C14-C15
3	B	401	A1EJP	C16-C13-C14-C15
3	D	401	A1EJP	C16-C13-C14-C15
3	A	402[B]	A1EJP	N5-C12-C13-C16



There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	401	A1EJP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/325 (79%)	0.23	6 (2%) 61 54	33, 59, 88, 112	0
1	B	256/325 (78%)	0.40	8 (3%) 51 46	37, 64, 93, 109	0
1	C	250/325 (76%)	0.89	21 (8%) 18 16	52, 84, 113, 129	0
1	D	255/325 (78%)	1.07	32 (12%) 9 9	51, 89, 118, 137	0
All	All	1020/1300 (78%)	0.64	67 (6%) 26 22	33, 72, 110, 137	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	46	HIS	4.4
1	D	251	PRO	4.2
1	D	254	PRO	3.5
1	D	290	LEU	3.5
1	C	221	ALA	3.4
1	C	273	ASN	3.3
1	D	58	ILE	3.3
1	D	206	TYR	3.2
1	D	210	ILE	3.2
1	A	182	GLY	3.2
1	B	32	GLY	3.1
1	C	295	VAL	3.1
1	D	307	SER	3.0
1	D	255	GLY	3.0
1	D	297	ALA	3.0
1	C	189	PRO	3.0
1	D	245	GLU	2.9
1	D	91	GLN	2.9
1	C	184	LEU	2.9
1	D	142	ARG	2.9
1	B	255	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	140	LEU	2.9
1	C	241	PRO	2.8
1	D	127	LEU	2.8
1	D	280	CYS	2.7
1	A	183	THR	2.7
1	B	165	SER	2.7
1	D	130	CYS	2.7
1	D	33	PHE	2.7
1	D	191	LEU	2.6
1	D	247	PRO	2.6
1	D	249	GLY	2.6
1	C	142	ARG	2.6
1	C	255	GLY	2.6
1	C	61	GLU	2.6
1	C	136	ASP	2.5
1	D	240	ARG	2.5
1	B	167	PHE	2.5
1	D	291	VAL	2.5
1	D	221	ALA	2.5
1	C	249	GLY	2.4
1	D	277	PHE	2.4
1	B	184	LEU	2.4
1	C	245	GLU	2.4
1	B	307	SER	2.4
1	D	278	GLN	2.3
1	C	307	SER	2.3
1	D	276	SER	2.3
1	A	273	ASN	2.3
1	C	265	HIS	2.3
1	D	32	GLY	2.3
1	D	136	ASP	2.3
1	D	92	ALA	2.3
1	C	244	THR	2.2
1	C	162	PHE	2.2
1	D	119	CYS	2.2
1	C	186	TYR	2.2
1	C	87	PHE	2.2
1	C	53	VAL	2.1
1	B	136	ASP	2.1
1	A	31	GLY	2.1
1	A	222	GLU	2.1
1	C	247	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	243	LEU	2.0
1	C	81	GLU	2.0
1	A	305	TYR	2.0
1	D	279	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A1EJP	D	401	27/27	0.73	0.23	41,70,83,92	0
2	A1EJO	A	401[A]	27/27	0.78	0.27	44,51,55,57	27
3	A1EJP	B	401	27/27	0.79	0.19	26,55,66,73	0
3	A1EJP	A	402[B]	27/27	0.79	0.21	43,51,56,58	27
3	A1EJP	C	401	27/27	0.81	0.20	55,73,83,88	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.