



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:01 AM UTC

PDB ID : 6LHX / pdb_00006lhx
Title : Crystal structure of ThsA
Authors : Bae, E.; Ka, D.; Oh, H.
Deposited on : 2019-12-10
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	 16% 83% 14%
1	B	476	 13% 75% 10% 15%
1	C	476	 14% 83% 13%
1	D	476	 17% 72% 12% 16%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 13332 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ThsA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	461	Total 3556	C 2273	N 597	O 677	S 2	Se 7	0	0	0
1	B	406	Total 3170	C 2034	N 531	O 596	S 1	Se 8	0	0	0
1	C	458	Total 3535	C 2262	N 587	O 677	S 2	Se 7	0	0	0
1	D	401	Total 3035	C 1944	N 514	O 569	S 1	Se 7	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP J8G6Z1
A	2	HIS	-	expression tag	UNP J8G6Z1
B	1	GLY	-	expression tag	UNP J8G6Z1
B	2	HIS	-	expression tag	UNP J8G6Z1
C	1	GLY	-	expression tag	UNP J8G6Z1
C	2	HIS	-	expression tag	UNP J8G6Z1
D	1	GLY	-	expression tag	UNP J8G6Z1
D	2	HIS	-	expression tag	UNP J8G6Z1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	15	Total 15	O 15	0	0
2	B	9	Total 9	O 9	0	0
2	C	7	Total 7	O 7	0	0
2	D	5	Total 5	O 5	0	0

● Molecule 1: ThsA



● Molecule 1: ThsA



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	180.42Å 168.58Å 93.61Å 90.00° 106.71° 90.00°	Depositor
Resolution (Å)	31.64 – 2.50 31.64 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.2 (31.64-2.50) 88.5 (31.64-2.50)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.43 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.242 , 0.273 0.246 , 0.277	Depositor DCC
R_{free} test set	1996 reflections (2.17%)	wwPDB-VP
Wilson B-factor (Å ²)	49.5	Xtrriage
Anisotropy	0.179	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 70.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13332	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/3610	0.77	1/4874 (0.0%)
1	B	0.30	0/3219	0.80	3/4339 (0.1%)
1	C	0.29	0/3594	0.76	1/4867 (0.0%)
1	D	0.30	0/3083	0.80	2/4172 (0.0%)
All	All	0.29	0/13506	0.78	7/18252 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	292	VAL	N-CA-C	-6.86	106.35	111.90
1	A	292	VAL	N-CA-C	-5.68	107.28	112.96
1	D	313	LEU	N-CA-C	-5.68	105.17	111.36
1	B	93	LEU	N-CA-C	5.60	118.57	110.28
1	B	385	TYR	CB-CA-C	-5.47	109.80	117.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3556	0	3330	42	0
1	B	3170	0	3017	34	0
1	C	3535	0	3298	36	0
1	D	3035	0	2793	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	15	0	0	0	0
2	B	9	0	0	0	0
2	C	7	0	0	0	0
2	D	5	0	0	0	0
All	All	13332	0	12438	136	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 136 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:142:VAL:HG12	1:C:143:LYS:HG2	1.65	0.77
1:A:76:ARG:NH2	1:A:167:ASP:OD1	2.20	0.74
1:A:69:CYS:SG	1:A:170:LYS:NZ	2.63	0.72
1:B:283:LYS:NZ	1:B:317:ASP:O	2.23	0.71
1:A:228:GLU:HG2	1:A:242:ARG:HB3	1.72	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	443/476 (93%)	430 (97%)	13 (3%)	0	100	100
1	B	392/476 (82%)	377 (96%)	15 (4%)	0	100	100
1	C	448/476 (94%)	431 (96%)	17 (4%)	0	100	100
1	D	385/476 (81%)	371 (96%)	13 (3%)	1 (0%)	36	55
All	All	1668/1904 (88%)	1609 (96%)	58 (4%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	419	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	356/417 (85%)	352 (99%)	4 (1%)	65 84
1	B	323/417 (78%)	320 (99%)	3 (1%)	70 87
1	C	355/417 (85%)	352 (99%)	3 (1%)	73 88
1	D	293/417 (70%)	289 (99%)	4 (1%)	59 81
All	All	1327/1668 (80%)	1313 (99%)	14 (1%)	65 84

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	314	ILE
1	C	409	GLU
1	D	423	ILE
1	D	211	ARG
1	D	312	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	432	ASN
1	D	2	HIS
1	D	411	ASN
1	D	265	ASN
1	D	267	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	453/476 (95%)	1.06	74 (16%) 4 3	45, 77, 119, 138	0
1	B	398/476 (83%)	1.00	64 (16%) 4 4	44, 78, 114, 135	0
1	C	450/476 (94%)	1.09	69 (15%) 5 4	44, 82, 121, 145	0
1	D	394/476 (82%)	1.23	81 (20%) 2 2	51, 88, 128, 160	0
All	All	1695/1904 (89%)	1.09	288 (16%) 4 3	44, 81, 121, 160	0

The worst 5 of 288 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	457	ALA	6.6
1	C	453	LEU	5.9
1	D	458	LEU	5.8
1	B	442	ALA	5.6
1	B	458	LEU	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.