

wwPDB EM Validation Summary Report (i)

Oct 27, 2024 – 01:31 PM JST

PDB ID : 6LID

EMDB ID : EMD-0904

Title: Heteromeric amino acid transporter b0,+AT-rBAT complex

Authors: Yan, R.H.; Li, Y.N.; Lei, J.L.; Zhou, Q.

Deposited on : 2019-12-10

Resolution : 2.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : 1.9.13

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
-		200	32%		
1	A	699	82%	7%	11%
1		COO	32%		
1	С	699	82%	7%	11%
2	В	507	90%		
	D	307	81% 90%	9%	• 10%
2	D	507		100/	100/
	D	307	80%	10%	10%
3	Е	2	50% 50%		
		_	100%		
3	F	2	100%		
			100%		
3	G	2	100%		
			100%		
3	Н	2	50% 50%		



 $Continued\ from\ previous\ page...$

Mol	Chain	Length	Quality of	of chain
			100%	6
3	I	2	50%	50%
			100%	6
3	J	2	50%	50%
			100%	6
3	K	2	100%	ó
			100%	0
3	${ m L}$	2	100%	6
			100%	6
3	M	2	50%	50%
			100%	6
3	N	2	50%	50%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 17964 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Neutral and basic amino acid transport protein rBAT.

Mol	Chain	Residues	\mathbf{Atoms}			AltConf	Trace		
1	A	623	Total 5084	C 3248	N 859	O 952	S 25	0	0
1	С	623	Total 5084	C 3248	N 859	O 952	S 25	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	expression tag	UNP Q07837
A	-12	ALA	-	expression tag	UNP Q07837
A	-11	HIS	-	expression tag	UNP Q07837
A	-10	HIS	-	expression tag	UNP Q07837
A	-9	HIS	-	expression tag	UNP Q07837
A	-8	HIS	-	expression tag	UNP Q07837
A	-7	HIS	-	expression tag	UNP Q07837
A	-6	HIS	-	expression tag	UNP Q07837
A	-5	HIS	-	expression tag	UNP Q07837
A	-4	HIS	-	expression tag	UNP Q07837
A	-3	HIS	-	expression tag	UNP Q07837
A	-2	HIS	-	expression tag	UNP Q07837
A	-1	SER	-	expression tag	UNP Q07837
A	0	GLY	-	expression tag	UNP Q07837
A	1	ARG	-	expression tag	UNP Q07837
С	-13	MET	-	expression tag	UNP Q07837
С	-12	ALA	-	expression tag	UNP Q07837
С	-11	HIS	-	expression tag	UNP Q07837
С	-10	HIS	-	expression tag	UNP Q07837
С	-9	HIS	-	expression tag	UNP Q07837
С	-8	HIS	-	expression tag	UNP Q07837
С	-7	HIS	-	expression tag	UNP Q07837
С	-6	HIS	-	expression tag	UNP Q07837
С	-5	HIS	-	expression tag	UNP Q07837
С	-4	HIS	-	expression tag	UNP Q07837
С	-3	HIS	-	expression tag	UNP Q07837



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	HIS	-	expression tag	UNP Q07837
С	-1	SER	-	expression tag	UNP Q07837
С	0	GLY	-	expression tag	UNP Q07837
С	1	ARG	ı	expression tag	UNP Q07837

• Molecule 2 is a protein called b(0,+)-type amino acid transporter 1.

Mol	Chain	Residues	${f Atoms}$			AltConf	Trace		
2	В	458		C 2368		O 607	S 20	0	0
2	D	458	Total 3538	_		O 607	S 20	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-19	MET	-	expression tag	UNP P82251
В	-18	ALA	_	expression tag	UNP P82251
В	-17	ASP	_	expression tag	UNP P82251
В	-16	TYR	-	expression tag	UNP P82251
В	-15	LYS	-	expression tag	UNP P82251
В	-14	ASP	-	expression tag	UNP P82251
В	-13	ASP	-	expression tag	UNP P82251
В	-12	ASP	-	expression tag	UNP P82251
В	-11	ASP	-	expression tag	UNP P82251
В	-10	LYS	-	expression tag	UNP P82251
В	-9	SER	-	expression tag	UNP P82251
В	-8	GLY	-	expression tag	UNP P82251
В	-7	PRO	-	expression tag	UNP P82251
В	-6	ASP	-	expression tag	UNP P82251
В	-5	GLU	-	expression tag	UNP P82251
В	-4	VAL	-	expression tag	UNP P82251
В	-3	ASP	-	expression tag	UNP P82251
В	-2	ALA	-	expression tag	UNP P82251
В	-1	SER	-	expression tag	UNP P82251
В	0	GLY	-	expression tag	UNP P82251
В	1	ARG	-	expression tag	UNP P82251
D	-19	MET	-	expression tag	UNP P82251
D	-18	ALA	-	expression tag	UNP P82251
D	-17	ASP	-	expression tag	UNP P82251
D	-16	TYR	-	expression tag	UNP P82251
D	-15	LYS	-	expression tag	UNP P82251



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-14	ASP	-	expression tag	UNP P82251
D	-13	ASP	-	expression tag	UNP P82251
D	-12	ASP	-	expression tag	UNP P82251
D	-11	ASP	-	expression tag	UNP P82251
D	-10	LYS	-	expression tag	UNP P82251
D	-9	SER	-	expression tag	UNP P82251
D	-8	GLY	-	expression tag	UNP P82251
D	-7	PRO	-	expression tag	UNP P82251
D	-6	ASP	-	expression tag	UNP P82251
D	-5	GLU	-	expression tag	UNP P82251
D	-4	VAL	-	expression tag	UNP P82251
D	-3	ASP	-	expression tag	UNP P82251
D	-2	ALA	-	expression tag	UNP P82251
D	-1	SER	-	expression tag	UNP P82251
D	0	GLY	-	expression tag	UNP P82251
D	1	ARG	-	expression tag	UNP P82251

 \bullet Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	AltConf	Trace
3	Е	2	Total C N O 28 16 2 10	0	0
3	F	2	Total C N O 28 16 2 10	0	0
3	G	2	Total C N O 28 16 2 10	0	0
3	Н	2	Total C N O 28 16 2 10	0	0
3	I	2	Total C N O 28 16 2 10	0	0
3	J	2	Total C N O 28 16 2 10	0	0
3	K	2	Total C N O 28 16 2 10	0	0
3	L	2	Total C N O 28 16 2 10	0	0




Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf	Trace
3	M	2	Total C N O 28 16 2 10	0	0
3	N	2	Total C N O 28 16 2 10	0	0

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	AltConf
4	A	1	Total Ca 1 1	0
4	С	1	Total Ca 1 1	0

• Molecule 5 is 1,2-DIACYL-GLYCEROL-3-SN-PHOSPHATE (three-letter code: 3PH) (formula: $C_{39}H_{77}O_8P$).



Mol	Chain	Residues	Atoms	AltConf
5	В	1	Total C O P	0
9	Ъ	1	48 39 8 1	U
5	В	1	Total C O P	0
9	Б	1	48 39 8 1	U
5	D	1	Total C O P	0
9	ט	1	48 39 8 1	U
5	D	1	Total C O P	0
	ע	1	48 39 8 1	U

• Molecule 6 is water.




Mol	Chain	Residues	Atoms	AltConf
6	A	116	Total O 116 116	0
6	В	7	Total O 7 7	0
6	С	116	Total O 116 116	0
6	D	7	Total O 7 7	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Neutral and basic amino acid transport protein rBAT












 \bullet Molecule 2: b(0,+)-type amino acid transporter 1

Chaim B:

| 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |





 \bullet Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



		100%			
Chain E:	50%		50%		
NAG2 NAG2					
• Molecule 3 opyranose	3: 2-acetamido-2-deoxy-	-beta-D-glucopyr	ranose-(1-4)-2-a	cetamido-2-deox	cy-beta-D-gluc
Chain F:		100%			
Cham F:		100%			
NAG1					
• Molecule 3 opyranose	3: 2-acetamido-2-deoxy-	-beta-D-glucopyı	ranose-(1-4)-2-ac	cetamido-2-deox	cy-beta-D-gluc
Olasia O		100%			
Chain G:		100%			
NAG1 NAG2 NAG2					
• Molecule 3 opyranose	3: 2-acetamido-2-deoxy-	-beta-D-glucopy	ranose-(1-4)-2-ac	cetamido-2-deox	cy-beta-D-gluc
C1		100%			
Chain H:	50%		50%		
NAG1 NAG2 NAG2					
• Molecule apyranose	3: 2-acetamido-2-deoxy-		ranose-(1-4)-2-ao	cetamido-2-deox	cy-beta-D-gluc
Chain I:	50%	100%	50%		
Cham i.	30 %		30 %		
NAG1 WAG2 WAG2					
• Molecule : opyranose	3: 2-acetamido-2-deoxy-	-beta-D-glucopyı	ranose-(1-4)-2-ac	cetamido-2-deox	cy-beta-D-gluc
O1		100%			
Chain J:	50%		50%		
NAG2 •					



• Molecule 3:	$2\hbox{-acetamido-}2\hbox{-deoxy-beta-D-glucopyranose-} (1\hbox{-}4)\hbox{-}$	2-acetamido-2-deoxy-beta-D-gluo
opyranose		

Chain K: 100%



 \bullet Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L: 100%



 \bullet Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M: 50% 50%



 $\bullet \ \, \text{Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-acetamido$

Chain N: 50% 50%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	127377	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.285	Depositor
Minimum map value	-0.141	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.08	Depositor
Map size (Å)	349.12, 349.12, 349.12	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.091, 1.091, 1.091	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 3PH, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	\mathbf{angles}
IVIOI	Mioi Chain		# Z > 5	RMSZ	# Z > 5
1	A	0.32	0/5219	0.50	0/7086
1	С	0.32	0/5219	0.50	0/7086
2	В	0.37	0/3633	0.58	0/4968
2	D	0.37	0/3633	0.58	0/4968
All	All	0.34	0/17704	0.53	0/24108

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5084	0	4904	31	0
1	С	5084	0	4904	30	0
2	В	3538	0	3677	34	0
2	D	3538	0	3677	34	0
3	Е	28	0	25	0	0
3	F	28	0	25	0	0
3	G	28	0	25	0	0
3	Н	28	0	25	0	0
3	I	28	0	25	0	0



$\alpha \cdots$	C		
Continued	trom	nremons	naae
Conducta	110110	production	payc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	J	28	0	25	0	0
3	K	28	0	25	0	0
3	L	28	0	25	0	0
3	M	28	0	25	0	0
3	N	28	0	25	0	0
4	A	1	0	0	0	0
4	С	1	0	0	0	0
5	В	96	0	150	5	0
5	D	96	0	150	5	0
6	A	116	0	0	3	0
6	В	7	0	0	0	0
6	С	116	0	0	3	0
6	D	7	0	0	0	0
All	All	17964	0	17712	125	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 125 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance} & (ext{Å}) \end{array}$	$egin{aligned} \operatorname{Clash} \ \operatorname{overlap}\ (\mathring{\mathbf{A}}) \end{aligned}$
2:B:456:HIS:CE1	2:B:457:TYR:HE1	1.36	1.43
2:D:456:HIS:CE1	2:D:457:TYR:HE1	1.39	1.39
2:D:456:HIS:ND1	2:D:457:TYR:CE1	1.89	1.37
2:B:456:HIS:ND1	2:B:457:TYR:CE1	1.89	1.37
2:B:456:HIS:CE1	2:B:457:TYR:CE1	2.21	1.27

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	621/699 (89%)	596 (96%)	25 (4%)	0	100	100
1	С	621/699 (89%)	596 (96%)	25 (4%)	0	100	100
2	В	456/507 (90%)	428 (94%)	27 (6%)	1 (0%)	44	68
2	D	456/507~(90%)	428 (94%)	27 (6%)	1 (0%)	44	68
All	All	2154/2412 (89%)	2048 (95%)	104 (5%)	2 (0%)	50	73

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	В	483	GLU
2	D	483	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	562/629 (89%)	561 (100%)	1 (0%)	92 98
1	\mathbf{C}	562/629 (89%)	561 (100%)	1 (0%)	92 98
2	В	384/427 (90%)	366 (95%)	18 (5%)	22 49
2	D	384/427 (90%)	365 (95%)	19 (5%)	21 47
All	All	1892/2112 (90%)	1853 (98%)	39 (2%)	49 76

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	295	ASP
2	D	431	SER
2	D	323	THR
2	D	372	ASN
2	D	464	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:



Mol	Chain	Res	Type
1	С	220	HIS
1	С	442	ASN
2	D	247	ASN
1	С	598	ASN
1	С	370	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	E	1	1,3	14,14,15	0.65	1 (7%)	17,19,21	1.19	1 (5%)
3	NAG	Е	2	3	14,14,15	0.62	0	17,19,21	0.38	0
3	NAG	F	1	1,3	14,14,15	0.49	0	17,19,21	1.20	2 (11%)
3	NAG	F	2	3	14,14,15	0.85	1 (7%)	17,19,21	2.26	4 (23%)
3	NAG	G	1	1,3	14,14,15	1.11	1 (7%)	17,19,21	1.36	1 (5%)
3	NAG	G	2	3	14,14,15	0.95	1 (7%)	17,19,21	2.23	4 (23%)
3	NAG	Н	1	1,3	14,14,15	0.93	1 (7%)	17,19,21	2.30	3 (17%)
3	NAG	Н	2	3	14,14,15	0.61	0	17,19,21	0.37	0
3	NAG	I	1	1,3	14,14,15	0.72	0	17,19,21	2.28	3 (17%)
3	NAG	I	2	3	14,14,15	0.46	0	17,19,21	0.38	0
3	NAG	J	1	1,3	14,14,15	0.64	1 (7%)	17,19,21	1.18	1 (5%)
3	NAG	J	2	3	14,14,15	0.62	0	17,19,21	0.37	0



Mol	Tuno	Chain	Res	Link	Вс	ond leng	ths	Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	K	1	1,3	14,14,15	0.51	0	17,19,21	1.20	2 (11%)
3	NAG	K	2	3	14,14,15	0.85	1 (7%)	17,19,21	2.26	4 (23%)
3	NAG	L	1	1,3	14,14,15	1.12	1 (7%)	17,19,21	1.36	1 (5%)
3	NAG	L	2	3	14,14,15	0.96	1 (7%)	17,19,21	2.23	4 (23%)
3	NAG	M	1	1,3	14,14,15	0.94	1 (7%)	17,19,21	2.30	3 (17%)
3	NAG	M	2	3	14,14,15	0.60	0	17,19,21	0.37	0
3	NAG	N	1	1,3	14,14,15	0.72	0	17,19,21	2.28	3 (17%)
3	NAG	N	2	3	14,14,15	0.46	0	17,19,21	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	Е	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	F	2	3	-	5/6/23/26	0/1/1/1
3	NAG	G	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	G	2	3	-	5/6/23/26	0/1/1/1
3	NAG	Н	1	1,3	-	5/6/23/26	0/1/1/1
3	NAG	Н	2	3	-	2/6/23/26	0/1/1/1
3	NAG	I	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	I	2	3	-	2/6/23/26	0/1/1/1
3	NAG	J	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	J	2	3	-	0/6/23/26	0/1/1/1
3	NAG	K	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	K	2	3	-	5/6/23/26	0/1/1/1
3	NAG	L	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	L	2	3	-	5/6/23/26	0/1/1/1
3	NAG	M	1	1,3	-	5/6/23/26	0/1/1/1
3	NAG	M	2	3	-	2/6/23/26	0/1/1/1
3	NAG	N	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	N	2	3	-	2/6/23/26	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
3	L	1	NAG	O5-C1	3.60	1.49	1.43
3	G	1	NAG	O5-C1	3.57	1.49	1.43
3	L	2	NAG	C1-C2	2.95	1.56	1.52
3	G	2	NAG	C1-C2	2.91	1.56	1.52
3	F	2	NAG	C1-C2	2.56	1.56	1.52

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	Н	1	NAG	C2-N2-C7	7.89	134.14	122.90
3	M	1	NAG	C2-N2-C7	7.88	134.12	122.90
3	I	1	NAG	C2-N2-C7	7.86	134.09	122.90
3	N	1	NAG	C2-N2-C7	7.83	134.06	122.90
3	F	2	NAG	C2-N2-C7	7.75	133.94	122.90

There are no chirality outliers.

5 of 62 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	F	1	NAG	O5-C5-C6-O6
3	K	1	NAG	O5-C5-C6-O6
3	G	2	NAG	C4-C5-C6-O6
3	L	2	NAG	C4-C5-C6-O6
3	F	2	NAG	O5-C5-C6-O6

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















































5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Iol Type Chain Res			Link	Вс	nd leng	ths	Bond angles		
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	3PH	D	701	-	47,47,47	0.91	2 (4%)	51,52,52	1.12	3 (5%)
5	3PH	В	702	-	47,47,47	0.92	2 (4%)	51,52,52	1.21	4 (7%)
5	3PH	D	702	-	47,47,47	0.92	2 (4%)	51,52,52	1.21	4 (7%)



Mol	Type	e Chain Res		Chain	Chain	Ros	Link	Bo	ond leng	ths	В	ond ang	les
MIOI	Moi Type Chain		rtes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2			
5	3PH	В	701	-	47,47,47	0.91	2 (4%)	51,52,52	1.12	3 (5%)			

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	3PH	D	701	-	-	13/49/49/49	-
5	3PH	В	702	-	-	17/49/49/49	-
5	3PH	D	702	-	-	16/49/49/49	-
5	3PH	В	701	-	-	13/49/49/49	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\mathring{A})$	$\operatorname{Ideal}(\text{\AA})$
5	В	702	3PH	O31-C31	4.15	1.45	1.33
5	D	702	3PH	O31-C31	4.13	1.45	1.33
5	В	701	3PH	O31-C31	4.05	1.45	1.33
5	D	701	3PH	O31-C31	4.04	1.45	1.33
5	D	701	3PH	O21-C21	3.96	1.45	1.34

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
5	В	702	3PH	O21-C21-C22	4.50	121.20	111.50
5	D	702	3PH	O21-C21-C22	4.50	121.19	111.50
5	В	701	3PH	O21-C21-C22	3.83	119.75	111.50
5	D	701	3PH	O21-C21-C22	3.80	119.69	111.50
5	D	702	3PH	C2-O21-C21	-3.08	110.20	117.79

There are no chirality outliers.

5 of 59 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	В	701	3PH	C1-O11-P-O12
5	В	702	3PH	C1-O11-P-O12
5	D	701	3PH	C1-O11-P-O12
5	D	702	3PH	C1-O11-P-O12
5	В	701	3PH	C21-C22-C23-C24




There are no ring outliers.


4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	701	3PH	4	0
5	В	702	3PH	2	0
5	D	702	3PH	2	0
5	В	701	3PH	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-0904. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map






The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.08. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 23 nm^3 ; this corresponds to an approximate mass of 20 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



^{*}Reported resolution corresponds to spatial frequency of 0.370 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-0904 and PDB model 6LID. Per-residue inclusion information can be found in section 3 on page 9.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.08 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.08).



9.4 Atom inclusion (i)



At the recommended contour level, 32% of all backbone atoms, 31% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.08) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.3060	0.5240
A	0.5270	0.6260
В	0.0110	0.3950
С	0.5270	0.6260
D	0.0100	0.3900
Е	0.0000	0.2690
F	0.0000	0.2540
G	0.0000	0.1300
Н	0.0000	0.1600
I	0.0710	0.3510
J	0.0000	0.2580
K	0.0000	0.2560
L	0.0000	0.1300
M	0.0000	0.1480
N	0.0710	0.3450



