



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 17, 2025 – 06:13 PM JST

PDB ID : 9LKG / pdb_00009lkg
Title : Ornithine decarboxylase from *Lactocaseibacillus rhamnosus*
Authors : Kim, D.S.; Park, J.H.; Adiko, N.N.; Seo, H.T.
Deposited on : 2025-01-16
Resolution : 2.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.011 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	693	 3% 87% 12%
1	B	693	 4% 87% 11% •
1	C	693	 3% 86% 14%
1	D	693	 3% 87% 12% •

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 22182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ornithine decarboxylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	C	693	5324	3410	912	993	1	8	15	0	0
1	A	693	5324	3410	912	993	1	8	15	0	0
1	B	692	5324	3408	912	995	1	8	15	0	0
1	D	692	5324	3408	912	995	1	8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	17	ASP	ASN	conflict	UNP A0AAN4BJQ6
C	66	TYR	HIS	conflict	UNP A0AAN4BJQ6
C	102	GLU	GLY	conflict	UNP A0AAN4BJQ6
C	455	VAL	ALA	conflict	UNP A0AAN4BJQ6
C	620	ASP	GLU	conflict	UNP A0AAN4BJQ6
C	654	GLU	PRO	conflict	UNP A0AAN4BJQ6
A	17	ASP	ASN	conflict	UNP A0AAN4BJQ6
A	66	TYR	HIS	conflict	UNP A0AAN4BJQ6
A	102	GLU	GLY	conflict	UNP A0AAN4BJQ6
A	455	VAL	ALA	conflict	UNP A0AAN4BJQ6
A	620	ASP	GLU	conflict	UNP A0AAN4BJQ6
A	654	GLU	PRO	conflict	UNP A0AAN4BJQ6
B	17	ASP	ASN	conflict	UNP A0AAN4BJQ6
B	66	TYR	HIS	conflict	UNP A0AAN4BJQ6
B	102	GLU	GLY	conflict	UNP A0AAN4BJQ6
B	455	VAL	ALA	conflict	UNP A0AAN4BJQ6
B	620	ASP	GLU	conflict	UNP A0AAN4BJQ6
B	654	GLU	PRO	conflict	UNP A0AAN4BJQ6
D	17	ASP	ASN	conflict	UNP A0AAN4BJQ6
D	66	TYR	HIS	conflict	UNP A0AAN4BJQ6
D	102	GLU	GLY	conflict	UNP A0AAN4BJQ6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	455	VAL	ALA	conflict	UNP A0AAN4BJQ6
D	620	ASP	GLU	conflict	UNP A0AAN4BJQ6
D	654	GLU	PRO	conflict	UNP A0AAN4BJQ6


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	218	Total O 218 218	0	0
2	A	220	Total O 220 220	0	0
2	B	232	Total O 232 232	0	0
2	D	216	Total O 216 216	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ornithine decarboxylase




- Molecule 1: Ornithine decarboxylase





- Molecule 1: Ornithine decarboxylase





● Molecule 1: Ornithine decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	71.78Å 80.61Å 121.82Å 86.75° 80.61° 70.83°	Depositor
Resolution (Å)	47.25 – 2.09 47.25 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.5 (47.25-2.09) 96.5 (47.25-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.278 , 0.301 0.281 , 0.302	Depositor DCC
R_{free} test set	7220 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	22.4	Xtrriage
Anisotropy	0.181	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.23$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	22182	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 93.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.1645e-09. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	2/5437 (0.0%)	0.85	3/7435 (0.0%)
1	B	0.56	2/5436 (0.0%)	0.87	4/7433 (0.1%)
1	C	0.57	1/5437 (0.0%)	0.85	3/7435 (0.0%)
1	D	0.56	1/5436 (0.0%)	0.85	3/7433 (0.0%)
All	All	0.56	6/21746 (0.0%)	0.85	13/29736 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	6
1	C	0	3
1	D	0	2
All	All	0	15

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	216	HIS	CE1-NE2	5.52	1.38	1.32
1	D	216	HIS	CE1-NE2	5.50	1.38	1.32
1	A	4	HIS	CE1-NE2	5.17	1.37	1.32
1	C	384	HIS	CE1-NE2	5.11	1.37	1.32
1	A	384	HIS	CE1-NE2	5.10	1.37	1.32

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	270	ASN	N-CA-C	-12.03	98.97	112.57
1	D	231	PRO	N-CA-C	6.87	125.22	111.69
1	B	231	PRO	N-CA-C	6.76	125.01	111.69
1	D	68	PRO	N-CA-CB	6.68	110.27	103.25
1	A	231	PRO	N-CA-C	6.47	125.80	112.47

There are no chirality outliers.

5 of 15 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	256	ARG	Sidechain
1	A	533	ARG	Sidechain
1	C	431	ARG	Sidechain
1	C	533	ARG	Sidechain
1	C	616	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5324	0	5219	44	0
1	B	5324	0	5231	50	0
1	C	5324	0	5219	58	0
1	D	5324	0	5231	47	0
2	A	220	0	0	4	0
2	B	232	0	0	4	0
2	C	218	0	0	4	0
2	D	216	0	0	1	0
All	All	22182	0	20900	196	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 196 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:262:ALA:HB2	1:D:269:ALA:HB2	1.46	0.96

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:465:HIS:HD2	1:A:467:LEU:H	1.16	0.94
1:C:465:HIS:HD2	1:C:467:LEU:H	1.16	0.93
1:D:262:ALA:HB2	1:D:269:ALA:CB	1.99	0.93
1:B:465:HIS:HD2	1:B:467:LEU:H	1.15	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	690/693 (100%)	674 (98%)	15 (2%)	1 (0%)	48	51
1	B	689/693 (99%)	669 (97%)	16 (2%)	4 (1%)	22	19
1	C	690/693 (100%)	673 (98%)	15 (2%)	2 (0%)	37	37
1	D	689/693 (99%)	667 (97%)	19 (3%)	3 (0%)	30	29
All	All	2758/2772 (100%)	2683 (97%)	65 (2%)	10 (0%)	30	29

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	67	ALA
1	B	68	PRO
1	D	67	ALA
1	B	69	GLU
1	B	354	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	544/555 (98%)	532 (98%)	12 (2%)	47	53
1	B	547/555 (99%)	532 (97%)	15 (3%)	40	44
1	C	544/555 (98%)	532 (98%)	12 (2%)	47	53
1	D	547/555 (99%)	531 (97%)	16 (3%)	37	41
All	All	2182/2220 (98%)	2127 (98%)	55 (2%)	42	47

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	256	ARG
1	B	548	ASN
1	D	674	THR
1	D	533	ARG
1	B	267	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 83 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	540	GLN
1	D	346	HIS
1	B	584	GLN
1	D	139	HIS
1	D	379	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LLP	B	347	1	23,24,25	0.47	0	25,32,34	0.66	0
1	LLP	C	347	1	23,24,25	0.47	0	25,32,34	0.64	0
1	LLP	D	347	1	23,24,25	0.50	0	25,32,34	0.70	0
1	LLP	A	347	1	23,24,25	0.46	0	25,32,34	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	B	347	1	-	7/16/17/19	0/1/1/1
1	LLP	C	347	1	-	4/16/17/19	0/1/1/1
1	LLP	D	347	1	-	2/16/17/19	0/1/1/1
1	LLP	A	347	1	-	5/16/17/19	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	347	LLP	C4-C4'-NZ-CE
1	A	347	LLP	C4-C4'-NZ-CE
1	B	347	LLP	C4-C4'-NZ-CE
1	B	347	LLP	C5'-OP4-P-OP1
1	B	347	LLP	C5'-OP4-P-OP2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	347	LLP	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	692/693 (99%)	-0.00	21 (3%) 52 54	14, 25, 44, 71	0
1	B	691/693 (99%)	-0.01	25 (3%) 46 48	14, 25, 42, 65	0
1	C	692/693 (99%)	0.01	21 (3%) 52 54	14, 25, 45, 67	0
1	D	691/693 (99%)	-0.03	23 (3%) 49 51	14, 25, 42, 64	0
All	All	2766/2772 (99%)	-0.01	90 (3%) 49 51	14, 25, 44, 71	0

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	68	PRO	13.0
1	B	270	ASN	11.0
1	B	68	PRO	9.4
1	D	269	ALA	8.0
1	B	269	ALA	6.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	C	347	24/25	0.92	0.08	16,20,20,20	15
1	LLP	A	347	24/25	0.92	0.08	16,20,20,20	15
1	LLP	D	347	24/25	0.92	0.12	16,21,24,25	0
1	LLP	B	347	24/25	0.95	0.06	16,20,20,20	15

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.