

# wwPDB NMR Structure Validation Summary Report (i)

Dec 24, 2024 – 04:40 PM EST

PDB ID : 2LN0 BMRB ID : 18142

Title : Structure of MOZ

Authors : Qiu, Y. Deposited on : 2011-12-15

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV :  $\overline{\text{Wang et al.}}$  (2010)

 $\begin{array}{ccc} wwPDB\text{-ShiftChecker} &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$ 

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment is 83%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{array}{c} { m NMR \ archive} \ (\#{ m Entries}) \end{array}$
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	112	71%	24%	• • •



# 2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues				
Well-defined core   Residue range (total)   Backbone RMSD (Å)   Medoid mode				
1	A:207-A:313 (107)	0.91	10	

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 4, 7, 8, 10, 14, 15
2	1, 3, 9, 13, 17, 18
3	5, 6, 12, 16
4	11, 19, 20



# 3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1691 atoms, of which 828 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Histone acetyltransferase KAT6A.

Mol	Chain	Residues	Atoms			Trace			
1	Λ	110	Total	С	Н	N	О	S	0
1	A	110	1687	525	828	156	159	19	U

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	202	HIS	-	expression tag	UNP Q92794
A	203	MET	-	expression tag	UNP Q92794

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms
2	A	4	Total Zn 4 4




# 4 Residue-property plots (i)

#### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Histone acetyltransferase KAT6A



# 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 10. Colouring as in section 4.1 above.

• Molecule 1: Histone acetyltransferase KAT6A





#### Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 500 calculated structures, 20 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
CNS	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1181
Number of shifts mapped to atoms	1175
Number of unparsed shifts	0
Number of shifts with mapping errors	6
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	83%



# 6 Model quality (i)

## 6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	835	804	804	14±3
All	All	16780	16080	16080	286

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

5 of 99 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	$Distance(\mathring{A})$	${f Models}$	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:275:ALA:HB1	1:A:289:HIS:NE2	0.78	1.93	9	1
1:A:252:VAL:HA	1:A:255:LEU:HD23	0.71	1.62	7	6
1:A:227:LEU:HD23	1:A:236:SER:HB2	0.69	1.63	18	3
1:A:264:THR:HG21	1:A:271:GLN:CG	0.69	2.17	5	1
1:A:227:LEU:HD23	1:A:238:HIS:CD2	0.65	2.27	4	2



#### 6.3 Torsion angles (i)

#### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	106/112 (95%)	97±2 (92±2%)	9±2 (8±2%)	0±0 (0±0%)	45	81
All	All	2120/2240 (95%)	1941 (92%)	174 (8%)	5 (0%)	45	81

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	207	PRO	3
1	A	273	LYS	1
1	A	301	PRO	1

#### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric		Percentiles		
1	A	98/103 (95%)	81±3 (83±3%)	17±3 (17±3%)	4	38	
All	All	1960/2060~(95%)	1628 (83%)	332 (17%)	4	38	

5 of 50 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	266	SER	20
1	A	285	ASP	18
1	A	222	LYS	17
1	A	308	GLN	14
1	A	216	LYS	13



#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 6.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

#### 6.7 Other polymers (i)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 83% for the well-defined parts and 81% for the entire structure.

#### 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: assigned\_chem\_shift\_list\_1

#### 7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1181
Number of shifts mapped to atoms	1175
Number of unparsed shifts	0
Number of shifts with mapping errors	6
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

• No matching atom found in the structure. First 5 (of 6) occurrences are reported below.

List ID	Chain	Pog	Type	Atom	Shift Data		
LIST ID	Chain	nes	Type	Atom	Value	Uncertainty	Ambiguity
1	A	202	HIS	CA	56.238		1
1	A	202	HIS	СВ	31.142		1
1	A	203	MET	CA	54.557	•	1
1	A	203	MET	CB	32.624	•	1
1	A	203	MET	Н	8.147	•	1
1	A	203	MET	N	122.382		1

#### 7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, $ppm$	Suggested action
$^{13}\mathrm{C}_{\alpha}$	109	$-0.65 \pm 0.28$	Should be checked
$^{13}\mathrm{C}_{\beta}$	102	$0.24 \pm 0.27$	None needed ( $< 0.5 \text{ ppm}$ )

Continued on next page...



Continued from previous page...

Nucleus	# values	Correction $\pm$ precision, $ppm$	Suggested action
<sup>13</sup> C′	100	$-2.20 \pm 0.31$	Should be applied
$^{15}N$	97	$-0.07 \pm 0.67$	None needed (< 0.5 ppm)

#### 7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 83%, i.e. 1175 atoms were assigned a chemical shift out of a possible 1409. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	512/525 (98%)	209/212 (99%)	207/214 (97%)	96/99 (97%)
Sidechain	633/806 (79%)	440/517 (85%)	186/249 (75%)	7/40 (18%)
Aromatic	30/78 (38%)	28/40 (70%)	0/34~(0%)	2/4~(50%)
Overall	1175/1409 (83%)	677/769 (88%)	393/497 (79%)	105/143 (73%)

#### 7.1.4 Statistically unusual chemical shifts (i)


There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:







# 8 NMR restraints analysis (i)

## 8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1916
Intra-residue ( $ i-j =0$ )	561
Sequential ( $ i-j =1$ )	516
Medium range ( $ i-j >1$ and $ i-j <5$ )	337
Long range ( i-j ≥5)	482
Inter-chain	0
Hydrogen bond restraints	20
Disulfide bond restraints	0
Total dihedral-angle restraints	111
Number of unmapped restraints	0
Number of restraints per residue	17.5
Number of long range restraints per residue <sup>1</sup>	4.3

<sup>&</sup>lt;sup>1</sup>Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

## 8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

## 8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	1.1	0.14
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



## 8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than  $1^\circ$  are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	0.1	1.14
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



# 9 Distance violation analysis (i)

## 9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1~Å are not included in the statistics.

Doctroints type	Count	<b>%</b> <sup>1</sup>	Vio	${f Violated^3}$			Consistently Violated <sup>4</sup>		
Restraints type	Count	70-	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$	
Intra-residue ( i-j =0)	561	29.3	3	0.5	0.2	0	0.0	0.0	
Backbone-Backbone	85	4.4	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	342	17.8	3	0.9	0.2	0	0.0	0.0	
Sidechain-Sidechain	134	7.0	0	0.0	0.0	0	0.0	0.0	
Sequential ( i-j =1)	516	26.9	3	0.6	0.2	1	0.2	0.1	
Backbone-Backbone	207	10.8	1	0.5	0.1	0	0.0	0.0	
Backbone-Sidechain	265	13.8	2	0.8	0.1	1	0.4	0.1	
Sidechain-Sidechain	44	2.3	0	0.0	0.0	0	0.0	0.0	
Medium range ( $ i-j >1 \&  i-j <5$ )	337	17.6	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	121	6.3	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	154	8.0	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	62	3.2	0	0.0	0.0	0	0.0	0.0	
Long range ( $ i-j  \ge 5$ )	482	25.2	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	105	5.5	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	247	12.9	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	130	6.8	0	0.0	0.0	0	0.0	0.0	
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Hydrogen bond	20	1.0	0	0.0	0.0	0	0.0	0.0	
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0	
Total	1916	100.0	6	0.3	0.3	1	0.1	0.1	
Backbone-Backbone	518	27.0	1	0.2	0.1	0	0.0	0.0	
Backbone-Sidechain	1028	53.7	5	0.5	0.3	1	0.1	0.1	
Sidechain-Sidechain	370	19.3	0	0.0	0.0	0	0.0	0.0	

<sup>&</sup>lt;sup>1</sup> percentage calculated with respect to the total number of distance restraints, <sup>2</sup> percentage calculated with respect to the number of restraints in a particular restraint category, <sup>3</sup> violated in at least one model, <sup>4</sup> violated in all the models



#### 9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

#### 9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1~Å are not included in the statistics.

Model ID		Nun	nber o	f viola	ations	5	Mean (Å)	Max (Å)	$SD^6$ (Å)	Median (Å)
Wiodei 1D	$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
1	0	1	0	0	0	1	0.12	0.12	0.0	0.12
2	0	1	0	0	0	1	0.13	0.13	0.0	0.13
3	1	1	0	0	0	2	0.11	0.11	0.0	0.11
4	0	1	0	0	0	1	0.11	0.11	0.0	0.11
5	1	2	0	0	0	3	0.11	0.11	0.0	0.11
6	1	2	0	0	0	3	0.12	0.14	0.02	0.11
7	1	1	0	0	0	2	0.11	0.12	0.01	0.11
8	0	1	0	0	0	1	0.12	0.12	0.0	0.12
9	0	2	0	0	0	2	0.12	0.14	0.02	0.12
10	0	1	0	0	0	1	0.12	0.12	0.0	0.12

Continued on next page...




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	;	Mean (Å)	Max (Å)	${ m SD}^6$ (Å)	Median (Å)
Model 1D	$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
11	0	1	0	0	0	1	0.11	0.11	0.0	0.11
12	0	1	0	0	0	1	0.11	0.11	0.0	0.11
13	1	1	0	0	0	2	0.12	0.13	0.02	0.12
14	1	1	0	0	0	2	0.13	0.14	0.01	0.13
15	0	1	0	0	0	1	0.12	0.12	0.0	0.12
16	1	1	0	0	0	2	0.11	0.12	0.01	0.11
17	0	1	0	0	0	1	0.14	0.14	0.0	0.14
18	0	1	0	0	0	1	0.12	0.12	0.0	0.12
19	0	1	0	0	0	1	0.1	0.1	0.0	0.1
20	0	1	0	0	0	1	0.12	0.12	0.0	0.12

 $<sup>^1</sup>$ Intra-residue restraints,  $^2$ Sequential restraints,  $^3$ Medium range restraints,  $^4$ Long range restraints,  $^5$ Inter-chain restraints,  $^6$ Standard deviation

#### 9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



#### 9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1890(IR:558, SQ:513, MR:337, LR:482, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	restra	aints	Fraction of the ensemble		
$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Count <sup>6</sup>	%	
2	1	0	0	0	3	1	5.0	
0	1	0	0	0	1	2	10.0	
0	0	0	0	0	0	3	15.0	
0	0	0	0	0	0	4	20.0	
1	0	0	0	0	1	5	25.0	
0	0	0	0	0	0	6	30.0	
0	0	0	0	0	0	7	35.0	
0	0	0	0	0	0	8	40.0	
0	0	0	0	0	0	9	45.0	
0	0	0	0	0	0	10	50.0	
0	0	0	0	0	0	11	55.0	
0	0	0	0	0	0	12	60.0	
0	0	0	0	0	0	13	65.0	
0	0	0	0	0	0	14	70.0	
0	0	0	0	0	0	15	75.0	
0	0	0	0	0	0	16	80.0	
0	0	0	0	0	0	17	85.0	
0	0	0	0	0	0	18	90.0	
0	0	0	0	0	0	19	95.0	
0	1	0	0	0	1	20	100.0	

 $<sup>^1</sup>$ Intra-residue restraints,  $^2$ Sequential restraints,  $^3$ Medium range restraints,  $^4$ Long range restraints,  $^5$ Inter-chain restraints,  $^6$  Number of models with violations



#### 9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




## 9.4 Most violated distance restraints in the ensemble (i)

## 9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





#### 9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	$oxed{ ext{Models}^1}$	Mean (Å)	$\mathbf{SD}^1$ (Å)	Median (Å)
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB2	20	0.12	0.01	0.12
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB3	20	0.12	0.01	0.12
(1,1494)	1:255:A:LEU:HA	1:255:A:LEU:HG	5	0.1	0.0	0.1
(1,1642)	1:277:A:ASN:HA	1:276:A:ASP:HA	2	0.12	0.02	0.12


<sup>&</sup>lt;sup>1</sup>Number of violated models, <sup>2</sup>Standard deviation

## 9.5 All violated distance restraints (i)

#### 9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





#### 9.5.2 Table: All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)	
(1,1642)	1:277:A:ASN:HA	1:276:A:ASP:HA	9	0.14	
(1,1611)	1:269:A:ARG:HA	1:269:A:ARG:HD2	6	0.14	
(1,1611)	1:269:A:ARG:HA	1:269:A:ARG:HD3	6	0.14	
(1,1343)	1:243:A:LYS:HA	1:243:A:LYS:HD2	14	0.14	
(1,1343)	1:243:A:LYS:HA	1:243:A:LYS:HD3	14	0.14	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB2	17	0.14	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB3	17	0.14	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB2	2	0.13	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB3	2	0.13	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB2	13	0.13	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB3	13	0.13	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB2	1	0.12	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB3	1	0.12	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB2	7	0.12	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB3	7	0.12	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB2	8	0.12	
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB3	8	0.12	

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,733)	1:278:A:MET:H	1:277:A:ASN:HB2	10	0.12



# 10 Dihedral-angle violation analysis (i)


## 10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle true	Count	Count %1		lated	3	Consistently Violated <sup>4</sup>		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
PHI	56	50.5	1	1.8	0.9	0	0.0	0.0
PSI	55	49.5	2	3.6	1.8	0	0.0	0.0
Total	111	100.0	3	2.7	2.7	0	0.0	0.0

 $<sup>^1</sup>$  percentage calculated with respect to total number of dihedral-angle restraints,  $^2$  percentage calculated with respect to number of restraints in a particular dihedral-angle type,  $^3$  violated in at least one model,  $^4$  violated in all the models

#### 10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




## 10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Num	iber o	f violations	Moon (°)	Mov (°)	SD (°)	Modian (°)
Wiodei 1D	PHI	PSI	Total	$Mean (^{\circ})$	$\mathbf{Max} (^{\circ})$	$\mathbf{SD}$ (°)	$\mid$ Median (°) $\mid$
1	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0.0	0.0	0.0	0.0
5	0	1	1	1.14	1.14	0.0	1.14
6	0	1	1	1.03	1.03	0.0	1.03
7	0	0	0	0.0	0.0	0.0	0.0
8	1	0	1	1.11	1.11	0.0	1.11
9	0	0	0	0.0	0.0	0.0	0.0
10	0	0	0	0.0	0.0	0.0	0.0
11	0	0	0	0.0	0.0	0.0	0.0
12	0	0	0	0.0	0.0	0.0	0.0
13	0	0	0	0.0	0.0	0.0	0.0
14	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0.0	0.0	0.0	0.0
16	0	0	0	0.0	0.0	0.0	0.0
17	0	0	0	0.0	0.0	0.0	0.0
18	0	0	0	0.0	0.0	0.0	0.0
19	0	0	0	0.0	0.0	0.0	0.0
20	0	0	0	0.0	0.0	0.0	0.0



#### 10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

## 10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Num	ıber o	f violated restraints	Fractio	n of the ensemble
PHI	PSI	Total	Count <sup>1</sup>	%
1	2	3	1	5.0
0	0	0	2	10.0
0	0	0	3	15.0
0	0	0	4	20.0
0	0	0	5	25.0
0	0	0	6	30.0
0	0	0	7	35.0
0	0	0	8	40.0
0	0	0	9	45.0
0	0	0	10	50.0
0	0	0	11	55.0

Continued on next page...




Continued from previous page...

Num	ber o	f violated restraints	Fractio	n of the ensemble
PHI	PSI	Total	Count <sup>1</sup>	%
0	0	0	12	60.0
0	0	0	13	65.0
0	0	0	14	70.0
0	0	0	15	75.0
0	0	0	16	80.0
0	0	0	17	85.0
0	0	0	18	90.0
0	0	0	19	95.0
0	0	0	20	100.0

<sup>&</sup>lt;sup>1</sup> Number of models with violations

## 10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)



## 10.4 Most violated dihedral-angle restraints in the ensemble (i)

No violations found



## 10.5 All violated dihedral-angle restraints (i)

#### 10.5.1 Histogram: Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.

Data insufficient to plot histogram

#### 10.5.2 Table: All violated dihedral-angle restraints (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,13)	1:219:A:ASN:N	1:219:A:ASN:CA	1:219:A:ASN:C	1:220:A:ARG:N	5	1.14
(1,54)	1:258:A:GLN:C	1:259:A:CYS:N	1:259:A:CYS:CA	1:259:A:CYS:C	8	1.11
(1,64)	1:266:A:SER:N	1:266:A:SER:CA	1:266:A:SER:C	1:267:A:SER:N	6	1.03

