



## wwPDB EM Validation Summary Report ⓘ

Jan 11, 2026 – 01:02 AM JST

PDB ID : 9LUE / pdb\_00009lue  
EMDB ID : EMD-63395  
Title : Cryo-EM structure of the Nipah G head domain in complex with three Fabs  
Authors : Wang, Y.; Deng, Z.; Zhao, H.  
Deposited on : 2025-02-08  
Resolution : 3.03 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : **FAILED**  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4-5-2 with Phenix2.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
EM percentile statistics : **NOT EXECUTED**  
MapQ : **FAILED**  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.47

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	G	426	
2	L	107	
3	A	119	
4	B	111	
5	C	120	
6	D	106	
7	H	119	

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 8693 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoprotein G.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	G	426	3355	2138	566	630	21	0	0

- Molecule 2 is a protein called NIV LN3D3 Fab Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	L	103	793	493	134	162	4	0	0

- Molecule 3 is a protein called NiV S2B10 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	A	118	945	602	156	183	4	0	0

- Molecule 4 is a protein called NiV S2B10 Fab Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	B	108	830	516	145	165	4	0	0

- Molecule 5 is a protein called NiV S1E2 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	C	120	967	612	159	191	5	0	0


- Molecule 6 is a protein called NiV S1E2 Fab Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	D	105	792	501	128	158	5	0	0

- Molecule 7 is a protein called NIV LN3D3 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	H	119	941	599	148	189	5	0	0

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
8	G	1	14	8	1	5	0
8	G	1	14	8	1	5	0
8	G	1	14	8	1	5	0
8	G	1	14	8	1	5	0
8	G	1	14	8	1	5	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycoprotein G




- Molecule 2: NiV LN3D3 Fab Light Chain



- Molecule 3: NiV S2B10 Fab Heavy Chain




- Molecule 4: NiV S2B10 Fab Light Chain



- Molecule 5: NiV S1E2 Fab Heavy Chain





- Molecule 6: NiV S1E2 Fab Light Chain

Chain D: 73% 21% 6%

A horizontal bar chart showing the quality distribution for Chain D. The bar is 73% green, 21% yellow, and 6% orange.



- Molecule 7: NIV LN3D3 Fab Heavy Chain

Chain H: 84% 16%

A horizontal bar chart showing the quality distribution for Chain H. The bar is 84% green and 16% yellow.



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	145635	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	JEOL CRYO ARM 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	40	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	G	0.20	0/3436	0.48	3/4675 (0.1%)
2	L	0.15	0/813	0.43	0/1101
3	A	0.34	0/972	0.56	0/1325
4	B	0.14	0/850	0.37	0/1154
5	C	0.16	0/994	0.39	0/1350
6	D	0.77	2/813 (0.2%)	0.79	3/1104 (0.3%)
7	H	0.35	0/967	0.62	0/1315
All	All	0.32	2/8845 (0.0%)	0.52	6/12024 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
7	H	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	14	SER	C-N	15.59	1.51	1.33
6	D	2	ILE	C-O	-5.26	1.18	1.24

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	14	SER	CA-C-N	8.69	129.94	120.13
6	D	14	SER	C-N-CA	8.69	129.94	120.13
6	D	14	SER	O-C-N	-6.24	113.72	121.64
1	G	392	PRO	CA-N-CD	-5.20	104.72	112.00

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	569	LYS	CA-C-N	5.11	131.29	121.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
7	H	97	TYR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	3355	0	3301	53	0
2	L	793	0	744	14	0
3	A	945	0	900	17	0
4	B	830	0	792	10	0
5	C	967	0	909	15	0
6	D	792	0	767	23	0
7	H	941	0	889	10	0
8	G	70	0	65	0	0
All	All	8693	0	8367	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 137 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:D:11:MET:CE	6:D:19:VAL:HG23	1.62	1.27
6:D:11:MET:HE2	6:D:19:VAL:CG2	1.64	1.27
6:D:11:MET:HE2	6:D:19:VAL:HG23	1.05	1.04
3:A:6:GLU:OE2	3:A:92:CYS:N	2.14	0.80
1:G:414:LEU:HD22	1:G:426:VAL:HG12	1.71	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	424/426 (100%)	400 (94%)	24 (6%)	0	100	100
2	L	101/107 (94%)	95 (94%)	6 (6%)	0	100	100
3	A	116/119 (98%)	109 (94%)	7 (6%)	0	100	100
4	B	106/111 (96%)	98 (92%)	8 (8%)	0	100	100
5	C	118/120 (98%)	111 (94%)	7 (6%)	0	100	100
6	D	103/106 (97%)	99 (96%)	3 (3%)	1 (1%)	13	42
7	H	117/119 (98%)	109 (93%)	7 (6%)	1 (1%)	14	45
All	All	1085/1108 (98%)	1021 (94%)	62 (6%)	2 (0%)	45	74

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
7	H	98	PRO
6	D	8	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	381/381 (100%)	380 (100%)	1 (0%)	91	95
2	L	92/96 (96%)	92 (100%)	0	100	100
3	A	105/105 (100%)	104 (99%)	1 (1%)	73	87
4	B	91/94 (97%)	91 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	C	108/108 (100%)	107 (99%)	1 (1%)	75	88
6	D	90/91 (99%)	84 (93%)	6 (7%)	13	40
7	H	104/104 (100%)	103 (99%)	1 (1%)	73	87
All	All	971/979 (99%)	961 (99%)	10 (1%)	71	87

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	D	12	SER
6	D	14	SER
7	H	3	LYS
6	D	3	VAL
6	D	5	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	77	ASN
4	B	89	GLN
1	G	570	ASN
1	G	600	GLN
2	L	3	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	NAG	G	1701	1	14,14,15	0.30	0	17,19,21	0.49	0
8	NAG	G	1608	1	14,14,15	0.29	0	17,19,21	0.45	0
8	NAG	G	1607	1	14,14,15	0.26	0	17,19,21	0.48	0
8	NAG	G	1606	1	14,14,15	0.55	0	17,19,21	0.54	0
8	NAG	G	1605	1	14,14,15	0.25	0	17,19,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	G	1701	1	-	3/6/23/26	0/1/1/1
8	NAG	G	1608	1	-	3/6/23/26	0/1/1/1
8	NAG	G	1607	1	-	0/6/23/26	0/1/1/1
8	NAG	G	1606	1	-	2/6/23/26	0/1/1/1
8	NAG	G	1605	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


5 of 12 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
8	G	1605	NAG	O5-C5-C6-O6
8	G	1605	NAG	C4-C5-C6-O6
8	G	1605	NAG	C8-C7-N2-C2
8	G	1605	NAG	O7-C7-N2-C2
8	G	1608	NAG	C8-C7-N2-C2



There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.