



wwPDB X-ray Structure Validation Summary Report ⓘ

May 3, 2025 – 11:34 AM EDT

PDB ID : 3MGP / pdb_00003mgp
Title : Binding of Cobalt ions to the Nucleosome Core Particle
Authors : Mohideen, K.; Muhammad, R.; Davey, C.A.
Deposited on : 2010-04-07
Resolution : 2.44 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2124 (2.46-2.42)
Clashscore	180529	2259 (2.46-2.42)
Ramachandran outliers	177936	2244 (2.46-2.42)
Sidechain outliers	177891	2244 (2.46-2.42)
RSRZ outliers	164620	2124 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	5% 61% 12% • 27%
1	E	135	3% 56% 16% • 27%
2	B	102	3% 70% 7% • 23%
2	F	102	% 68% 13% • • 15%
3	C	119	% 78% 8% • 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	119	
4	D	125	
4	H	125	
5	I	147	
6	J	147	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	CO	C	120	-	-	-	X
8	CO	D	123	-	-	-	X
8	CO	D	124	-	-	-	X
8	CO	H	123	-	-	-	X
8	CO	H	124	-	-	-	X
8	CO	I	74	-	-	-	X
8	CO	I	76	-	-	-	X
8	CO	I	77	-	-	-	X
8	CO	I	80	-	-	-	X
8	CO	I	82	-	-	-	X
8	CO	I	83	-	-	-	X
8	CO	I	84	-	-	-	X
8	CO	I	85	-	-	-	X
8	CO	I	86	-	-	-	X
8	CO	I	87	-	-	-	X
8	CO	I	88	-	-	-	X
8	CO	I	89	-	-	-	X
8	CO	I	94	-	-	-	X
8	CO	J	102	-	-	-	X
8	CO	J	74	-	-	-	X
8	CO	J	76	-	-	-	X
8	CO	J	77	-	-	-	X
8	CO	J	78	-	-	-	X
8	CO	J	80	-	-	-	X
8	CO	J	81	-	-	-	X
8	CO	J	82	-	-	-	X
8	CO	J	83	-	-	-	X
8	CO	J	84	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	CO	J	85	-	-	-	X
8	CO	J	86	-	-	-	X
8	CO	J	87	-	-	-	X
8	CO	J	88	-	-	-	X
8	CO	J	89	-	-	-	X
8	CO	J	90	-	-	-	X
8	CO	J	91	-	-	-	X
8	CO	J	92	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 12228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	99	Total	C	N	O	S	0	0	0
			817	515	158	141	3			
1	E	99	Total	C	N	O	S	0	0	0
			817	515	158	141	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	GLY	SEE REMARK 999	UNP P84233
E	102	ALA	GLY	SEE REMARK 999	UNP P84233

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	79	Total	C	N	O	S	0	0	0
			627	395	121	110	1			
2	F	87	Total	C	N	O	S	0	0	0
			703	442	142	118	1			

- Molecule 3 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	104	Total	C	N	O		0	0	0
			804	507	157	140				
3	G	106	Total	C	N	O		0	0	0
			818	516	160	142				

- Molecule 4 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	100	Total	C	N	O	S	0	0	0
			789	495	147	145	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	99	Total	C	N	O	S	0	0	0
			785	493	146	144	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	29	THR	SER	SEE REMARK 999	UNP P02281
H	29	THR	SER	SEE REMARK 999	UNP P02281

- Molecule 5 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	147	Total	C	N	O	P	0	0	0
			3011	1440	546	879	146			

- Molecule 6 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	J	147	Total	C	N	O	P	0	0	0
			3010	1440	543	881	146			

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		
7	D	1	Total	Cl	0	0
			1	1		
7	E	1	Total	Cl	0	0
			1	1		
7	G	1	Total	Cl	0	0
			1	1		

- Molecule 8 is COBALT (II) ION (CCD ID: CO) (formula: Co).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	1	Total	Co	0	0
			1	1		
8	D	2	Total	Co	0	0
			2	2		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	E	1	Total 1	Co 1	0	0
8	H	2	Total 2	Co 2	0	0
8	I	17	Total 17	Co 17	0	0
8	J	20	Total 20	Co 20	0	0

- Molecule 1: Histone H3.2





• Molecule 3: Histone H2A



• Molecule 4: Histone H2B 1.1



• Molecule 4: Histone H2B 1.1



• Molecule 5: DNA (147-MER)



• Molecule 6: DNA (147-MER)





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.50Å 109.94Å 183.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.60 – 2.44 52.60 – 2.44	Depositor EDS
% Data completeness (in resolution range)	91.3 (52.60-2.44) 91.3 (52.60-2.44)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.45Å)	Xtriage
Refinement program	REFMAC Rigid Body	Depositor
R, R_{free}	0.239 , 0.282 0.242 , 0.280	Depositor DCC
R_{free} test set	1466 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	61.2	Xtriage
Anisotropy	0.405	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12228	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.69	0/829	0.96	0/1111
1	E	0.87	0/829	1.02	2/1111 (0.2%)
2	B	0.76	0/634	1.00	0/848
2	F	0.92	1/711 (0.1%)	1.09	0/948
3	C	0.82	0/814	1.03	0/1099
3	G	0.65	0/828	0.96	2/1117 (0.2%)
4	D	0.81	0/800	1.04	4/1070 (0.4%)
4	H	0.67	0/796	0.99	0/1065
5	I	0.30	0/3378	0.94	3/5212 (0.1%)
6	J	0.30	0/3376	0.95	3/5209 (0.1%)
All	All	0.58	1/12995 (0.0%)	0.98	14/18790 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	H	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	50	ILE	CA-CB	5.62	1.61	1.54

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	58	ILE	CB-CA-C	-6.94	102.95	112.04
3	G	116	LEU	CA-C-N	6.76	128.29	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	116	LEU	C-N-CA	6.76	128.29	119.84
4	D	98	LEU	N-CA-C	6.64	118.32	111.14
5	I	-46	DA	P-O3'-C3'	5.87	129.01	120.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	H	28	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	817	0	858	14	0
1	E	817	0	858	14	0
2	B	627	0	663	5	0
2	F	703	0	755	10	0
3	C	804	0	859	7	0
3	G	818	0	877	11	0
4	D	789	0	828	8	0
4	H	785	0	825	8	0
5	I	3011	0	1662	32	0
6	J	3010	0	1663	34	0
7	A	1	0	0	1	0
7	D	1	0	0	0	0
7	E	1	0	0	0	0
7	G	1	0	0	0	0
8	C	1	0	0	0	0
8	D	2	0	0	0	0
8	E	1	0	0	0	0
8	H	2	0	0	0	0
8	I	17	0	0	0	0
8	J	20	0	0	0	0
All	All	12228	0	9848	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:26:ARG:HA	6:J:31:DT:OP1	1.71	0.89
5:I:-69:DA:H2''	5:I:-68:DT:H5''	1.72	0.72
2:F:31:LYS:HE2	2:F:35:ARG:HH21	1.54	0.72
4:D:91:ILE:O	4:D:95:VAL:HG23	1.92	0.68
5:I:14:DG:H3'	5:I:15:DC:O4'	1.94	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/135 (72%)	96 (99%)	1 (1%)	0	100	100
1	E	97/135 (72%)	93 (96%)	4 (4%)	0	100	100
2	B	77/102 (76%)	74 (96%)	2 (3%)	1 (1%)	10	9
2	F	85/102 (83%)	80 (94%)	3 (4%)	2 (2%)	5	2
3	C	102/119 (86%)	94 (92%)	8 (8%)	0	100	100
3	G	104/119 (87%)	98 (94%)	5 (5%)	1 (1%)	13	13
4	D	98/125 (78%)	90 (92%)	5 (5%)	3 (3%)	3	1
4	H	97/125 (78%)	93 (96%)	3 (3%)	1 (1%)	13	13
All	All	757/962 (79%)	718 (95%)	31 (4%)	8 (1%)	12	12

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	18	HIS
4	H	101	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	24	LYS
4	D	121	ALA
2	F	19	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/110 (78%)	84 (98%)	2 (2%)	45	58
1	E	86/110 (78%)	80 (93%)	6 (7%)	12	15
2	B	64/78 (82%)	64 (100%)	0	100	100
2	F	72/78 (92%)	66 (92%)	6 (8%)	9	10
3	C	83/92 (90%)	77 (93%)	6 (7%)	12	14
3	G	84/92 (91%)	83 (99%)	1 (1%)	67	79
4	D	85/105 (81%)	81 (95%)	4 (5%)	22	30
4	H	85/105 (81%)	79 (93%)	6 (7%)	12	14
All	All	645/770 (84%)	614 (95%)	31 (5%)	21	29

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	63	ARG
4	H	84	SER
1	E	134	ARG
4	H	103	LEU
3	G	71	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	110	ASN
4	H	79	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	39	HIS
1	E	76	GLN
2	F	18	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 47 ligands modelled in this entry, 47 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/135 (73%)	0.34	7 (7%) 23 23	42, 59, 89, 103	0
1	E	99/135 (73%)	0.03	4 (4%) 43 43	34, 47, 76, 96	0
2	B	79/102 (77%)	0.28	3 (3%) 44 44	42, 53, 69, 80	0
2	F	87/102 (85%)	-0.11	1 (1%) 77 79	36, 44, 64, 101	0
3	C	104/119 (87%)	0.08	1 (0%) 79 81	33, 50, 65, 80	0
3	G	106/119 (89%)	0.37	4 (3%) 44 44	42, 59, 82, 98	0
4	D	100/125 (80%)	0.33	4 (4%) 43 43	40, 53, 99, 113	0
4	H	99/125 (79%)	0.63	10 (10%) 14 13	48, 60, 101, 118	0
5	I	147/147 (100%)	0.20	2 (1%) 73 75	58, 111, 144, 167	0
6	J	147/147 (100%)	0.16	0 100 100	64, 109, 147, 156	0
All	All	1067/1256 (84%)	0.23	36 (3%) 48 48	33, 59, 135, 167	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	135	ALA	6.2
2	B	102	GLY	6.0
1	E	135	ALA	5.2
3	G	13	LYS	5.0
2	B	101	GLY	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	CO	J	92	1/1	-0.61	0.72	114,114,114,114	1
8	CO	I	83	1/1	-0.59	0.48	211,211,211,211	1
8	CO	J	102	1/1	-0.57	0.64	86,86,86,86	1
8	CO	J	85	1/1	-0.56	0.63	146,146,146,146	1
8	CO	J	90	1/1	-0.40	0.80	26,26,26,26	1
8	CO	I	81	1/1	-0.34	0.39	91,91,91,91	1
8	CO	I	80	1/1	-0.34	0.45	139,139,139,139	1
8	CO	J	89	1/1	-0.31	0.49	133,133,133,133	1
8	CO	J	80	1/1	-0.31	0.66	114,114,114,114	1
8	CO	H	123	1/1	-0.29	0.91	99,99,99,99	1
8	CO	J	91	1/1	-0.29	0.96	93,93,93,93	1
8	CO	D	124	1/1	-0.27	0.70	96,96,96,96	1
8	CO	I	82	1/1	-0.26	0.83	94,94,94,94	1
8	CO	I	86	1/1	-0.19	0.75	106,106,106,106	1
8	CO	I	87	1/1	-0.18	0.92	28,28,28,28	1
8	CO	C	120	1/1	-0.11	0.75	97,97,97,97	1
8	CO	J	84	1/1	-0.04	0.59	132,132,132,132	1
8	CO	J	86	1/1	-0.02	0.91	85,85,85,85	1
8	CO	H	124	1/1	-0.02	0.41	81,81,81,81	1
8	CO	I	89	1/1	0.05	0.41	152,152,152,152	1
8	CO	J	87	1/1	0.06	1.11	101,101,101,101	1
8	CO	J	78	1/1	0.06	0.52	114,114,114,114	1
8	CO	D	123	1/1	0.10	0.45	89,89,89,89	1
8	CO	J	83	1/1	0.11	0.51	105,105,105,105	1
8	CO	I	78	1/1	0.11	0.39	124,124,124,124	1
8	CO	I	94	1/1	0.15	0.69	126,126,126,126	1
8	CO	I	85	1/1	0.18	0.69	76,76,76,76	1
8	CO	J	88	1/1	0.24	0.41	161,161,161,161	1
8	CO	I	88	1/1	0.24	1.62	32,32,32,32	1
8	CO	J	82	1/1	0.25	0.43	107,107,107,107	1
8	CO	I	76	1/1	0.28	0.49	85,85,85,85	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	CO	I	84	1/1	0.28	1.11	48,48,48,48	1
8	CO	J	79	1/1	0.33	0.38	128,128,128,128	1
8	CO	I	77	1/1	0.37	0.48	117,117,117,117	1
8	CO	J	81	1/1	0.42	0.57	91,91,91,91	1
8	CO	J	74	1/1	0.46	0.41	89,89,89,89	1
8	CO	I	74	1/1	0.48	0.56	116,116,116,116	1
8	CO	I	75	1/1	0.58	0.36	97,97,97,97	1
8	CO	J	77	1/1	0.59	0.49	110,110,110,110	1
8	CO	I	79	1/1	0.64	0.29	62,62,62,62	1
8	CO	J	76	1/1	0.74	0.45	65,65,65,65	1
8	CO	J	75	1/1	0.85	0.37	79,79,79,79	1
8	CO	E	136	1/1	0.86	0.38	79,79,79,79	0
7	CL	G	3145	1/1	0.93	0.08	61,61,61,61	0
7	CL	A	3147	1/1	0.93	0.14	59,59,59,59	1
7	CL	D	3146	1/1	0.96	0.06	65,65,65,65	0
7	CL	E	3148	1/1	0.96	0.07	52,52,52,52	1

6.5 Other polymers ⓘ

There are no such residues in this entry.