



wwPDB EM Validation Summary Report i

Dec 13, 2025 – 06:12 PM EST

PDB ID : 9MGE / pdb_00009mge
EMDB ID : EMD-48253
Title : beta-barrel assembly machine from Escherichia coli in an early state of substrate assembly
Authors : Thomson, B.D.; Kahne, D.
Deposited on : 2024-12-10
Resolution : 3.60 Å(reported)
Based on initial models : 5LJO, 6V05

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev129
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY


The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	12797 (3.10 - 4.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain		
5	E	94		85%	10% 5%
6	F	568		11%	87%
7	G	6		100%	

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 22711 atoms, of which 10979 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer membrane protein assembly factor BamA.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
1	A	777	10588	3448	5005	987	1133	15	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	ALA	-	expression tag	UNP P0A940
A	14	GLU	-	expression tag	UNP P0A940
A	15	GLY	-	expression tag	UNP P0A940
A	16	HIS	-	expression tag	UNP P0A940
A	17	HIS	-	expression tag	UNP P0A940
A	18	HIS	-	expression tag	UNP P0A940
A	19	HIS	-	expression tag	UNP P0A940
A	20	HIS	-	expression tag	UNP P0A940
A	21	HIS	-	expression tag	UNP P0A940
A	22	HIS	-	expression tag	UNP P0A940
A	23	HIS	-	expression tag	UNP P0A940
A	512	CYS	ASP	engineered mutation	UNP P0A940

- Molecule 2 is a protein called Outer membrane protein assembly factor BamB.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
2	B	356	5216	1658	2574	454	524	6	0	0

- Molecule 3 is a protein called Outer membrane protein assembly factor BamC.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
3	C	61	872	277	435	74	85	1	0	0

- Molecule 4 is a protein called Outer membrane protein assembly factor BamD.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	D	217	Total	C	H	N	O	S	0	0
			3419	1098	1674	307	333	7		

- Molecule 5 is a protein called Outer membrane protein assembly factor BamE.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	E	89	Total	C	H	N	O	S	0	0
			1370	437	675	121	135	2		

- Molecule 6 is a protein called Outer membrane protein assembly factor BamA.

Mol	Chain	Residues	Atoms						AltConf	Trace
6	F	75	Total	C	H	N	O	S	0	0
			1189	397	590	94	104	4		

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	243	ALA	-	expression tag	UNP P0A940
F	244	TRP	-	expression tag	UNP P0A940
F	245	SER	-	expression tag	UNP P0A940
F	246	HIS	-	expression tag	UNP P0A940
F	247	PRO	-	expression tag	UNP P0A940
F	248	GLN	-	expression tag	UNP P0A940
F	249	PHE	-	expression tag	UNP P0A940
F	250	GLU	-	expression tag	UNP P0A940
F	251	LYS	-	expression tag	UNP P0A940
F	252	GLY	-	expression tag	UNP P0A940
F	253	GLY	-	expression tag	UNP P0A940
F	254	GLY	-	expression tag	UNP P0A940
F	255	SER	-	expression tag	UNP P0A940
F	256	GLY	-	expression tag	UNP P0A940
F	257	GLY	-	expression tag	UNP P0A940
F	258	GLY	-	expression tag	UNP P0A940
F	259	SER	-	expression tag	UNP P0A940
F	260	GLY	-	expression tag	UNP P0A940
F	261	GLY	-	expression tag	UNP P0A940
F	262	SER	-	expression tag	UNP P0A940
F	263	ALA	-	expression tag	UNP P0A940
F	264	TRP	-	expression tag	UNP P0A940
F	265	SER	-	expression tag	UNP P0A940
F	266	HIS	-	expression tag	UNP P0A940
F	267	PRO	-	expression tag	UNP P0A940

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	268	GLN	-	expression tag	UNP P0A940
F	269	PHE	-	expression tag	UNP P0A940
F	270	GLU	-	expression tag	UNP P0A940
F	271	LYS	-	expression tag	UNP P0A940
F	637	CYS	TYR	engineered mutation	UNP P0A940


- Molecule 7 is a protein called Unknown peptide.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	6	Total	C	H	N	O	0	0
			57	18	26	6	7		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Outer membrane protein assembly factor BamA




- Molecule 2: Outer membrane protein assembly factor BamB








- Molecule 3: Outer membrane protein assembly factor BamC




- Molecule 4: Outer membrane protein assembly factor BamD



- Molecule 5: Outer membrane protein assembly factor BamE



- Molecule 6: Outer membrane protein assembly factor BamA



GLY	VAL	THR	ASN	THR
SER	ALA	SER	LEU	VAL
ASP	MET	LEU	GLU	VAL
ASN	TRP	PHE	ALA	ASN
GLU	GLY	SER	ALA	ASP
TYR	ARG	ASN	GLU	GLU
TYR	TYR	PHE	GLY	ASP
TYR	LEU	GLY	VAL	ILE
LYS	TYR	PHE	ILE	ARG
VAL	SER	TYR	VAL	SER
THR	MET	ASN	ASN	ASN
LEU	GLY	ASP	GLY	THR
ASP	GLU	PHE	GLY	THR
ASP	HIS	GLN	GLU	ILE
ALA	ALA	ALA	GLU	ALA
PRO	SER	ASP	GLY	LEU
ASP	THR	ASP	GLY	PHE
ASP	VAL	ASP	GLY	VAL
ASP	VAL	TYR	GLY	ALA
ASP	VAL	ASP	VAL	THR
ASP	PRO	ASP	GLY	GLY
ILE	GLN	GLN	GLY	GLY
ASP	LEU	GLN	GLY	GLY
ASP	SER	ASP	GLY	ASP
ASP	ASN	ASP	GLY	ASP
ASP	SER	TYR	VAL	ASP
PHE	TYR	VAL	GLU	ASP
HIS	VAL	GLN	LYS	VAL
LYS	GLY	GLN	GLY	VAL
TRP	THR	LYS	GLY	ARG
VAL	ASP	ASP	ASP	VAL
VAL	ASP	TYR	ASP	LEU
LEU	LEU	LEU	ASP	LEU
GLY	GLY	GLY	PHE	ASP
ARG	TYR	THR	GLY	GLY
ASN	ASN	ASP	GLY	GLY
TYR	TRP	THR	VAL	VAL
GLY	GLY	GLY	LYS	VAL
GLY	TRP	TRP	VAL	VAL
CYS	CYS	THR	TYR	TYR
GLY	GLY	THR	TYR	TYR
ASN	ASN	ASP	PRO	ILE
TYR	ASN	VAL	ILE	SER
ARG	TRP	THR	ASP	ALA
TRP	GLY	GLY	ILE	LYS
GLY	TRP	GLY	GLY	VAL
GLY	CYS	THR	TYR	TYR
GLY	GLY	THR	PRO	ILE
ASN	ASN	ASP	ILE	ASN
ASP	ASN	TYR	ASP	ASN
GLY	LYS	TYR	ILE	ASN
LEU	LEU	VAL	GLU	GLY
GLY	ASP	ASP	GLU	GLY
GLY	ASP	TYR	GLY	GLY
ASN	ASN	ASP	GLU	GLY
GLU	GLU	GLY	GLY	GLY
MET	PHE	PHE	VAL	VAL
PRO	PRO	ALA	HIS	VAL
PHE	THR	GLY	VAL	ASP
GLY	ASP	TYR	ASP	ASP
GLY	ASP	ASP	LEU	ASP
LYS	TYR	ASP	LEU	ASP
GLU	GLU	GLY	PRO	LYS
SER	ASN	SER	TYR	LEU
PHE	ARG	VAL	VAL	ASP
TYR	VAL	HIS	VAL	ASP
ALA	ASN	ASN	THR	NET
ALA	ASN	SER	ASN	VAL
GLY	LEU	LEU	GLN	GLU
GLY	GLY	GLY	PRO	ASP
SER	SER	GLY	ILE	LYS
VAL	VAL	GLY	VAL	GLY
VAL	VAL	VAL	PRO	ASN
ASP	ASP	ASP	ASP	GLN
GLY	GLY	GLY	GLY	GLY

- Molecule 7: Unknown peptide

Chain G:

There are no outlier residues recorded for this chain.

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	96441	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	51.1	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.255	Depositor
Minimum map value	-0.158	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.023	Depositor
Map size (Å)	265.6, 265.6, 265.6	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.83000004, 0.83000004, 0.83000004	Depositor

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.21	0/5709	0.52	0/7681
2	B	0.27	0/2687	0.60	0/3666
3	C	0.29	0/447	0.65	0/612
4	D	0.32	0/1785	0.64	0/2427
5	E	0.27	0/710	0.54	0/968
6	F	0.43	0/617	0.82	0/833
All	All	0.26	0/11955	0.58	0/16187

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5583	5005	4862	63	0
2	B	2642	2574	2572	50	0
3	C	437	435	434	13	0
4	D	1745	1674	1673	25	0
5	E	695	675	674	6	0
6	F	599	590	587	6	0
7	G	31	26	10	0	0
All	All	11732	10979	10812	159	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:152:ARG:HG3	4:D:152:ARG:HH11	1.68	0.58
2:B:62:ASN:O	2:B:62:ASN:OD1	2.21	0.58
3:C:67:ILE:HD12	3:C:67:ILE:H	1.68	0.58
2:B:375:ASP:O	2:B:377:LYS:NZ	2.36	0.58
4:D:205:TYR:O	4:D:208:THR:HG22	2.04	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	773/798 (97%)	714 (92%)	59 (8%)	0	100 100
2	B	350/392 (89%)	317 (91%)	33 (9%)	0	100 100
3	C	59/320 (18%)	56 (95%)	3 (5%)	0	100 100
4	D	215/226 (95%)	198 (92%)	17 (8%)	0	100 100
5	E	87/94 (93%)	82 (94%)	5 (6%)	0	100 100
6	F	69/568 (12%)	64 (93%)	5 (7%)	0	100 100
All	All	1553/2398 (65%)	1431 (92%)	122 (8%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	533/680 (78%)	531 (100%)	2 (0%)	89	95
2	B	280/321 (87%)	277 (99%)	3 (1%)	70	83
3	C	44/258 (17%)	43 (98%)	1 (2%)	45	68
4	D	180/190 (95%)	178 (99%)	2 (1%)	70	83
5	E	77/82 (94%)	76 (99%)	1 (1%)	65	81
6	F	63/473 (13%)	59 (94%)	4 (6%)	15	44
All	All	1177/2004 (59%)	1164 (99%)	13 (1%)	69	83

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	117	MET
5	E	103	ASN
6	F	778	SER
6	F	726	SER
6	F	728	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	209	GLN
5	E	89	GLN
6	F	789	GLN
5	E	96	ASN
2	B	319	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-48253. These allow visual inspection of the internal detail of the map and identification of artifacts.




Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map




6.1.2 Raw map




The images above show the map projected in three orthogonal directions.

6.2 Central slices


6.2.1 Primary map



X Index: 160



Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 183




Y Index: 145




Z Index: 186


6.3.2 Raw map



X Index: 184



Y Index: 145




Z Index: 186


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [\(i\)](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.023. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [\(i\)](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 92 nm^3 ; this corresponds to an approximate mass of 83 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.278\AA^{-1}

8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.278 \AA^{-1}

8.2 Resolution estimates [\(i\)](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.60	-	-
Author-provided FSC curve	3.56	3.97	3.60
Unmasked-calculated*	3.99	4.28	4.02

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.99 differs from the reported value 3.6 by more than 10 %

9 Map-model fit i


This section contains information regarding the fit between EMDB map EMD-48253 and PDB model 9MGE. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay i




The images above show the 3D surface view of the map at the recommended contour level 0.023 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.023).

9.4 Atom inclusion [\(i\)](#)



At the recommended contour level, 72% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.023) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7380	0.4020
A	0.7450	0.4070
B	0.7190	0.3850
C	0.7190	0.3980
D	0.8110	0.4170
E	0.8560	0.4510
F	0.6580	0.3260
G	0.7740	0.4130

