

Full wwPDB X-ray Structure Validation Report (i)

Feb 27, 2025 - 05:19 PM EST

PDB ID	:	9MGR
Title	:	Crystal structure of PRMT5:MEP50 in complex with MTA and compound 51
Authors	:	Whittington, D.A.
Deposited on	:	2024-12-11
Resolution	:	2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	164625	3436 (2.08-2.04)
Clashscore	180529	3661 (2.08-2.04)
Ramachandran outliers	177936	3649 (2.08-2.04)
Sidechain outliers	177891	3649 (2.08-2.04)
RSRZ outliers	164620	3436 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	645	87%	9% •
2	В	350	31% 75% 10%	15%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7720 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein arginine N-methyltransferase 5.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	624	Total 5146	C 3286	N 892	0 943	$\begin{array}{c} \mathrm{S} \\ \mathrm{25} \end{array}$	0	10	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-7	MET	-	initiating methionine	UNP 014744
А	-6	ASP	-	expression tag	UNP 014744
А	-5	TYR	-	expression tag	UNP 014744
А	-4	LYS	-	expression tag	UNP 014744
А	-3	ASP	-	expression tag	UNP 014744
А	-2	ASP	-	expression tag	UNP 014744
А	-1	ASP	-	expression tag	UNP 014744
А	0	ASP	-	expression tag	UNP 014744
A	1	LYS	-	expression tag	UNP 014744

• Molecule 2 is a protein called Methylosome protein 50.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	В	298	Total 2317	C 1450	N 396	O 457	S 14	0	8	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-7	MET	-	initiating methionine	UNP Q9BQA1
В	-6	HIS	-	expression tag	UNP Q9BQA1
В	-5	HIS	-	expression tag	UNP Q9BQA1
В	-4	HIS	-	expression tag	UNP Q9BQA1
В	-3	HIS	-	expression tag	UNP Q9BQA1
В	-2	HIS	-	expression tag	UNP Q9BQA1
В	-1	HIS	-	expression tag	UNP Q9BQA1



Continued from	previous	page
----------------	----------	------

Chain	Residue	Modelled	Actual	Comment	Reference
В	0	HIS	-	expression tag	UNP Q9BQA1
В	1	HIS	-	expression tag	UNP Q9BQA1

• Molecule 3 is 6-[(1-acetylazetidin-3-yl)amino]-N-[(2R)-2-hydroxy-2-{(3S)-7-[(4-methyl-1,3 -oxazol-5-yl)methoxy]-1,2,3,4-tetrahydroisoquinolin-3-yl}ethyl]-2-(4-methylpiperidin-1-yl)pyrimidine-4-carboxamide (three-letter code: A1BLF) (formula: $C_{32}H_{42}N_8O_5$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	А	1	Total 45	C 32	N 8	O 5	0	0

• Molecule 4 is 5'-DEOXY-5'-METHYLTHIOADENOSINE (three-letter code: MTA) (formula: $C_{11}H_{15}N_5O_3S$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	А	1	Total 20	C 11	N 5	O 3	S 1	0	0

• Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	115	Total O 115 115	0	0
6	В	17	Total O 17 17	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein arginine N-methyltransferase 5







4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants	102.95Å 137.89Å 178.61Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution(A)	109.15 - 2.07	Depositor
Resolution (A)	109.15 - 2.07	EDS
% Data completeness	47.6 (109.15-2.07)	Depositor
(in resolution range)	47.6(109.15-2.07)	EDS
R_{merge}	0.16	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.53 (at 2.07 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
B B.	0.222 , 0.273	Depositor
Π, Π_{free}	0.227 , 0.271	DCC
R_{free} test set	3854 reflections $(4.95%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	35.6	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.37, 36.9	EDS
L-test for $twinning^2$	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7720	wwPDB-VP
Average B, all atoms $(Å^2)$	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, A1BLF, MTA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.29	0/5286	0.60	0/7182
2	В	0.30	0/2369	0.58	0/3231
All	All	0.30	0/7655	0.60	0/10413

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	193[A]	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	5146	0	5044	47	0
2	В	2317	0	2223	29	0
3	А	45	0	0	3	0
4	А	20	0	15	3	0
5	А	56	0	84	5	0
5	В	4	0	6	0	0
6	А	115	0	0	2	0
6	В	17	0	0	0	0
All	All	7720	0	7372	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:B:135[B]:SER:HB2	2:B:180[B]:ASP:OD1	1.56	1.06
1:A:299:LEU:HA	1:A:302:LYS:HE3	1.61	0.80
1:A:189:TRP:CE2	1:A:193[A]:ARG:HD3	2.20	0.75
1:A:94:SER:O	1:A:100[A]:ARG:HD3	1.93	0.68
2:B:157[B]:GLN:O	2:B:157[B]:GLN:HG3	1.97	0.65
1:A:314:PRO:HD2	6:A:881:HOH:O	1.99	0.63
2:B:102:ALA:HB2	2:B:122:TYR:CD1	2.33	0.62
1:A:301:ALA:HB1	1:A:505:ARG:HG2	1.83	0.61
1:A:299:LEU:HD23	1:A:302:LYS:CE	2.31	0.61
1:A:285:GLU:O	1:A:289:GLN:OE1	2.21	0.58
2:B:135[B]:SER:CB	2:B:180[B]:ASP:OD1	2.44	0.58
1:A:100[B]:ARG:HH12	5:A:708:EDO:C1	2.17	0.58
1:A:371:LEU:HD11	1:A:435:GLU:HB2	1.87	0.56
1:A:347:ASP:OD2	1:A:561[A]:GLN:NE2	2.38	0.56
2:B:157[B]:GLN:O	2:B:157[B]:GLN:CG	2.53	0.56
1:A:333:LYS:HE3	1:A:579:TRP:CH2	2.41	0.55
1:A:301:ALA:HB2	1:A:580:PHE:CZ	2.41	0.55
1:A:100[B]:ARG:HH12	5:A:708:EDO:H12	1.71	0.54
2:B:156:GLN:O	2:B:157[B]:GLN:HG2	2.06	0.54
1:A:311:PRO:HG3	1:A:503:VAL:HG23	1.90	0.54
1:A:299:LEU:HD23	1:A:302:LYS:HE2	1.91	0.53
2:B:171:THR:HG21	2:B:216:THR:HA	1.90	0.53
1:A:301:ALA:HA	1:A:304:TYR:HD2	1.74	0.53
1:A:300:PHE:O	1:A:301:ALA:HB3	2.09	0.53
3:A:701:A1BLF:C25	4:A:702:MTA:HCS3	2.39	0.52
1:A:301:ALA:HB2	1:A:580:PHE:CE1	2.44	0.52



Atom-1	Atom-2	Interatomic $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	Clash
		distance (A)	overlap (A)
1:A:07:1HK:0G1	1:A:08:ARG:N	2.39	0.51
1:A:3/1:LEU:HD13	1:A:455:VAL:HG12	1.92	0.51
2:B:30:GLN:HG2	2:B:49:LEU:HD12	1.92	0.51
2:B:102:ALA:HB2	2:B:122:1YR:CE1	2.45	0.51
2:B:107:GLU:HB2	2:B:118:LYS:HE2	1.93	0.51
1:A:299:LEU:HA	1:A:302:LYS:CE	2.37	0.50
2:B:147[A]:ILE:HG22	2:B:147[A]:ILE:O	2.12	0.50
1:A:310:SER:H	5:A:707:EDO:H12	1.77	0.49
1:A:189:TRP:CZ2	1:A:193[A]:ARG:HD3	2.47	0.49
1:A:369:GLY:N	1:A:370:PRO:CD	2.74	0.49
2:B:86:LEU:HA	2:B:95:LEU:O	2.12	0.49
2:B:180[A]:ASP:O	2:B:180[A]:ASP:CG	2.52	0.48
2:B:146:ASP:O	2:B:147[B]:ILE:HB	2.13	0.48
1:A:553:GLY:HA3	1:A:582:ILE:HG22	1.94	0.48
2:B:301:ARG:HD2	2:B:318:TRP:NE1	2.30	0.47
1:A:302:LYS:HB2	5:A:716:EDO:H11	1.96	0.47
1:A:149:SER:OG	1:A:201[A]:ARG:NH2	2.43	0.47
2:B:281:ALA:HB2	2:B:292:ARG:HE	1.79	0.47
1:A:519:PHE:CE1	1:A:534:ARG:HB3	2.50	0.47
2:B:136:SER:N	2:B:180[B]:ASP:OD1	2.48	0.46
1:A:527:ASP:OD1	1:A:528:PRO:HD2	2.14	0.46
1:A:313:GLN:HB3	6:A:801:HOH:O	2.14	0.45
1:A:446:SER:N	1:A:447:PRO:CD	2.79	0.45
3:A:701:A1BLF:C29	4:A:702:MTA:CS	2.95	0.44
1:A:100[B]:ARG:HH12	5:A:708:EDO:H21	1.82	0.44
2:B:121:LYS:HE3	2:B:157[B]:GLN:HG3	1.99	0.43
2:B:254:HIS:CG	2:B:275:SER:HB2	2.53	0.43
2:B:153:ASP:OD2	2:B:156:GLN:HB2	2.18	0.43
1:A:100[B]:ARG:NH1	1:A:100[B]:ARG:HG2	2.32	0.43
1:A:178:GLU:O	1:A:182:GLU:HG3	2.19	0.43
1:A:298:GLU:C	1:A:300:PHE:H	2.22	0.42
1:A:100[B]:ARG:CG	1:A:100[B]:ARG:HH11	2.33	0.42
1:A:301:ALA:O	1:A:305:GLU:HG3	2.20	0.42
1:A:362:MET:SD	1:A:429:ALA:HB2	2.60	0.42
2:B:263:PHE:HA	2:B:270:PHE:O	2.19	0.42
2:B:265:PRO:HB2	2:B:310:HIS:CE1	2.55	0.42
1:A:627:ASN:N	1:A:628:PRO:CD	2.83	0.42
2:B:181:SER:O	2:B:196:ASP:HA	2.20	0.42
1:A:107:MET:SD	1:A:134:LEU:HD12	2.60	0.42
2:B:43:LEU:HD12	2:B:59:TRP:O	2.20	0.41
2:B:147[B]:ILE:HG22	2:B:147[B]:ILE:O	2.19	0.41

Continued from previous page...



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:361:LEU:HD11	1:A:431:ILE:HD12	2.01	0.41
1:A:92:PRO:O	1:A:100[A]:ARG:HG3	2.20	0.41
2:B:35:ARG:NH1	2:B:86:LEU:O	2.54	0.41
1:A:25:GLU:OE1	2:B:53[B]:CYS:SG	2.75	0.41
2:B:289:GLU:OE2	2:B:292:ARG:HB2	2.21	0.41
1:A:232:PRO:O	1:A:235:ILE:HG12	2.21	0.41
3:A:701:A1BLF:C29	4:A:702:MTA:HCS2	2.51	0.41
1:A:97:GLU:OE2	1:A:101:ARG:NH1	2.49	0.40
1:A:311:PRO:HD3	1:A:502:TYR:HA	2.03	0.40
1:A:617:VAL:O	1:A:622:CYS:HA	2.21	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	630/645~(98%)	606~(96%)	24~(4%)	0	100 100
2	В	302/350~(86%)	290~(96%)	8~(3%)	4 (1%)	10 3
All	All	932/995~(94%)	896 (96%)	32 (3%)	4 (0%)	44 23

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	В	147[A]	ILE
2	В	147[B]	ILE
2	В	257[A]	CYS
2	В	257[B]	CYS



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Outliers		Percentiles		
1	А	567/570~(100%)	565 (100%)	2(0%)	89	90	
2	В	262/298~(88%)	262 (100%)	0	100	100	
All	All	829/868~(96%)	827 (100%)	2 (0%)	92	93	

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	178	GLU
1	А	312	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such side chains are listed below:

Mol	Chain	Res	Type
1	А	128	GLN
1	А	140	ASN
1	А	215	ASN
1	А	282	GLN
1	А	309	GLN
1	А	627	ASN
2	В	78	GLN
2	В	111	ASN
2	В	169	GLN
2	В	266	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tuno	Chain	Dog	Link	Bo	ond leng	ths	B	ond ang	gles
	туре	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
5	EDO	А	705	-	3,3,3	0.06	0	2,2,2	0.08	0
5	EDO	А	715	-	3,3,3	0.08	0	2,2,2	0.17	0
5	EDO	А	709	-	3,3,3	0.07	0	$2,\!2,\!2$	0.17	0
5	EDO	А	706	-	3,3,3	0.17	0	2,2,2	0.52	0
5	EDO	А	703	-	3,3,3	0.07	0	2,2,2	0.26	0
4	MTA	А	702	-	18,22,22	0.68	0	18,32,32	0.79	1 (5%)
5	EDO	А	714	-	3,3,3	0.10	0	2,2,2	0.18	0
5	EDO	А	713	-	3,3,3	0.10	0	$2,\!2,\!2$	0.19	0
5	EDO	А	716	-	3,3,3	0.07	0	2,2,2	0.20	0
5	EDO	А	704	-	3,3,3	0.12	0	2,2,2	0.32	0
5	EDO	А	710	-	3,3,3	0.10	0	2,2,2	0.20	0
5	EDO	А	708	-	3,3,3	0.12	0	2,2,2	0.11	0
5	EDO	В	401	-	3,3,3	0.07	0	2,2,2	0.17	0
5	EDO	А	712	-	3,3,3	0.08	0	2,2,2	0.26	0
3	A1BLF	А	701	-	44,50,50	0.74	2 (4%)	53,71,71	1.80	11 (20%)
5	EDO	А	711	-	3,3,3	0.08	0	2,2,2	0.16	0
5	EDO	А	707	-	3,3,3	0.06	0	2,2,2	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	А	705	-	-	0/1/1/1	-
5	EDO	А	715	-	-	0/1/1/1	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	А	709	-	-	1/1/1/1	-
5	EDO	А	706	-	-	1/1/1/1	-
5	EDO	А	703	-	-	1/1/1/1	-
4	MTA	А	702	-	-	1/3/23/23	0/3/3/3
5	EDO	А	714	-	-	0/1/1/1	-
5	EDO	А	713	-	-	0/1/1/1	-
5	EDO	А	716	-	-	0/1/1/1	-
5	EDO	А	704	-	-	1/1/1/1	-
5	EDO	А	710	-	-	1/1/1/1	-
5	EDO	А	708	-	-	0/1/1/1	-
5	EDO	В	401	-	-	1/1/1/1	-
5	EDO	А	712	-	-	1/1/1/1	-
3	A1BLF	А	701	-	_	7/28/57/57	0/6/6/6
5	EDO	A	711	_	_	0/1/1/1	-
5	EDO	A	707	-	-	0/1/1/1	-

Continued from previous page...

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
3	А	701	A1BLF	C26-C28	-2.39	1.36	1.42
3	А	701	A1BLF	C7-N1	2.03	1.40	1.35

AII (12) bond ai	igie ou	thers are i	listed below.			
Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$ $ Ideal $(^{o})$
3	А	701	A1BLF	C15-N5-C7	7.53	121.51	115.85
3	А	701	A1BLF	C25-C26-C28	4.42	132.45	127.17
3	А	701	A1BLF	N5-C7-N2	-3.48	120.09	126.27
3	А	701	A1BLF	C5-N1-C7	-2.74	116.26	121.67
3	А	701	A1BLF	N5-C7-N1	2.70	120.96	117.12
3	А	701	A1BLF	C25-O4-C24	2.57	123.75	117.62
3	А	701	A1BLF	C18-C19-N8	-2.38	106.76	110.85
3	А	701	A1BLF	C11-N4-C10	-2.36	93.04	94.69
4	А	702	MTA	C5-C6-N6	2.22	123.69	120.31
3	А	701	A1BLF	C14-C15-N5	-2.21	119.57	123.41
3	А	701	A1BLF	C14-C8-N2	-2.18	118.83	123.35
3	А	701	A1BLF	C4-N1-C7	-2.15	117.42	121.67

All (12) bond angle outliers are listed below:

There are no chirality outliers.

All (15) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	А	702	MTA	C4'-C5'-S5'-CS
3	А	701	A1BLF	C14-C15-C16-N6
3	А	701	A1BLF	C14-C15-C16-O2
3	А	701	A1BLF	N5-C15-C16-O2
3	А	701	A1BLF	N5-C15-C16-N6
3	А	701	A1BLF	N6-C17-C18-C19
5	А	712	EDO	O1-C1-C2-O2
5	А	704	EDO	O1-C1-C2-O2
5	А	703	EDO	O1-C1-C2-O2
5	А	709	EDO	O1-C1-C2-O2
5	А	710	EDO	O1-C1-C2-O2
5	А	706	EDO	O1-C1-C2-O2
5	В	401	EDO	O1-C1-C2-O2
3	А	701	A1BLF	N5-C7-N1-C4
3	А	701	A1BLF	N2-C7-N1-C4

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	А	702	MTA	3	0
5	А	716	EDO	1	0
5	А	708	EDO	3	0
3	А	701	A1BLF	3	0
5	А	707	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2		$OWAB(Å^2)$	Q < 0.9	
1	А	624/645~(96%)	0.70	80~(12%)	9	9	11, 34, 76, 111	10 (1%)
2	В	298/350~(85%)	1.70	107~(35%)	1	0	18, 53, 77, 111	8 (2%)
All	All	922/995~(92%)	1.03	187 (20%)	3	3	11, 41, 77, 111	18 (1%)

All (187) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	312	LEU	7.5
1	А	299	LEU	6.5
2	В	248	VAL	6.5
1	А	311	PRO	5.8
2	В	249	LEU	5.8
1	А	314	PRO	5.8
2	В	287	LEU	5.6
1	А	301	ALA	5.5
1	А	300	PHE	5.4
1	А	243	PHE	5.2
1	А	297	TYR	5.0
2	В	245	THR	5.0
1	А	277	PHE	4.8
2	В	246	SER	4.7
2	В	116	VAL	4.5
1	А	292	PRO	4.4
1	А	293	PRO	4.4
2	В	41	ALA	4.3
2	В	114	LEU	4.3
2	В	312	LEU	4.2
1	A	302	LYS	4.1
2	В	243	LYS	4.1
2	В	157[A]	GLN	4.0
1	A	147	HIS	4.0



Continued from previous page								
Mol	Chain	Res	Type	RSRZ				
2	В	268	VAL	3.9				
2	В	303	ALA	3.9				
1	А	294	PRO	3.9				
1	А	296	ALA	3.9				
2	В	313	LEU	3.9				
2	В	308	LEU	3.8				
1	А	406	PHE	3.8				
1	А	354	LYS	3.8				
2	В	72	PHE	3.7				
2	В	61	PHE	3.7				
1	А	240	LYS	3.6				
1	А	242	GLY	3.5				
1	А	175	HIS	3.5				
2	В	282	VAL	3.5				
1	А	529	MET	3.5				
2	В	108	LEU	3.5				
2	В	64	PRO	3.5				
2	В	34	ALA	3.4				
1	А	275	LYS	3.4				
2	В	247	CYS	3.4				
1	А	286	TYR	3.4				
1	А	356	THR	3.3				
2	В	27	MET	3.3				
2	В	326	VAL	3.3				
2	В	266	HIS	3.3				
2	В	285	SER	3.3				
2	В	270	PHE	3.2				
2	В	43	LEU	3.2				
2	В	206	ILE	3.2				
2	В	322	VAL	3.2				
1	А	307	TYR	3.2				
1	А	148	SER	3.1				
2	В	90	GLY	3.1				
1	А	174	THR	3.1				
2	В	63	ASP	3.1				
2	В	158	VAL	3.1				
1	А	346	LEU	3.1				
2	В	115	ILE	3.1				
2	В	244	SER	3.1				
2	В	40	GLY	3.0				
1	А	290	ASN	3.0				
2	В	59	TRP	3.0				



Mol	Chain	Res	Type	RSRZ
1	A	283	TYR	3.0
2	B	328	PRO	3.0
$\frac{2}{2}$	B	62	LVS	3.0
$\frac{2}{2}$	B	327	VAL	3.0
1	Δ	$\frac{521}{284}$	LEU	3.0
1		633	TVP	3.0
1	A D	000		2.0
2	D	200		3.0
<u></u>		216	VAL	2.9
1	A	310		2.9
1	A	380		2.9
1	A	527	ASP	2.9
2	В	87	THR	2.9
1	A	404	TRP	2.9
1	A	241	LYS	2.9
1	A	244	PRO	2.8
2	В	106	TRP	2.8
1	А	56	GLN	2.8
1	А	528	PRO	2.8
2	В	110	GLU	2.8
2	В	242	THR	2.8
2	В	218	LEU	2.8
2	В	66	ALA	2.8
1	А	287	LEU	2.8
1	А	315	LEU	2.8
2	В	286	SER	2.8
2	В	225	SER	2.7
2	В	306	SER	2.7
2	В	323	VAL	2.7
2	В	112	GLU	2.7
1	А	273	SER	2.7
2	В	155	ALA	2.7
1	А	361	LEU	2.7
1	А	272	HIS	2.7
2	В	226	GLU	2.6
2	В	68	PRO	2.6
1	Ā	281	LEU	2.6
2	B	102	ALA	2.6
2	B	229	VAL	2.6
2	B	269	PRO	$\frac{2.0}{2.6}$
2	B	117	SER	2.0
2	B	288	SER	2.0
2 9	R	60	LEII	2.0
4		00	ULLU	U

Continued from previous page...



Mol	Chain	Res	\mathbf{Tvpe}	RSRZ
2	B	280	LEU	2.6
1	A	13	ARG	2.0
2	B	77	VAL	2.5
$\frac{-2}{2}$	B	36	TYR	2.5
1	A	249	MET	2.5
2	B	263	PHE	2.5
$\frac{2}{2}$	B	200	PHE	2.5
2	B	92	ARG	2.5
2	B	307	PRO	2.5
2	B	239	LEU	2.5
2	B	65	CYS	2.5
1	A	382	ALA	2.5
2	B	262	VAL	2.5
1	A	357	ASN	2.5
2	B	119	PHE	2.5
2	B	105	LEU	2.5
1	A	304	TYR	2.4
1	А	145	GLY	2.4
2	В	154	LEU	2.4
1	A	345	LEU	2.4
1	A	342	TYR	2.4
1	А	526	ARG	2.4
2	В	311	SER	2.4
1	А	177	GLU	2.3
1	А	237	LEU	2.3
2	В	118	LYS	2.3
2	В	310	HIS	2.3
2	В	177	PRO	2.3
2	В	240	VAL	2.3
2	В	137	GLY	2.3
2	В	230	PHE	2.3
2	В	42	LEU	2.3
2	В	138	THR	2.3
2	В	273	SER	2.3
2	В	89	VAL	2.3
1	А	388	LEU	2.3
1	А	68	ARG	2.3
2	В	55	ALA	2.3
2	В	224	GLN	2.3
1	А	245	VAL	2.3
1	А	409	TRP	2.2

VAL Continued on next page...

413

2.2

1

А



Mol	Chain	Res	Type	RSRZ
2	В	227	VAL	2.2
1	А	274	GLU	2.2
2	В	113	THR	2.2
1	А	310	SER	2.2
2	В	212	GLY	2.2
1	А	349	VAL	2.2
2	В	151	VAL	2.2
1	А	255	PHE	2.2
2	В	58	LEU	2.2
2	В	219	ALA	2.2
2	В	31	LEU	2.1
1	А	355	ASP	2.1
2	В	222	PRO	2.1
2	В	94	ILE	2.1
2	В	95	LEU	2.1
2	В	267	SER	2.1
1	А	31	GLY	2.1
1	А	358	VAL	2.1
1	А	530	ILE	2.1
2	В	147[A]	ILE	2.1
1	А	309	GLN	2.1
1	А	278	CYS	2.1
1	А	149	SER	2.1
2	В	274	LEU	2.1
1	А	251	GLN	2.1
1	А	313	GLN	2.1
2	В	33	ALA	2.0
2	В	71	GLY	2.0
1	А	280	TYR	2.0
2	В	121	LYS	2.0
2	В	38	SER	2.0
1	А	176	THR	2.0
1	А	282	GLN	2.0
2	В	223	GLN	2.0
1	А	270	ASN	2.0
1	A	385	ARG	2.0
2	В	297	ARG	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B -factors($Å^2$)	Q<0.9
5	EDO	А	716	4/4	0.62	0.20	44,44,46,46	0
5	EDO	А	706	4/4	0.63	0.29	44,45,45,47	0
5	EDO	А	714	4/4	0.70	0.17	36,36,37,37	0
5	EDO	А	711	4/4	0.78	0.20	42,42,43,43	0
5	EDO	А	708	4/4	0.80	0.18	47,48,48,49	0
5	EDO	А	707	4/4	0.81	0.19	44,44,44,44	0
5	EDO	А	713	4/4	0.82	0.17	38,39,40,41	0
5	EDO	А	709	4/4	0.83	0.16	42,43,43,43	0
5	EDO	В	401	4/4	0.83	0.19	59,59,60,61	0
5	EDO	А	710	4/4	0.84	0.12	36, 36, 36, 37	0
5	EDO	А	703	4/4	0.84	0.20	40,40,40,40	0
5	EDO	А	704	4/4	0.84	0.22	38,38,39,39	0
5	EDO	А	712	4/4	0.89	0.20	34,34,34,35	0
5	EDO	А	715	4/4	0.90	0.14	53,53,54,54	0
5	EDO	A	705	4/4	0.92	0.08	27,28,28,28	0
3	A1BLF	А	701	45/45	0.94	0.08	25,27,40,45	0
4	MTA	А	702	20/20	0.96	0.08	25,26,28,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

