



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2025 – 12:30 AM EST

PDB ID : 9MQN
EMDB ID : EMD-48535
Title : AngV-F Pre-fusion Protein
Authors : Lella, M.; Acharya, P.
Deposited on : 2025-01-04
Resolution : 5.90 Å(reported)
Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev117
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 5.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	541	
1	B	541	
1	C	541	
1	D	541	
1	E	541	
1	F	541	
2	G	4	
2	H	4	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	4	100% 100%
2	J	4	100% 100%
2	K	4	100% 100%
2	L	4	50% 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	D	603	X	-	-	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 21258 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	456	Total	C	N	O	S	0	0
			3464	2194	574	676	20		
1	B	456	Total	C	N	O	S	0	0
			3464	2194	574	676	20		
1	C	456	Total	C	N	O	S	0	0
			3464	2194	574	676	20		
1	D	449	Total	C	N	O	S	0	0
			3412	2163	566	664	19		
1	E	449	Total	C	N	O	S	0	0
			3412	2163	566	664	19		
1	F	449	Total	C	N	O	S	0	0
			3412	2163	566	664	19		

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
2	G	4	Total	C	N	O	0	0
			49	28	2	19		
2	H	4	Total	C	N	O	0	0
			49	28	2	19		
2	I	4	Total	C	N	O	0	0
			49	28	2	19		
2	J	4	Total	C	N	O	0	0
			49	28	2	19		
2	K	4	Total	C	N	O	0	0
			49	28	2	19		
2	L	4	Total	C	N	O	0	0
			49	28	2	19		

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
3	A	1	Total	C	N	O	0
			14	8	1	5	
3	A	1	Total	C	N	O	0
			14	8	1	5	
3	A	1	Total	C	N	O	0
			14	8	1	5	
3	A	1	Total	C	N	O	0
			14	8	1	5	
3	B	1	Total	C	N	O	0
			14	8	1	5	
3	B	1	Total	C	N	O	0
			14	8	1	5	
3	B	1	Total	C	N	O	0
			14	8	1	5	
3	B	1	Total	C	N	O	0
			14	8	1	5	
3	C	1	Total	C	N	O	0
			14	8	1	5	
3	C	1	Total	C	N	O	0
			14	8	1	5	
3	C	1	Total	C	N	O	0
			14	8	1	5	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
3	D	1	Total	C	N	O	0
			14	8	1	5	
3	D	1	Total	C	N	O	0
			14	8	1	5	
3	D	1	Total	C	N	O	0
			14	8	1	5	
3	D	1	Total	C	N	O	0
			14	8	1	5	
3	E	1	Total	C	N	O	0
			14	8	1	5	
3	E	1	Total	C	N	O	0
			14	8	1	5	
3	E	1	Total	C	N	O	0
			14	8	1	5	
3	E	1	Total	C	N	O	0
			14	8	1	5	
3	F	1	Total	C	N	O	0
			14	8	1	5	
3	F	1	Total	C	N	O	0
			14	8	1	5	
3	F	1	Total	C	N	O	0
			14	8	1	5	
3	F	1	Total	C	N	O	0
			14	8	1	5	

Year	Country	Value	Category
2024	Y22	100	Green
2023	K38	95	Yellow
2022	L39	90	Yellow
2021	Y40	85	Yellow
2020	K41	80	Yellow
2019	P46	75	Yellow
2018	D50	70	Yellow
2017	T51	65	Yellow
2016	V52	60	Yellow
2015	M68	55	Yellow
2014	V60	50	Yellow
2013	R92	45	Yellow
2012	N93	40	Yellow
2011	N94	35	Yellow
2010	G98	30	Green
2009	N101	25	Yellow
2008	F105	20	Green
2007	I108	15	Yellow
2006	V109	10	Yellow
2005	A113	5	Yellow
2004	N132	0	Yellow
2003	Q133	-5	Yellow
2002	N134	-10	Yellow
2001	R142	-15	Yellow
2000	D143	-20	Yellow
1999	S144	-25	Yellow
1998	L155	-30	Yellow
1997	S156	-35	Yellow
1996	L157	-40	Yellow
1995	G158	-45	Yellow
1994	M159	-50	Yellow
1993	Q160	-55	Yellow
1992	E161	-60	Yellow
1991	T162	-65	Yellow
1990	L212	-70	Yellow
1989	R213	-75	Yellow
1988	E234	-80	Yellow
1987	T237	-85	Yellow
1986	T242	-90	Yellow
1985	K254	-95	Green
1984	D264	-100	Yellow
1983	I271	-105	Yellow
1982	N275	-110	Yellow
1981	W302	-115	Yellow
1980	L319	-120	Yellow
1979	K329	-125	Yellow
1978	R330	-130	Yellow
1977	S331	-135	Yellow
1976	R336	-140	Yellow
1975	D337	-145	Yellow
1974	Y338	-150	Yellow
1973	M342	-155	Yellow
1972	T354	-160	Yellow
1971	S355	-165	Yellow
1970	R359	-170	Yellow
1969	Y370	-175	Yellow
1968	S373	-180	Yellow
1967	V376	-185	Yellow
1966	I377	-190	Yellow
1965	Y378	-195	Yellow
1964	A379	-200	Yellow
1963	C386	-205	Yellow
1962	I395	-210	Yellow
1961	V396	-215	Yellow
1960	Q397	-220	Yellow
1959	S398	-225	Yellow
1958	T399	-230	Yellow
1957	S400	-235	Yellow
1956	T404	-240	Yellow
1955	M405	-245	Yellow
1954	I406	-250	Yellow
1953	D407	-255	Yellow
1952	L427	-260	Yellow
1951	S431	-265	Yellow
1950	P432	-270	Yellow
1949	M433	-275	Yellow
1948	M434	-280	Yellow
1947	L435	-285	Yellow
1946	T436	-290	Yellow
1945	I437	-295	Yellow
1944	E438	-300	Yellow



• Molecule 1: Fusion protein



• Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%
100%



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%
100%



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  50%
100%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	240865	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE; Micrographs were curated through contrast transfer function (CTF) where greater than 30 angstrom were discarded	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	28	Depositor
Minimum defocus (nm)	1700	Depositor
Maximum defocus (nm)	2800	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	3.024	Depositor
Minimum map value	-0.757	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.086	Depositor
Recommended contour level	0.847	Depositor
Map size (\AA)	492.80002, 492.80002, 492.80002	wwPDB
Map dimensions	448, 448, 448	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.1, 1.1, 1.1	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/3515	0.62	5/4788 (0.1%)
1	B	0.27	0/3515	0.60	2/4788 (0.0%)
1	C	0.27	0/3515	0.57	1/4788 (0.0%)
1	D	0.28	0/3463	0.59	1/4718 (0.0%)
1	E	0.34	0/3463	0.60	1/4718 (0.0%)
1	F	0.30	0/3463	0.58	1/4718 (0.0%)
All	All	0.29	0/20934	0.59	11/28518 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	1
All	All	0	6

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	461	LEU	CA-CB-CG	8.20	134.15	115.30
1	E	462	SER	O-C-N	-7.83	110.17	122.70
1	A	474	ILE	CG1-CB-CG2	-6.23	97.69	111.40
1	F	68	MET	CA-CB-CG	6.11	123.69	113.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	173	ILE	CG1-CB-CG2	-5.51	99.28	111.40

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	462	SER	Mainchain
1	B	462	SER	Mainchain
1	C	462	SER	Mainchain
1	D	462	SER	Mainchain
1	E	462	SER	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3464	0	3547	64	0
1	B	3464	0	3547	54	0
1	C	3464	0	3547	56	0
1	D	3412	0	3496	55	0
1	E	3412	0	3496	52	0
1	F	3412	0	3496	44	0
2	G	49	0	43	0	0
2	H	49	0	43	0	0
2	I	49	0	43	0	0
2	J	49	0	43	0	0
2	K	49	0	43	0	0
2	L	49	0	43	0	0
3	A	56	0	52	0	0
3	B	56	0	52	1	0
3	C	56	0	52	0	0
3	D	56	0	52	1	0
3	E	56	0	52	0	0
3	F	56	0	52	0	0
All	All	21258	0	21699	288	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 288 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:389:ALA:HB1	1:C:101:ASN:ND2	1.28	1.48
1:A:389:ALA:CB	1:C:101:ASN:HD21	1.32	1.43
1:A:389:ALA:CB	1:C:101:ASN:ND2	1.81	1.38
1:A:389:ALA:HB3	1:C:101:ASN:HD21	1.24	1.03
1:A:389:ALA:CA	1:C:101:ASN:HD21	1.85	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/541 (84%)	422 (93%)	30 (7%)	2 (0%)	30	68
1	B	454/541 (84%)	426 (94%)	26 (6%)	2 (0%)	30	68
1	C	454/541 (84%)	421 (93%)	32 (7%)	1 (0%)	44	78
1	D	447/541 (83%)	410 (92%)	37 (8%)	0	100	100
1	E	447/541 (83%)	416 (93%)	30 (7%)	1 (0%)	44	78
1	F	447/541 (83%)	419 (94%)	28 (6%)	0	100	100
All	All	2703/3246 (83%)	2514 (93%)	183 (7%)	6 (0%)	45	78

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	101	ASN
1	A	429	GLN
1	B	429	GLN
1	A	108	ILE
1	B	101	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	405/468 (86%)	405 (100%)	0	100	100
1	B	405/468 (86%)	403 (100%)	2 (0%)	86	89
1	C	405/468 (86%)	404 (100%)	1 (0%)	92	94
1	D	398/468 (85%)	398 (100%)	0	100	100
1	E	398/468 (85%)	392 (98%)	6 (2%)	60	75
1	F	398/468 (85%)	394 (99%)	4 (1%)	73	82
All	All	2409/2808 (86%)	2396 (100%)	13 (0%)	85	89

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	463	LYS
1	E	469	LYS
1	F	104	ILE
1	F	97	ASP
1	F	103	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	456	ASN
1	C	380	ASN
1	E	453	GLN
1	E	456	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	G	1	1,2	14,14,15	0.68	0	17,19,21	2.49	4 (23%)
2	NAG	G	2	2	14,14,15	0.73	0	17,19,21	1.74	3 (17%)
2	BMA	G	3	2	11,11,12	0.93	1 (9%)	15,15,17	2.63	5 (33%)
2	FUC	G	4	2	10,10,11	0.76	0	14,14,16	0.99	1 (7%)
2	NAG	H	1	1,2	14,14,15	0.67	0	17,19,21	2.63	4 (23%)
2	NAG	H	2	2	14,14,15	0.73	0	17,19,21	1.73	3 (17%)
2	BMA	H	3	2	11,11,12	0.92	1 (9%)	15,15,17	2.63	4 (26%)
2	FUC	H	4	2	10,10,11	0.74	0	14,14,16	1.00	1 (7%)
2	NAG	I	1	1,2	14,14,15	0.72	0	17,19,21	1.83	3 (17%)
2	NAG	I	2	2	14,14,15	0.71	0	17,19,21	1.78	3 (17%)
2	BMA	I	3	2	11,11,12	0.93	1 (9%)	15,15,17	2.65	5 (33%)
2	FUC	I	4	2	10,10,11	0.73	0	14,14,16	1.00	1 (7%)
2	NAG	J	1	1,2	14,14,15	0.71	0	17,19,21	2.44	5 (29%)
2	NAG	J	2	2	14,14,15	0.73	0	17,19,21	1.72	3 (17%)
2	BMA	J	3	2	11,11,12	0.94	1 (9%)	15,15,17	2.65	5 (33%)
2	FUC	J	4	2	10,10,11	0.73	0	14,14,16	1.01	1 (7%)
2	NAG	K	1	1,2	14,14,15	0.73	0	17,19,21	1.95	5 (29%)
2	NAG	K	2	2	14,14,15	0.72	0	17,19,21	1.75	3 (17%)
2	BMA	K	3	2	11,11,12	0.92	1 (9%)	15,15,17	2.70	5 (33%)
2	FUC	K	4	2	10,10,11	0.76	0	14,14,16	0.96	1 (7%)
2	NAG	L	1	1,2	14,14,15	0.67	0	17,19,21	2.39	4 (23%)
2	NAG	L	2	2	14,14,15	0.72	0	17,19,21	1.72	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	L	3	2	11,11,12	0.92	1 (9%)	15,15,17	2.65	4 (26%)
2	FUC	L	4	2	10,10,11	0.75	0	14,14,16	0.99	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	G	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
2	BMA	G	3	2	-	2/2/19/22	0/1/1/1
2	FUC	G	4	2	-	-	0/1/1/1
2	NAG	H	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	H	2	2	-	2/6/23/26	0/1/1/1
2	BMA	H	3	2	-	1/2/19/22	0/1/1/1
2	FUC	H	4	2	-	-	0/1/1/1
2	NAG	I	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	I	2	2	-	2/6/23/26	0/1/1/1
2	BMA	I	3	2	-	1/2/19/22	0/1/1/1
2	FUC	I	4	2	-	-	0/1/1/1
2	NAG	J	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	J	2	2	-	2/6/23/26	0/1/1/1
2	BMA	J	3	2	-	1/2/19/22	0/1/1/1
2	FUC	J	4	2	-	-	0/1/1/1
2	NAG	K	1	1,2	-	3/6/23/26	0/1/1/1
2	NAG	K	2	2	-	2/6/23/26	0/1/1/1
2	BMA	K	3	2	-	1/2/19/22	0/1/1/1
2	FUC	K	4	2	-	-	0/1/1/1
2	NAG	L	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	L	2	2	-	2/6/23/26	0/1/1/1
2	BMA	L	3	2	-	1/2/19/22	0/1/1/1
2	FUC	L	4	2	-	-	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	3	BMA	C2-C3	2.27	1.56	1.52
2	I	3	BMA	C2-C3	2.23	1.55	1.52
2	G	3	BMA	C2-C3	2.21	1.55	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	3	BMA	C2-C3	2.20	1.55	1.52
2	K	3	BMA	C2-C3	2.20	1.55	1.52

The worst 5 of 77 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	3	BMA	C1-O5-C5	8.60	123.72	112.19
2	I	3	BMA	C1-O5-C5	8.42	123.47	112.19
2	L	3	BMA	C1-O5-C5	8.40	123.45	112.19
2	J	3	BMA	C1-O5-C5	8.38	123.42	112.19
2	H	3	BMA	C1-O5-C5	8.30	123.31	112.19

There are no chirality outliers.



5 of 31 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	G	1	NAG	C3-C2-N2-C7
2	H	1	NAG	C3-C2-N2-C7
2	G	3	BMA	O5-C5-C6-O6
2	J	3	BMA	O5-C5-C6-O6
2	I	3	BMA	O5-C5-C6-O6



There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	601	1	14,14,15	0.72	0	17,19,21	0.96	1 (5%)
3	NAG	C	601	1	14,14,15	0.70	0	17,19,21	0.96	1 (5%)
3	NAG	C	604	1	14,14,15	0.79	0	17,19,21	0.92	0
3	NAG	C	602	1	14,14,15	0.70	0	17,19,21	1.30	2 (11%)
3	NAG	E	602	1	14,14,15	0.68	0	17,19,21	1.27	2 (11%)
3	NAG	F	601	1	14,14,15	0.70	0	17,19,21	0.96	1 (5%)
3	NAG	C	603	1	14,14,15	0.70	0	17,19,21	1.21	2 (11%)
3	NAG	F	604	1	14,14,15	0.74	0	17,19,21	1.29	1 (5%)
3	NAG	E	603	1	14,14,15	0.71	0	17,19,21	1.26	2 (11%)
3	NAG	D	602	1	14,14,15	0.70	0	17,19,21	1.17	2 (11%)
3	NAG	D	603	1	14,14,15	0.38	0	17,19,21	0.50	0
3	NAG	A	602	1	14,14,15	0.68	0	17,19,21	1.26	2 (11%)
3	NAG	B	604	1	14,14,15	0.72	0	17,19,21	1.22	1 (5%)
3	NAG	D	604	1	14,14,15	0.74	1 (7%)	17,19,21	2.61	4 (23%)
3	NAG	E	604	1	14,14,15	0.36	0	17,19,21	0.49	0
3	NAG	A	603	1	14,14,15	0.69	0	17,19,21	1.24	2 (11%)
3	NAG	B	603	1	14,14,15	0.72	0	17,19,21	1.21	1 (5%)
3	NAG	B	602	1	14,14,15	0.71	0	17,19,21	1.28	2 (11%)
3	NAG	B	601	1	14,14,15	0.71	0	17,19,21	0.94	1 (5%)
3	NAG	A	601	1	14,14,15	0.70	0	17,19,21	0.91	1 (5%)
3	NAG	E	601	1	14,14,15	0.70	0	17,19,21	0.91	0
3	NAG	F	603	1	14,14,15	0.70	0	17,19,21	1.21	2 (11%)
3	NAG	A	604	1	14,14,15	0.93	1 (7%)	17,19,21	2.70	5 (29%)
3	NAG	F	602	1	14,14,15	0.73	0	17,19,21	1.40	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	601	1	-	0/6/23/26	0/1/1/1
3	NAG	C	601	1	-	0/6/23/26	0/1/1/1
3	NAG	C	604	1	-	0/6/23/26	0/1/1/1
3	NAG	C	602	1	-	2/6/23/26	0/1/1/1
3	NAG	E	602	1	-	2/6/23/26	0/1/1/1
3	NAG	F	601	1	-	0/6/23/26	0/1/1/1
3	NAG	C	603	1	-	2/6/23/26	0/1/1/1
3	NAG	F	604	1	-	0/6/23/26	0/1/1/1
3	NAG	E	603	1	-	2/6/23/26	0/1/1/1
3	NAG	D	602	1	-	2/6/23/26	0/1/1/1
3	NAG	D	603	1	1/1/5/7	2/6/23/26	0/1/1/1
3	NAG	A	602	1	-	2/6/23/26	0/1/1/1
3	NAG	B	604	1	-	2/6/23/26	0/1/1/1
3	NAG	D	604	1	-	3/6/23/26	0/1/1/1
3	NAG	E	604	1	-	0/6/23/26	0/1/1/1
3	NAG	A	603	1	-	2/6/23/26	0/1/1/1
3	NAG	B	603	1	-	2/6/23/26	0/1/1/1
3	NAG	B	602	1	-	2/6/23/26	0/1/1/1
3	NAG	B	601	1	-	0/6/23/26	0/1/1/1
3	NAG	A	601	1	-	0/6/23/26	0/1/1/1
3	NAG	E	601	1	-	0/6/23/26	0/1/1/1
3	NAG	F	603	1	-	2/6/23/26	0/1/1/1
3	NAG	A	604	1	-	2/6/23/26	0/1/1/1
3	NAG	F	602	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	604	NAG	C1-C2	3.07	1.56	1.52
3	D	604	NAG	C1-C2	2.23	1.55	1.52

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	604	NAG	C1-O5-C5	8.65	123.78	112.19
3	D	604	NAG	C1-O5-C5	8.46	123.52	112.19
3	D	604	NAG	C2-N2-C7	4.76	129.28	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	604	NAG	C2-N2-C7	4.69	129.19	122.90
3	F	604	NAG	C1-O5-C5	3.97	117.51	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	D	603	NAG	C1

5 of 31 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	D	603	NAG	C8-C7-N2-C2
3	D	603	NAG	O7-C7-N2-C2
3	A	602	NAG	C8-C7-N2-C2
3	A	602	NAG	O7-C7-N2-C2
3	A	603	NAG	C8-C7-N2-C2



There are no ring outliers.



2 monomers are involved in 2 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	603	NAG	1	0
3	B	604	NAG	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-48535. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 224




Y Index: 224




Z Index: 224


6.2.2 Raw map



X Index: 224



Y Index: 224




Z Index: 224


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 232




Y Index: 227




Z Index: 252


6.3.2 Raw map



X Index: 224



Y Index: 227




Z Index: 253


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.847. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 356 nm^3 ; this corresponds to an approximate mass of 321 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.169 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.169 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.90	-	-
Author-provided FSC curve	7.84	9.17	8.03
Unmasked-calculated*	7.81	9.07	7.96


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from author-provided FSC intersecting FSC 0.143 CUT-OFF 7.84 differs from the reported value 5.9 by more than 10 %

The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.81 differs from the reported value 5.9 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-48535 and PDB model 9MQN. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.847 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.847).

9.4 Atom inclusion [i](#)



At the recommended contour level, 97% of all backbone atoms, 86% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.847) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.8600	<div></div> 0.1940
A	<div></div> 0.8770	<div></div> 0.1990
B	<div></div> 0.8550	<div></div> 0.1920
C	<div></div> 0.8790	<div></div> 0.1910
D	<div></div> 0.8700	<div></div> 0.1990
E	<div></div> 0.8590	<div></div> 0.1940
F	<div></div> 0.8760	<div></div> 0.1860
G	<div></div> 0.1430	<div></div> 0.2670
H	<div></div> 0.4900	<div></div> 0.2960
I	<div></div> 0.1020	<div></div> 0.2450
J	<div></div> 0.1220	<div></div> 0.1780
K	<div></div> 0.0200	<div></div> 0.1740
L	<div></div> 0.3060	<div></div> 0.2940

