



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 24, 2025 – 07:05 pm BST

PDB ID : 5MSJ / pdb_00005msj
Title : Mouse PA28alpha
Authors : Huber, E.M.; Groll, M.
Deposited on : 2017-01-05
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1094 (3.56-3.44)
Clashscore	180529	1045 (3.54-3.46)
Ramachandran outliers	177936	1032 (3.54-3.46)
Sidechain outliers	177891	1033 (3.54-3.46)
RSRZ outliers	164620	1093 (3.56-3.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	
1	B	249	
1	C	249	
1	D	249	
1	E	249	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	249	
1	G	249	
1	H	249	
1	I	249	
1	J	249	
1	K	249	
1	L	249	
1	M	249	
1	N	249	
1	O	249	
1	P	249	
1	Q	249	
1	R	249	
1	S	249	
1	T	249	
1	U	249	
1	V	249	
1	W	249	
1	X	249	
1	Y	249	
1	Z	249	
1	a	249	
1	b	249	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 46128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome activator complex subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1690	1081	288	315	6			
1	B	204	Total	C	N	O	S	0	0	0
			1646	1051	282	307	6			
1	C	205	Total	C	N	O	S	0	0	0
			1653	1056	283	308	6			
1	D	203	Total	C	N	O	S	0	0	0
			1635	1045	278	306	6			
1	E	208	Total	C	N	O	S	0	0	0
			1675	1070	286	313	6			
1	F	201	Total	C	N	O	S	0	0	0
			1619	1037	274	302	6			
1	G	205	Total	C	N	O	S	0	0	0
			1649	1054	280	309	6			
1	H	208	Total	C	N	O	S	0	0	0
			1676	1071	286	313	6			
1	I	195	Total	C	N	O	S	0	0	0
			1577	1006	269	297	5			
1	J	202	Total	C	N	O	S	0	0	0
			1635	1047	280	303	5			
1	K	207	Total	C	N	O	S	0	0	0
			1667	1066	285	310	6			
1	L	209	Total	C	N	O	S	0	0	0
			1682	1075	287	314	6			
1	M	201	Total	C	N	O	S	0	0	0
			1622	1038	276	302	6			
1	N	208	Total	C	N	O	S	0	0	0
			1675	1071	286	312	6			
1	O	203	Total	C	N	O	S	0	0	0
			1632	1044	276	306	6			
1	P	208	Total	C	N	O	S	0	0	0
			1675	1071	286	312	6			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	203	Total	C	N	O	S	0	0	0
			1635	1045	278	306	6			
1	R	200	Total	C	N	O	S	0	0	0
			1612	1032	273	301	6			
1	S	199	Total	C	N	O	S	0	0	0
			1602	1024	272	300	6			
1	T	206	Total	C	N	O	S	0	0	0
			1661	1062	284	309	6			
1	U	212	Total	C	N	O	S	0	0	0
			1711	1092	290	323	6			
1	V	202	Total	C	N	O	S	0	0	0
			1625	1039	275	305	6			
1	W	204	Total	C	N	O	S	0	0	0
			1647	1053	282	307	5			
1	X	200	Total	C	N	O	S	0	0	0
			1612	1031	273	303	5			
1	Y	201	Total	C	N	O	S	0	0	0
			1622	1037	276	304	5			
1	Z	204	Total	C	N	O	S	0	0	0
			1651	1057	282	307	5			
1	a	209	Total	C	N	O	S	0	0	0
			1683	1076	287	314	6			
1	b	206	Total	C	N	O	S	0	0	0
			1659	1062	284	307	6			

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome activator complex subunit 1

Chain A: 




- Molecule 1: Proteasome activator complex subunit 1

Chain B: 




- Molecule 1: Proteasome activator complex subunit 1


Chain C: 




- Molecule 1: Proteasome activator complex subunit 1

Chain D: 











- Molecule 1: Proteasome activator complex subunit 1

Chain T: 75% 7% 17%




- Molecule 1: Proteasome activator complex subunit 1

Chain U: 78% 7% 15%



- Molecule 1: Proteasome activator complex subunit 1

Chain V: 76% 5% 19%



- Molecule 1: Proteasome activator complex subunit 1

Chain W: 78% . 18%



- Molecule 1: Proteasome activator complex subunit 1

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	87.79Å 119.80Å 196.67Å 94.21° 98.52° 87.98°	Depositor
Resolution (Å)	29.65 – 3.50 29.65 – 3.50	Depositor EDS
% Data completeness (in resolution range)	93.1 (29.65-3.50) 93.1 (29.65-3.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 3.47Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.250 , 0.283 0.251 , 0.283	Depositor DCC
R_{free} test set	4638 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	91.5	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	46128	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/1722	0.85	0/2329
1	B	0.55	0/1675	0.86	0/2262
1	C	0.55	0/1683	0.86	0/2274
1	D	0.55	0/1665	0.85	0/2250
1	E	0.54	0/1705	0.85	0/2304
1	F	0.54	0/1649	0.87	0/2228
1	G	0.55	0/1679	0.83	0/2270
1	H	0.56	0/1707	0.85	0/2307
1	I	0.55	0/1604	0.86	0/2165
1	J	0.55	0/1664	0.85	0/2248
1	K	0.54	0/1698	0.84	0/2296
1	L	0.56	0/1715	0.85	0/2320
1	M	0.55	0/1652	0.84	0/2232
1	N	0.54	0/1706	0.83	0/2306
1	O	0.54	0/1662	0.84	0/2245
1	P	0.55	0/1706	0.86	0/2306
1	Q	0.55	0/1664	0.86	0/2248
1	R	0.55	0/1641	0.84	0/2217
1	S	0.54	0/1630	0.87	0/2200
1	T	0.54	0/1691	0.85	0/2285
1	U	0.54	0/1742	0.84	0/2352
1	V	0.55	0/1654	0.84	0/2234
1	W	0.54	0/1677	0.84	0/2266
1	X	0.53	0/1640	0.86	0/2214
1	Y	0.55	0/1651	0.84	0/2230
1	Z	0.54	0/1681	0.86	0/2271
1	a	0.55	0/1714	0.85	0/2318
1	b	0.54	0/1690	0.84	0/2284
All	All	0.55	0/46967	0.85	0/63461

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1690	0	1718	4	0
1	B	1646	0	1672	5	0
1	C	1653	0	1679	7	0
1	D	1635	0	1657	0	0
1	E	1675	0	1705	8	0
1	F	1619	0	1650	9	0
1	G	1649	0	1673	4	0
1	H	1676	0	1702	8	0
1	I	1577	0	1594	6	0
1	J	1635	0	1669	6	0
1	K	1667	0	1695	4	0
1	L	1682	0	1705	13	0
1	M	1622	0	1649	3	0
1	N	1675	0	1705	6	0
1	O	1632	0	1659	7	0
1	P	1675	0	1705	5	0
1	Q	1635	0	1659	7	0
1	R	1612	0	1642	8	0
1	S	1602	0	1629	6	0
1	T	1661	0	1690	10	0
1	U	1711	0	1734	11	0
1	V	1625	0	1651	8	0
1	W	1647	0	1674	6	0
1	X	1612	0	1639	6	0
1	Y	1622	0	1645	4	0
1	Z	1651	0	1683	4	0
1	a	1683	0	1710	3	0
1	b	1659	0	1692	7	0
All	All	46128	0	46885	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:124:VAL:HG21	1:H:219:ILE:HG23	1.63	0.78
1:E:135:LEU:HD11	1:E:237:LEU:HD22	1.66	0.78
1:F:124:VAL:HG21	1:F:219:ILE:HG23	1.69	0.75
1:L:124:VAL:HG21	1:L:219:ILE:HG23	1.69	0.73
1:U:135:LEU:HD11	1:U:237:LEU:HD22	1.76	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/249 (83%)	201 (98%)	4 (2%)	1 (0%)	25	59
1	B	200/249 (80%)	194 (97%)	6 (3%)	0	100	100
1	C	201/249 (81%)	197 (98%)	3 (2%)	1 (0%)	25	59
1	D	199/249 (80%)	195 (98%)	3 (2%)	1 (0%)	25	59
1	E	204/249 (82%)	199 (98%)	3 (2%)	2 (1%)	13	46
1	F	197/249 (79%)	190 (96%)	7 (4%)	0	100	100
1	G	201/249 (81%)	199 (99%)	1 (0%)	1 (0%)	25	59
1	H	204/249 (82%)	199 (98%)	3 (2%)	2 (1%)	13	46
1	I	191/249 (77%)	189 (99%)	2 (1%)	0	100	100
1	J	198/249 (80%)	194 (98%)	4 (2%)	0	100	100
1	K	203/249 (82%)	200 (98%)	3 (2%)	0	100	100
1	L	205/249 (82%)	200 (98%)	5 (2%)	0	100	100
1	M	197/249 (79%)	194 (98%)	3 (2%)	0	100	100
1	N	204/249 (82%)	201 (98%)	3 (2%)	0	100	100
1	O	199/249 (80%)	195 (98%)	3 (2%)	1 (0%)	25	59
1	P	204/249 (82%)	200 (98%)	3 (2%)	1 (0%)	25	59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	199/249 (80%)	195 (98%)	3 (2%)	1 (0%)	25	59
1	R	196/249 (79%)	192 (98%)	4 (2%)	0	100	100
1	S	195/249 (78%)	190 (97%)	5 (3%)	0	100	100
1	T	202/249 (81%)	198 (98%)	4 (2%)	0	100	100
1	U	208/249 (84%)	203 (98%)	4 (2%)	1 (0%)	25	59
1	V	198/249 (80%)	193 (98%)	5 (2%)	0	100	100
1	W	200/249 (80%)	196 (98%)	4 (2%)	0	100	100
1	X	196/249 (79%)	192 (98%)	2 (1%)	2 (1%)	13	46
1	Y	197/249 (79%)	192 (98%)	4 (2%)	1 (0%)	25	59
1	Z	200/249 (80%)	199 (100%)	1 (0%)	0	100	100
1	a	205/249 (82%)	196 (96%)	8 (4%)	1 (0%)	25	59
1	b	202/249 (81%)	198 (98%)	3 (2%)	1 (0%)	25	59
All	All	5611/6972 (80%)	5491 (98%)	103 (2%)	17 (0%)	37	68

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	146	ASN
1	D	58	LYS
1	H	58	LYS
1	Y	147	ASN
1	b	146	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/223 (84%)	182 (97%)	6 (3%)	34	62
1	B	182/223 (82%)	178 (98%)	4 (2%)	47	70
1	C	183/223 (82%)	179 (98%)	4 (2%)	47	70
1	D	181/223 (81%)	177 (98%)	4 (2%)	47	70

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	186/223 (83%)	181 (97%)	5 (3%)	40	65
1	F	180/223 (81%)	177 (98%)	3 (2%)	56	75
1	G	183/223 (82%)	181 (99%)	2 (1%)	70	83
1	H	186/223 (83%)	182 (98%)	4 (2%)	47	70
1	I	174/223 (78%)	170 (98%)	4 (2%)	45	69
1	J	181/223 (81%)	174 (96%)	7 (4%)	27	56
1	K	185/223 (83%)	181 (98%)	4 (2%)	47	70
1	L	187/223 (84%)	182 (97%)	5 (3%)	40	65
1	M	180/223 (81%)	174 (97%)	6 (3%)	33	61
1	N	186/223 (83%)	182 (98%)	4 (2%)	47	70
1	O	181/223 (81%)	175 (97%)	6 (3%)	33	61
1	P	186/223 (83%)	182 (98%)	4 (2%)	47	70
1	Q	181/223 (81%)	178 (98%)	3 (2%)	56	75
1	R	179/223 (80%)	173 (97%)	6 (3%)	32	60
1	S	177/223 (79%)	172 (97%)	5 (3%)	38	65
1	T	184/223 (82%)	177 (96%)	7 (4%)	28	57
1	U	190/223 (85%)	188 (99%)	2 (1%)	70	83
1	V	180/223 (81%)	176 (98%)	4 (2%)	47	70
1	W	182/223 (82%)	180 (99%)	2 (1%)	70	83
1	X	178/223 (80%)	178 (100%)	0	100	100
1	Y	179/223 (80%)	176 (98%)	3 (2%)	56	75
1	Z	183/223 (82%)	178 (97%)	5 (3%)	40	65
1	a	187/223 (84%)	186 (100%)	1 (0%)	86	93
1	b	184/223 (82%)	181 (98%)	3 (2%)	58	76
All	All	5113/6244 (82%)	5000 (98%)	113 (2%)	47	70

5 of 113 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	159	LEU
1	b	4	LEU
1	P	148	PHE
1	a	173	GLN
1	W	13	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 94 such sidechains are listed below:

Mol	Chain	Res	Type
1	N	147	ASN
1	T	147	ASN
1	O	146	ASN
1	Q	146	ASN
1	U	191	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/249 (84%)	-0.16	1 (0%) 87 75	66, 89, 122, 152	0
1	B	204/249 (81%)	-0.15	1 (0%) 87 75	70, 95, 136, 141	0
1	C	205/249 (82%)	-0.13	0 100 100	71, 97, 133, 149	0
1	D	203/249 (81%)	-0.08	1 (0%) 87 75	76, 100, 148, 160	0
1	E	208/249 (83%)	0.00	1 (0%) 87 75	82, 105, 135, 148	0
1	F	201/249 (80%)	-0.15	0 100 100	73, 95, 137, 152	0
1	G	205/249 (82%)	-0.19	1 (0%) 87 75	71, 88, 122, 138	0
1	H	208/249 (83%)	-0.14	0 100 100	79, 100, 134, 162	0
1	I	195/249 (78%)	-0.07	0 100 100	79, 105, 143, 158	0
1	J	202/249 (81%)	-0.15	0 100 100	75, 98, 143, 154	0
1	K	207/249 (83%)	-0.14	0 100 100	75, 101, 144, 157	0
1	L	209/249 (83%)	-0.13	1 (0%) 87 75	76, 106, 137, 165	0
1	M	201/249 (80%)	-0.11	0 100 100	75, 107, 145, 157	0
1	N	208/249 (83%)	-0.14	0 100 100	73, 98, 137, 147	0
1	O	203/249 (81%)	-0.12	1 (0%) 87 75	71, 93, 125, 147	0
1	P	208/249 (83%)	-0.18	0 100 100	72, 95, 132, 144	0
1	Q	203/249 (81%)	-0.21	0 100 100	76, 100, 129, 135	0
1	R	200/249 (80%)	-0.16	0 100 100	81, 100, 140, 171	0
1	S	199/249 (79%)	-0.15	0 100 100	69, 94, 137, 159	0
1	T	206/249 (82%)	-0.08	0 100 100	70, 93, 131, 139	0
1	U	212/249 (85%)	-0.19	1 (0%) 87 75	69, 92, 144, 174	0
1	V	202/249 (81%)	-0.15	0 100 100	76, 95, 134, 161	0
1	W	204/249 (81%)	-0.20	0 100 100	73, 95, 137, 150	0
1	X	200/249 (80%)	-0.19	0 100 100	69, 94, 128, 136	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9	
1	Y	201/249 (80%)	-0.17	0	100	100	67, 96, 136, 162	0
1	Z	204/249 (81%)	-0.19	0	100	100	68, 95, 134, 159	0
1	a	209/249 (83%)	-0.11	0	100	100	71, 98, 133, 155	0
1	b	206/249 (82%)	-0.15	1 (0%)	87	75	71, 97, 133, 148	0
All	All	5723/6972 (82%)	-0.14	9 (0%)	92	86	66, 97, 137, 174	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	O	145	GLY	3.8
1	b	62	ASP	2.4
1	L	144	ASP	2.3
1	B	145	GLY	2.2
1	D	171	HIS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.